



OPEN ACCESS

APPROVED BY

Frontiers Editorial Office,
Frontiers Media SA, Switzerland

*CORRESPONDENCE

Young Woo Park
park1556@umn.edu
Christophe Lenglet
clenglet@umn.edu

SPECIALTY SECTION

This article was submitted to
Applied Neuroimaging,
a section of the journal
Frontiers in Neurology

RECEIVED 06 September 2022

ACCEPTED 07 September 2022

PUBLISHED 29 September 2022

CITATION

Park YW, Joers JM, Guo B, Hutter D, Bushara K, Adanyeguh IM, Eberly LE, Öz G and Lenglet C (2022) Corrigendum: Assessment of cerebral and cerebellar white matter microstructure in spinocerebellar ataxias 1, 2, 3, and 6 using diffusion MRI. *Front. Neurosci.* 13:1038298. doi: 10.3389/fneur.2022.1038298

COPYRIGHT

© 2022 Park, Joers, Guo, Hutter, Bushara, Adanyeguh, Eberly, Öz and Lenglet. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: Assessment of cerebral and cerebellar white matter microstructure in spinocerebellar ataxias 1, 2, 3, and 6 using diffusion MRI

Young Woo Park^{1*}, James M. Joers¹, Bin Guo², Diane Hutter¹, Khalaf Bushara³, Isaac M. Adanyeguh¹, Lynn E. Eberly^{1,2}, Gülin Öz¹ and Christophe Lenglet^{1*}

¹Department of Radiology, Center for Magnetic Resonance Research, University of Minnesota Medical School, Minneapolis, MN, United States, ²Division of Biostatistics, School of Public Health, University of Minnesota, Minneapolis, MN, United States, ³Department of Neurology, University of Minnesota Medical School, Minneapolis, MN, United States

KEYWORDS

SCA1, SCA2, SCA3, SCA6, diffusion MRI, Spinocerebellar ataxias

A corrigendum on

Assessment of cerebral and cerebellar white matter microstructure in spinocerebellar ataxias 1, 2, 3, and 6 using diffusion MRI

by Park, Y. W., Joers, J. M., Guo, B., Hutter, D., Bushara, K., Adanyeguh, I. M., Eberly, L. E., Öz, G., and Lenglet, C. (2020). *Front. Neurosci.* 11:411. doi: 10.3389/fneur.2020.00411

In the published article there was an error in the reference list as published. The reference list was submitted in the incorrect order. The revised reference list appears below.

The authors apologize for the above-mentioned errors and state that it does not affect the conclusions of the article in any way. The original version of this article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Ruano L, Melo C, Silva MC, Coutinho P. The global epidemiology of hereditary ataxia and spastic paraparesis: a systematic review of prevalence studies. *Neuroepidemiology*. (2014) 42:174–83. doi: 10.1159/000358801
- Sullivan R, Yau WY, O'Connor E, Houlden H. Spinocerebellar ataxia: an update. *J Neurol.* (2019) 266:533–44. doi: 10.1007/s00415-018-9076-4
- Schols L, Amoiridis G, Buttner T, Przuntek H, Epplen JT, Riess O. Autosomal dominant cerebellar ataxia: phenotypic differences in genetically defined subtypes? *Ann Neurol.* (1997) 42:924–32. doi: 10.1002/ana.410420615
- Seidel K, Siswanto S, Brunt ER, den Dunnen W, Korf HW, Rub U. Brain pathology of spinocerebellar ataxias. *Acta Neuropathol.* (2012) 124:1–21. doi: 10.1007/s00401-012-1000-x
- Rub U, Schols L, Paulson H, Auburger G, Kermer P, Jen JC, et al. Clinical features, neurogenetics and neuropathology of the polyglutamine spinocerebellar ataxias type 1, 2, 3, 6 and 7. *Prog Neurobiol.* (2013) 104:38–66. doi: 10.1016/j.pneurobio.2013.01.001
- Orr HT, Chung MY, Banfi S, Kwiatkowski TJ Jr, Servadio A, Beaudet AL, et al. Expansion of an unstable trinucleotide CAG repeat in spinocerebellar ataxia type 1. *Nat Genet.* (1993) 4:221–6. doi: 10.1038/ng0793-221
- Rub U, Burk K, Timmann D, den Dunnen W, Seidel K, Farrag K, et al. Spinocerebellar ataxia type 1 (SCA1): new pathoanatomical and clinicopathological insights. *Neuropathol Appl Neurobiol.* (2012) 38:665–80. doi: 10.1111/j.1365-2990.2012.01259.x
- Armstrong J, Bonaventura I, Rojo A, Gonzalez G, Corral J, Nadal N, et al. Spinocerebellar ataxia type 2 (SCA2) with white matter involvement. *Neurosci Lett.* (2005) 381:247–51. doi: 10.1016/j.neulet.2005.02.063
- Riess O, Rub U, Pastore A, Bauer P, Schols L. SCA3: neurological features, pathogenesis and animal models. *Cerebellum.* (2008) 7:125–37. doi: 10.1007/s12311-008-0013-4
- Ishikawa K, Watanabe M, Yoshizawa K, Fujita T, Iwamoto H, Yoshizawa T, et al. Clinical, neuropathological, and molecular study in two families with spinocerebellar ataxia type 6 (SCA6). *J Neurol Neurosurg Psychiatry*. (1999) 67:86–9.
- Ginestroni A, Della Nave R, Tessa C, Giannelli M, De Grandis D, Plasmati R, et al. Brain structural damage in spinocerebellar ataxia type 1: a VBM study. *J Neurol.* (2008) 255:1153–8. doi: 10.1007/s00415-008-0860-4
- Jung BC, Choi SI, Du AX, Cuzzocreo JL, Ying HS, Landman BA, et al. MRI shows a region-specific pattern of atrophy in spinocerebellar ataxia type 2. *Cerebellum.* (2012) 11:272–9. doi: 10.1007/s12311-011-0308-8
- Rezende TJR, de Paiva JLR, Martinez ARM, Lopes-Cendes I, Pedroso JL, Barsottini OGP, et al. Structural signature of SCA3: from presymptomatic to late disease stages. *Ann Neurol.* (2018) 84:401–8. doi: 10.1002/ana.25297
- Reetz K, Costa AS, Mirzazade S, Lehmann A, Juzek A, Rakowicz M, et al. Genotype-specific patterns of atrophy progression are more sensitive than clinical decline in SCA1, SCA3 and SCA6. *Brain.* (2013) 136:905–17. doi: 10.1093/brain/aws369
- Oz G, Iltis I, Hutter D, Thomas W, Bushara KO, Gomez CM. Distinct neurochemical profiles of spinocerebellar ataxias 1, 2, 6, and cerebellar multiple system atrophy. *Cerebellum.* (2011) 10:208–17. doi: 10.1007/s12311-010-0213-6
- Adanyeguh IM, Henry PG, Nguyen TM, Rinaldi D, Jauffret C, Valabregue R, et al. In vivo neurometabolic profiling in patients with spinocerebellar ataxia types 1, 2, 3, and 7. *Movement Disord.* (2015) 30:662–70. doi: 10.1002/mds.26181
- Joers JM, Deelchand DK, Lyu T, Emir UE, Hutter D, Gomez CM, et al. Neurochemical abnormalities in premanifest and early spinocerebellar ataxias. *Ann Neurol.* (2018) 83:816–29. doi: 10.1002/ana.25212
- Martins CR, Martinez ARM, Vasconcelos IF, de Rezende TJR, Casseb RF, Pedroso JL, et al. Structural signature in SCA1: clinical correlates, determinants and natural history. *J Neurol.* (2018) 265:2949–59. doi: 10.1007/s00415-018-9087-1
- Deelchand DK, Joers JM, Ravishankar A, Lyu T, Emir UE, Hutter D, et al. Sensitivity of volumetric magnetic resonance imaging and magnetic resonance spectroscopy to progression of spinocerebellar ataxia type 1. *Mov Disord Clin Pract.* (2019) 6:549–58. doi: 10.1002/mdc3.12804
- Mandelli ML, De Simone T, Minati L, Bruzzone MG, Mariotti C, Fancello R, et al. Diffusion tensor imaging of spinocerebellar ataxias types 1 and 2. *AJR Am J Neuroradiol.* (2007) 28:1996–2000. doi: 10.3174/ajnr.A0716
- Della Nave R, Ginestroni A, Tessa C, Salvatore E, De Grandis D, Plasmati R, et al. Brain white matter damage in SCA1 and SCA2. An in vivo study using voxel-based morphometry, histogram analysis of mean diffusivity and tract-based spatial statistics. *Neuroimage*. (2008) 43:10–9. doi: 10.1016/j.neuroimage.2008.06.036
- Prakash N, Hageman N, Hua X, Toga AW, Perlman SL, Salamon N. Patterns of fractional anisotropy changes in white matter of cerebellar peduncles distinguish spinocerebellar ataxia-1 from multiple system atrophy and other ataxia syndromes. *Neuroimage*. (2009) 47 (Suppl. 2):T72–81. doi: 10.1016/j.neuroimage.2009.05.013
- Alcauter S, Barrios FA, Diaz R, Fernandez-Ruiz J. Gray and white matter alterations in spinocerebellar ataxia type 7: an in vivo DTI and VBM study. *Neuroimage*. (2011) 55:1–7. doi: 10.1016/j.neuroimage.2010.12.014
- Kang JS, Klein JC, Baudrexel S, Deichmann R, Nolte D, Hilker R. White matter damage is related to ataxia severity in SCA3. *J Neurol.* (2014) 261:291–9. doi: 10.1007/s00415-013-7186-6
- Adanyeguh IM, Perlberg V, Henry PG, Rinaldi D, Petit E, Valabregue R, et al. Autosomal dominant cerebellar ataxias: imaging biomarkers with high effect sizes. *Neuroimage Clin.* (2018) 19:858–67. doi: 10.1016/j.nicl.2018.06.011
- Froeling M, Pullens P, Leemans A. DTI analysis methods: region of interest analysis. In: Van Hecke W, Emsell L, and Sunaert S, editors. *Diffusion Tensor Imaging: A Practical Handbook*. New York, NY: Springer New York (2016). p. 175–82. doi: 10.1007/978-1-4939-3118-7_9
- Falcon MI, Gomez CM, Chen EE, Shereen A, Solodkin A. Early cerebellar network shifting in spinocerebellar ataxia type 6. *Cereb Cortex*. (2016) 26:3205–18. doi: 10.1093/cercor/bhv154
- Smith SM, Jenkinson M, Johansen-Berg H, Rueckert D, Nichols TE, Mackay CE, et al. Tract-based spatial statistics: voxelwise analysis of multi-subject diffusion data. *Neuroimage*. (2006) 31:1487–505. doi: 10.1016/j.neuroimage.2006.02.024
- Tournier JD, Smith R, Raffelt D, Tabbara R, Dhollander T, Pietsch M, et al. MRtrix3: a fast, flexible and open software framework for medical image processing and visualisation. *Neuroimage*. (2019) 202:116137. doi: 10.1016/j.neuroimage.2019.116137
- Mascalchi M, Toschi N, Giannelli M, Ginestroni A, Della Nave R, Nicolai E, et al. Progression of microstructural damage in spinocerebellar ataxia type 2: a longitudinal DTI study. *AJNR Am J Neuroradiol.* (2015) 36:1096–01. doi: 10.3174/ajnr.A4343
- Mascalchi M, Marzi C, Giannelli M, Ciulli S, Bianchi A, Ginestroni A, et al. Histogram analysis of DTI-derived indices reveals pontocerebellar degeneration and its progression in SCA2. *PLoS ONE*. (2018) 13:200258. doi: 10.1371/journal.pone.0200258
- Schmitz-Hubsch T, du Montcel ST, Baliko L, Berciano J, Boesch S, Depont D, et al. Scale for the assessment and rating of ataxia: development of a new clinical scale. *Neurology*. (2006) 66:1717–20. doi: 10.1212/01.wnl.0000219042.60538.92
- Maas RP, van Gaalen J, Klockgether T, van de Warrenburg BP. The preclinical stage of spinocerebellar ataxias. *Neurology*. (2015) 85:96–103. doi: 10.1212/WNL.0000000000001711
- du Montcel ST, Durr A, Rakowicz M, Nanetti L, Charles P, Sulek A, et al. Prediction of the age at onset in spinocerebellar ataxia type 1, 2, 3 and 6. *J Med Genet*. (2014) 51:479–86. doi: 10.1136/jmedgenet-2013-102200
- Subramony SH, May W, Lynch D, Gomez C, Fischbeck K, Hallett M, et al. Measuring Friedreich ataxia: interrater reliability of a neurologic rating scale. *Neurology*. (2005) 64:1261–2. doi: 10.1212/01.WNL.0000156802.15466.79
- Patel M, Isaacs CJ, Seyer L, Brigatti K, Gelbard S, Strawser C, et al. Progression of Friedreich ataxia: quantitative characterization over 5 years. *Ann Clin Transl Neurol.* (2016) 3:684–94. doi: 10.1002/acn3.332
- Jenkinson M, Beckmann CF, Behrens TE, Woolrich MW, Smith SM. Fsl. *Neuroimage*. (2012) 62:782–90. doi: 10.1016/j.neuroimage.2011.09.015
- Hua K, Zhang J, Wakana S, Jiang H, Li X, Reich DS, et al. Tract probability maps in stereotaxic spaces: analyses of white matter anatomy and tract-specific quantification. *Neuroimage*. (2008) 39:336–47. doi: 10.1016/j.neuroimage.2007.07.053
- Keihaninejad S, Zhang H, Ryan NS, Malone IB, Modat M, Cardoso MJ, et al. An unbiased longitudinal analysis framework for tracking white matter changes using diffusion tensor imaging with application to Alzheimer's disease. *Neuroimage*. (2013) 72:153–63. doi: 10.1016/j.neuroimage.2013.01.044
- Zhang H, Avants BB, Yushkevich PA, Woo JH, Wang S, McCluskey LF, et al. High-dimensional spatial normalization of diffusion tensor images improves the detection of white matter differences: an example study using amyotrophic lateral sclerosis. *IEEE Trans Med Imaging*. (2007) 26:1585–97. doi: 10.1109/TMI.2007.906784
- Park YW, Joers JM, Hutter D, Bushara KO, Oz G, Lenglet C. Improved sensitivity to longitudinal changes with

- advanced DTI analysis in a rare neurodegenerative disease. In: ISMRM 27th Annual Meeting and Exhibition. Montreal, QC (2019).
42. Raffelt DA, Tournier JD, Smith RE, Vaughan DN, Jackson G, Ridgway GR, et al. Investigating white matter fibre density and morphology using fixel-based analysis. *Neuroimage*. (2017) 144 (Pt A):58–73. doi: 10.1016/j.neuroimage.2016.09.029
43. Schulz JB, Borkert J, Wolf S, Schmitz-Hubsch T, Rakowicz M, Mariotti C, et al. Visualization, quantification and correlation of brain atrophy with clinical symptoms in spinocerebellar ataxia types 1, 3 and 6. *Neuroimage*. (2010) 49:158–68. doi: 10.1016/j.neuroimage.2009.07.027
44. Diedrichsen J. A spatially unbiased atlas template of the human cerebellum. *Neuroimage*. (2006) 33:127–38. doi: 10.1016/j.neuroimage.2006.05.056
45. Romero JE, Coupe P, Giraud R, Ta VT, Fonov V, Park MTM, et al. CERES: a new cerebellum lobule segmentation method. *Neuroimage*. (2017) 147:916–24. doi: 10.1016/j.neuroimage.2016.11.003
46. Kinnunen KM, Greenwood R, Powell JH, Leech R, Hawkins PC, Bonnelle V, et al. White matter damage and cognitive impairment after traumatic brain injury. *Brain*. (2011) 134 (Pt 2):449–63. doi: 10.1093/brain/awq347
47. Della Nave R, Ginestroni A, Diciotti S, Salvatore E, Soricelli A, Mascalchi M. Axial diffusivity is increased in the degenerating superior cerebellar peduncles of Friedreich's ataxia. *Neuroradiology*. (2011) 53:367–72. doi: 10.1007/s00234-010-0807-1