Check for updates

OPEN ACCESS

EDITED AND REVIEWED BY Edoardo Malfatti, Hôpitaux Universitaires Henri Mondor, France

*CORRESPONDENCE Barry J. Byrne Barry.byrne@ufl.edu

RECEIVED 05 December 2024 ACCEPTED 12 December 2024 PUBLISHED 03 January 2025

CITATION

Byrne BJ, Parenti G, Schoser B, van der Ploeg AT, Do H, Fox B, Goldman M, Johnson FK, Kang J, Mehta N, Mondick J, Sheikh MO, Sitaraman Das S, Tuske S, Brudvig J, Weimer JM and Mozaffar T (2025) Corrigendum: Cipaglucosidase alfa plus miglustat: linking mechanism of action to clinical outcomes in late-onset Pompe disease. *Front. Neurol.* 15:1540452. doi: 10.3389/fneur.2024.1540452

COPYRIGHT

© 2025 Byrne, Parenti, Schoser, van der Ploeg, Do, Fox, Goldman, Johnson, Kang, Mehta, Mondick, Sheikh, Sitaraman Das, Tuske, Brudvig, Weimer and Mozaffar. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: Cipaglucosidase alfa plus miglustat: linking mechanism of action to clinical outcomes in late-onset Pompe disease

Barry J. Byrne^{1*}, Giancarlo Parenti^{2,3}, Benedikt Schoser⁴, Ans T. van der Ploeg⁵, Hung Do⁶, Brian Fox⁷, Mitchell Goldman⁷, Franklin K. Johnson⁷, Jia Kang⁸, Nickita Mehta⁷, John Mondick⁹, M. Osman Sheikh⁷, Sheela Sitaraman Das⁷, Steven Tuske⁷, Jon Brudvig⁷, Jill M. Weimer⁷ and Tahseen Mozaffar¹⁰

¹Department of Pediatrics in the College of Medicine, University of Florida, Gainesville, FL, United States, ²Metabolic Unit, Department of Translational Medical Sciences, University of Naples Federico II, Naples, Italy, ³Telethon Institute of Genetics and Medicine, Pozzuoli, Italy, ⁴Friedrich-Baur-Institute, Department of Neurology, LMU University Hospital, LMU Munich, Munich, Germany, ⁵Erasmus MC University Medical Center, Rotterdam, Netherlands, ⁶M6P-Therapeutics, St Louis, MO, United States, ⁷Amicus Therapeutics, Inc., Princeton, NJ, United States, ⁸Metrum Research Group, Tariffville, CT, United States, ⁹Incyte Corporation, Wilmington, DE, United States, ¹⁰Department of Neurology, University of California, Irvine, Irvine, CA, United States

KEYWORDS

Pompe disease, glycogen storage disease type II, lysosomal storage disorders, enzyme replacement therapy, *n*-butyldeoxynojirimycin

A Corrigendum on

Cipaglucosidase alfa plus miglustat: linking mechanism of action to clinical outcomes in late-onset Pompe disease

by Byrne, B. J., Parenti, G., Schoser, B., van der Ploeg, A. T., Do, H., Fox, B., Goldman, M., Johnson, F. K., Kang, J., Mehta, N., Mondick, J., Sheikh, M. O., Sitaraman Das, S., Tuske, S., Brudvig, J., Weimer, J. M., and Mozaffar. T. (2024). *Front. Neurol.* 15:1451512. doi: 10.3389/fneur.2024.1451512

In the published article, there was an error in Figure 13 as published. The error relates to the positive/negative value of some of the numbers on the axes for the lower MMT score and the GSGC total score in Figure 13A. For both lower MMT score and GSGC total score, the axis scale incorrectly read -3, -2, 1, 0, -1, -2, -3 from left to right, whereas the axis scale for lower MMT score should have been -3, -2, -1, 0, 1, 2, 3 and the axis scale for GSGC total score should have been 3, 2, 1, 0, -1, -2, -3. The corrected Figure 13 and its caption appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.



В

Endpoint	Overall patients (ERT experienced plus ERT naïve)				
	Cipaglucosidase alfa + miglustat		Alglucosidase alfa + placebo		Two-sided
	Baseline, mean	CFBL week 52 LOCF, mean (95% Cl)	Baseline, mean	CFBL week 52 LOCF, mean (95% Cl)	<i>p</i> value for LS mean difference
Hex4, mmol/mol creatine	4.61	-1.88 (-2.4, -1.36)	6.92	1.22 (-0.26, 2.70)	<0.001
CK, U/L	447.0	-130.5 (-180.4, -80.7)	527.8	60.2 (7.0, 113.3)	<0.001
Lower MMT, score	27.96	1.56 (0.72, 2.40)	27.65	0.88 (-0.02, 1.78)	0.191
6MWD, m	357.93	20.79 (11.56, 30.01)	350.95	7.24 (-6.19, 20.67)	0.071
GSGC, total score	14.51	-0.53 (-1.13, 0.06)	14.50	0.77 (0.09, 1.44)	0.009
FVC, % predicted	70.74	-0.93 (-2.29, 0.42)	69.68	-3.95 (-5.58, -2.32)	0.023

FIGURE 13

Change from baseline at week 52 of PROPEL—effect of cipaglucosidase alfa plus miglustat compared with alglucosidase alfa plus placebo in key efficacy outcomes. (A) Forest plot illustrating mean estimated treatment differences between cipaglucosidase alfa plus miglustat versus alglucosidase alfa plus placebo and corresponding 95% CIs are shown for the combined PROPEL study population for each outcome, with units as indicated on the x-axes. For all outcomes, right-sided directionality of treatment differences indicates favorable outcomes for cipaglucosidase alfa plus miglustat compared with alglucosidase alfa plus placebo. (B) The table shows baseline mean values and Week 52 CFBL values for cipaglucosidase alfa plus miglustat and alglucosidase alfa plus placebo. Shaded CFBL indicates nominally significant improvement (green) or nominally significant worsening (red) from baseline (i.e., the 95% CI does not include zero) within each treatment group. The *p*-values (two-tailed LS mean difference) shown in the far-right column are for the between-group treatment differences illustrated in the forest plot.