



# Angiotensin II-Type I Receptor Antagonism Does Not Influence the Chemoreceptor Reflex or Hypoxia-Induced Central Sleep Apnea in Men

Courtney V. Brown<sup>1</sup>, Lindsey M. Boulet<sup>1</sup>, Tyler D. Vermeulen<sup>1</sup>, Scott A. Sands<sup>2</sup>, Richard J. A. Wilson<sup>3</sup>, Najib T. Ayas<sup>4,5</sup>, John S. Floras<sup>6</sup> and Glen E. Foster<sup>1\*</sup>

<sup>1</sup> Centre for Heart, Lung, and Vascular Health, School of Health and Exercise Science, University of British Columbia – Okanagan, Kelowna, BC, Canada, <sup>2</sup> Division of Sleep and Circadian Disorders, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, United States, <sup>3</sup> Department of Physiology and Pharmacology, Hotchkiss Brain Institute, Cumming School of Medicine, University of Calgary, Calgary, AB, Canada, <sup>4</sup> Sleep Disorders Program, University of British Columbia, Vancouver, BC, Canada, <sup>5</sup> Respiratory and Critical Care Divisions, University of British Columbia, Vancouver, BC, Canada, <sup>6</sup> University Health Network and Sinai Health System Division of Cardiology, Department of Medicine, University of Toronto, Toronto, ON, Canada

## OPEN ACCESS

### Edited by:

Bernhard Schaller,  
University of Zurich, Switzerland

### Reviewed by:

Vincent Joseph,  
Laval University, Canada  
Noah J. Marcus,  
Des Moines University, United States

### \*Correspondence:

Glen E. Foster  
glen.foster@ubc.ca

### Specialty section:

This article was submitted to  
Autonomic Neuroscience,  
a section of the journal  
Frontiers in Neuroscience

Received: 27 January 2020

Accepted: 27 March 2020

Published: 28 April 2020

### Citation:

Brown CV, Boulet LM, Vermeulen TD, Sands SA, Wilson RJA, Ayas NT, Floras JS and Foster GE (2020) Angiotensin II-Type I Receptor Antagonism Does Not Influence the Chemoreceptor Reflex or Hypoxia-Induced Central Sleep Apnea in Men. *Front. Neurosci.* 14:382. doi: 10.3389/fnins.2020.00382

Components of the renin-angiotensin system (RAS) situated within the carotid body or central nervous system may promote hypoxia-induced chemoreceptor reflex sensitization or central sleep apnea (CSA). We determined if losartan, an angiotensin-II type-I receptor (AT<sub>1</sub>R) antagonist, would attenuate chemoreceptor reflex sensitivity before or after 8 h of nocturnal hypoxia, and consequently CSA severity. In a double-blind, randomized, placebo-controlled, crossover protocol, 14 men (age: 25 ± 2 years; BMI: 24.6 ± 1.1 kg/m<sup>2</sup>; means ± SEM) ingested 3 doses of either losartan (50 mg) or placebo every 8 h. Chemoreceptor reflex sensitivity was assessed during hypoxic and hyperoxic hypercapnic ventilatory response (HCVR) tests and during six-20s hypoxic apneas before and after 8 h of sleep in normobaric hypoxia (F<sub>I</sub>O<sub>2</sub> = 0.135). Loop gain was assessed from a ventilatory control model fitted to the ventilatory pattern of CSA recorded during polysomnography. Prior to nocturnal hypoxia, losartan had no effect on either the hyperoxic (losartan: 3.6 ± 1.1, placebo: 4.0 ± 0.6 l/min/mmHg; *P* = 0.9) or hypoxic HCVR (losartan: 5.3 ± 1.4, placebo: 5.7 ± 0.68 l/min/mmHg; *P* = 1.0). Likewise, losartan did not influence either the hyperoxic (losartan: 4.2 ± 1.3, placebo: 3.8 ± 1.1 l/min/mmHg; *P* = 0.5) or hypoxic HCVR (losartan: 6.6 ± 1.8, placebo: 6.3 ± 1.5 l/min/mmHg; *P* = 0.9) after nocturnal hypoxia. Cardiorespiratory responses to apnea and participants' apnea hypopnea indexes during placebo and losartan were similar (73 ± 15 vs. 75 ± 14 events/h; *P* = 0.9). Loop gain, which correlated with CSA severity (*r* = 0.94, *P* < 0.001), was similar between treatments. In summary, in young healthy men, hypoxia-induced CSA severity is strongly associated with loop gain, but the AT<sub>1</sub>R does not modulate chemoreceptor reflex sensitivity before or after 8 h of nocturnal hypoxia.

**Keywords:** chemoreceptor reflex, angiotensin receptor, hypoxia, sleep apnea, human

## INTRODUCTION

Angiotensin-II (ANG-II) is a physiologically active hormone of the renin-angiotensin system (RAS) and has target receptors in various tissues, including the carotid bodies and central nervous system (Allen, 1998; Li et al., 2006). Animal models have demonstrated that ANG-II primarily exerts its effect on the carotid body through the ANG-II type-I receptor (AT<sub>1</sub>R) (Allen, 1998). AT<sub>1</sub>Rs are found on chemosensitive glomus cells of mouse carotid bodies and are upregulated following ANG-II infusion (Allen, 1998). Administration of ANG-II consequently elevates peripheral chemoreceptor activity and augments the chemoreceptor response to hypoxia in animal models of heart failure (Allen, 1998; Li et al., 2006). In isolated rat carotid bodies, blockade of the AT<sub>1</sub>R with losartan, severely reduces the afferent response to hypoxic-hypercapnic challenges (Roy et al., 2018), suggesting the AT<sub>1</sub>R is critical for peripheral chemoreceptor sensitivity.

Exposure to sustained hypoxia also sensitizes the chemoreceptor reflex response in both animals and humans (Aaron and Powell, 1993; Richard et al., 2014). In rats exposed to chronic hypoxia, there is increased expression of angiotensinogen and AT<sub>1</sub>Rs in carotid body glomus cells, and activation by ANG-II is associated with increased excitatory response which is attenuated by AT<sub>1</sub>R blockade (Leung et al., 2000; Lam and Leung, 2003). In rats, administering an AT<sub>1</sub>R blocker attenuates the long-lasting activity in the carotid sinus nerve (CSN) following acute hypoxic-hypercapnic bouts, demonstrating a critical role for the AT<sub>1</sub>R in the chemoreceptor reflex response to intermittent hypercapnic hypoxia (Roy et al., 2018). While AT<sub>1</sub>R antagonists (i.e., losartan) effectively mitigate the lasting effects of intermittent hypoxia on muscle sympathetic nerve activity (MSNA) and blood pressure in humans (Foster et al., 2010; Jouett et al., 2017), it remains to be determined if AT<sub>1</sub>R blockade attenuates chemosensitivity (either peripheral or central) before or after hypoxia in humans.

Heightened chemosensitivity during hypoxia is understood to contribute to the development of periodic breathing with sojourn to high altitude. Mechanistically, the presence of arterial hypoxia acutely increases the ventilatory responsiveness to hypoxia and to hypercapnia (e.g., manifests as a reduced gap between eupneic partial pressure of carbon dioxide (PCO<sub>2</sub>) and the apneic threshold). This heightened peripheral chemoreceptor sensitivity, via an increase in the overall loop gain of the ventilatory control system, leads to reduced stability and overt patterns of cyclic apnea and hyperventilation particularly during sleep. Loop gain – a term that describes the overall stability of the ventilatory control system – combines the unique contributions of chemoreceptor sensitivity (controller gain) as well as how effectively ventilation increases arterial partial pressure of oxygen and reduces the arterial PCO<sub>2</sub> (plant gain) (White, 2005). While high loop gain is considered the leading mechanism of central sleep apnea (CSA) in patients with heart failure or at altitude, it is also the leading cause of obstructive sleep apnea (OSA) (White, 2005; Edwards et al., 2012). Although acetazolamide has been shown to lower both plant and loop gain, and improves sleep apnea severity (OSA

and CSA), no study has pharmacologically lowered loop gain via chemosensitivity and consequently improved sleep apnea (Edwards et al., 2012).

Here we tested whether AT<sub>1</sub>R blockade consequently reduces chemoreceptor sensitivity, loop gain, and the severity of CSA in human subjects. Specifically, we assessed the influence of AT<sub>1</sub>R blockade on chemosensitivity before and after 8 h of nocturnal hypoxia, and on the severity of hypoxia induced CSA as a human model of high-loop gain, chemoreflex-dependent sleep apnea. We hypothesized that losartan administration would attenuate the hypercapnic ventilatory response (HCVR) and the cardiorespiratory response to apnea before and after nocturnal hypoxia. Additionally, we hypothesized that losartan would reduce the severity of hypoxia induced CSA by reducing loop gain during 8 h of nocturnal hypoxia.

## MATERIALS AND METHODS

### Ethical Approval

This study was approved by the University of British Columbia Clinical Research Ethics Board (H17-02920), was registered as a clinical trial (ClinicalTrials.gov; NCT03335904) and conformed to the latest revision of the Declaration of Helsinki. Prior to enrollment in the study all participants provided written informed consent.

### Participants

Male participants ( $n = 14$ ) recruited from the University of British Columbia – Okanagan campus were screened to ensure they were normotensive (systolic blood pressure (SBP) < 140 mmHg, diastolic blood pressure (DBP) < 90 mmHg), had normal pulmonary function (>80% of predicted), and were free of sleep disordered breathing (apnea hypopnea index (AHI) < 5 events/hour) based on a home sleep study. Females were excluded from participation because they reportedly develop less severe CSA compared with males at high altitude (Lombardi et al., 2013) and in heart failure (Sin et al., 1999). Additionally, resting ventilation and blood pressure regulation are sensitive to menstrual cycle phase (Behan et al., 2003; Hart and Charkoudian, 2014). Participants were excluded if they have smoked within the past year and have a history of impaired renal function, cardiovascular disease, or respiratory disease. Participants were also excluded if they were taking any medication, prescribed or over the counter, or were obese (BMI > 30 Kg/m<sup>2</sup>).

### Protocol

Participants were asked to abstain from strenuous physical activity, alcohol or caffeine during the 12 h preceding each experimental session. Using a double-blind, placebo controlled, randomized, crossover design participants took either 50 mg of losartan, or a placebo pill three times over the course of 24 h (see section Pharmacological Intervention). Each experimental visit was separated by a 1 week washout period. Losartan was selected to target the AT<sub>1</sub>R as it has a high affinity for the AT<sub>1</sub>R and has been shown to be void of any agonist activity

(Ohtawa et al., 1993). The experimental protocol consisted of (1) a hyperoxic hypercapnic ventilatory response (HCVR) test, (2) a hypoxic HCVR test and (3) a hypoxic apnea response (HAR) test measured before and after a night of sleep in a normobaric hypoxia chamber (8850 SUMMIT+; Altitude Tech, Kingston, ON, Canada). In addition, venous blood samples were collected before and after a night of hypoxic sleep and assayed for plasma renin activity (PRA) and aldosterone to confirm effective dosing of losartan. Cardiovascular and respiratory variables were recorded continuously throughout the HCVR and HAR tests. Following the evening pre-tests, participants were instrumented with a sleep system (see section Monitoring Sleep in Normobaric Hypoxia) prior to entering the hypoxic chamber. The fraction of inspired  $O_2$  within the normobaric chamber was set to 0.135 which has been shown to sufficiently induce CSA in young, healthy men (Lombardi et al., 2013). Eight hours after entering the chamber, participants were awakened, exited the hypoxic chamber and took the third and final dose of their assigned intervention. They then completed the Lake Louise Acute Mountain Sickness (AMS) Scoring System and the ESQ-Cerebral Symptoms Questionnaire (Roach et al., 1993; Beidleman et al., 2007). The chemoreceptor reflex sensitivity tests were repeated 1 h after the last drug or placebo dose.

### Pharmacological Intervention

Participants were randomly assigned to orally ingest either losartan tablets (50 mg), or placebo pills (microcrystalline cellulose, identical in appearance and packaged in identical blister packages) at 8 h intervals. The first of the three doses was taken ~10 h before the experimental visit. The second dose was given 2 h before the evening chemoreceptor reflex tests while the third dose was given 1 h prior to morning testing. This timing was selected as plasma concentrations of both losartan and E-3174, its active metabolite, peak after a single dose and are not significantly different than concentrations after 7 days of losartan use (Ohtawa et al., 1993). Additionally, we elected the dosing protocol to ensure plasma concentrations of losartan and E-3147 remained sufficiently high throughout the entire protocol. Peak plasma concentration of losartan occurs an hour after ingestion while E-3147 reaches peak concentrations 2 h after ingestion and as such ventilatory response tests were performed within this time frame (Ohtawa et al., 1993).

### Chemoreceptor Reflex Test Instrumentation

All respiratory and cardiovascular parameters were acquired using an analog-to-digital converter (Powerlab/16SP ML 880; AD Instruments, Colorado Springs, CO, United States) interfaced with a personal computer. Commercially available software was used to analyze ventilatory and cardiovascular variables (LabChart V7.1, AD Instruments). During both sets of ventilatory tests, subjects breathed through a mouthpiece while wearing a nose clip, and a two-way non-rebreathing valve. Respired gas pressures were sampled at the mouth and analyzed for the partial pressure of end tidal oxygen and carbon dioxide

( $P_{ET}O_2$  and  $P_{ET}CO_2$  respectively) (ML206; AD Instruments). Expired gases were passed through a 4.7 L mixing chamber (MLA246; AD Instruments), as well as an oxygen ( $O_2$ ) and carbon dioxide ( $CO_2$ ) gas analyzer connected in series (S-3A and CD-3A, AEI Technologies, Pittsburgh, PA, United States) to measure the fraction of inspired  $O_2$  and fraction of inspired  $CO_2$  to calculate resting metabolic parameters including  $O_2$  and  $CO_2$  consumption as well as respiratory exchange rate. These parameters were necessary to determine the isometabolic hyperbola. Respiratory flow was also measured near the mouth using a pneumotachograph (HR 800L, Hans Rudolph, Shawnee, KS, United States) and a differential pressure amplifier (PA1 1110, Hans Rudolph). Heart rate (HR) was determined from a standard lead II electrocardiogram (ML132, AD Instruments). Beat-by-beat blood pressure was measured from a cuff placed on the right middle finger using pulse photoplethysmography (Finometer PRO; Finapres Medical Systems, Amsterdam, the Netherlands). A return to flow calibration was completed and the Finometer was referenced to manual blood pressures taken with an automated sphygmomanometer (Carescape V100; GE Medical Systems, Milwaukee, WI, United States) on the contralateral arm three times during baseline of each testing protocol.

### Hypercapnic Ventilatory Response (HCVR)

Two HCVR tests were administered to assess ventilatory and cardiovascular responses to  $CO_2$ , first on the background of hyperoxia and secondly on the background of hypoxia, with the latter maximizing peripheral chemoreflex activity. Each protocol was separated by 10 min. Performing the HCVR test in hyperoxia minimizes the peripheral chemoreceptor contribution to the chemoreflex response, while a background of hypoxia maximizes the peripheral chemoreceptor input (Duffin, 2007). Prior to starting the HCVR tests, participants rested supine for 10 min while breathing through a mouthpiece with their nose-clamped to collect baseline values of all variables. Throughout the test, participants listened to relaxing music with no prominent rhythm. Dynamic end-tidal forcing (see section Dynamic End-Tidal Forcing) was used to clamp  $P_{ET}O_2$  and  $P_{ET}CO_2$  values throughout the HCVR tests (Tymko et al., 2016). In short,  $P_{ET}O_2$  was clamped at 350 mmHg for the hyperoxic HCVR test and at 50 mmHg for the hypoxic HCVR.  $P_{ET}CO_2$  was clamped for 3 min each at 0, +2, +4, and +6 mmHg respective to each subject's baseline value.

### Dynamic End-Tidal Forcing

Control of  $P_{ET}O_2$  and  $P_{ET}CO_2$  was accomplished through an end-tidal forcing system which uses independent gas solenoid valves for  $O_2$ ,  $CO_2$ , and nitrogen ( $N_2$ ) to deliver a precise quantity of each gas into an inspiratory reservoir where it is humidified (Tymko et al., 2015, 2016). Breath-by-breath measures of  $P_{ET}O_2$ ,  $P_{ET}CO_2$ , tidal volume, breathing frequency and minute ventilation were done online using specifically designed software (Labview 13.0, National Instruments, Austin, TX, United States). The forcing system uses  $P_{ET}O_2$ ,  $P_{ET}CO_2$ , inspired and expired tidal volumes feedback and alters the

inspired gas to bring end-tidal gas levels to the target value. Feed-forward control of the inspire is based on estimates of metabolic  $O_2$  consumption and  $CO_2$  production and utilizes the alveolar gas equation to determine the required fraction of inspired  $O_2$  and fraction of inspired  $CO_2$ . Feedback control is accomplished using a proportional and integral error reduction control system.

### Calculating Chemoreflex Sensitivity

For both hyperoxic and hypoxic HCVR tests minute ventilation was plotted against  $P_{ET}CO_2$  for all participants. Individual data were regressed using a linear mixed effects model (see section Statistical Analyses) from which individual coefficients (i.e., slopes and intercepts) were calculated from random effects. Reported group means were determined from the global model coefficients representing the group's response to condition and drug.

### Hypoxic Apnea Response (HAR)

Following the HCVR protocols, participants remained instrumented and performed the hypoxic apnea response (HAR) protocol. A 5 min baseline was collected prior to performing six hypoxic apneas during which all respiratory and cardiovascular variables were recorded. Each apneic cycle was repeated 6 times and consisted of (1) 2–3 breaths through a three-way valve connected to a reservoir containing 100%  $N_2$ , (2) a 20 s end-expiratory breath-hold, and (3) 40 s of room air breathing. A nadir arterial oxyhemoglobin saturation ( $SpO_2$ ) between 85 and 90% was targeted during each apnea, and the number of  $N_2$  breaths prior to each apnea was adjusted to achieve this range. Following the hypoxic apnea cycles, all variables were recorded throughout a 5 min recovery period. The HAR test was performed to assess if ventilatory and cardiovascular responses to apnea were influenced by losartan.

### Monitoring Sleep in Normobaric Hypoxia

Sleep disordered breathing was measured using a continuous, overnight, cardiopulmonary monitoring system (Somte PSG, Compumedics, VIC, Australia). The device consists of an oximeter to record  $SpO_2$ , an electrocardiogram to record HR, a pressure transducer to record nasal airflow, chest and abdomen bands to measure respiratory effort, as well as a body position sensor. Additionally, monitoring of FP1-A2 and FP2-A1 by electroencephalography was combined with electrooculography of the left and right eye to assess sleep vs. wakefulness. Total sleep time was produced once wakefulness and sleep had been scored. The data were manually scored by the same investigator (Profusion 4, Compumedics, VIC, Australia) for the calculation of the AHI, the oxygen desaturation index (ODI) and the peak and nadir oxyhemoglobin saturation according to the criteria established by the American Academy of Sleep Medicine (Berry et al., 2012). In short, apneas ( $\geq 90\%$  reduction in peak nasal pressure) and hypopneas ( $\geq 30\%$  reduction in peak nasal pressure) were required to be at least 10 s in duration. Additionally, hypopneas had to be associated with a  $\geq 4\%$  desaturation. Arousals and sleep staging were not scored.

### Dynamic Loop Gain During Sleep

Loop gain is the dynamic ventilatory drive response that occurs consequent to a reduction in ventilation. We modified an established method to estimate loop gain during sleep (Terrill et al., 2015). The method involved (1) extracting scored respiratory events, and (2) generating a ventilation signal (tidal volume  $\times$  respiratory rate, uncalibrated) based on the overnight airflow signal (nasal pressure, linearized). For each window of data (3 min, modified from the usual 7 min to fit the faster cycling of hypoxia-induced CSA), ventilation data were used to model (i.e., explain) future values of ventilatory drive, where ventilatory drive was considered to be equal to ventilation between events (during central apneas ventilatory drive was considered to be subthreshold; i.e., late in the apnea, chemical drive increases and passes zero at the point of central apnea cessation). Using this approach, the average dynamic loop gain was calculated (for each participant, on each experimental night) by taking the median of the values from each 3 min window.

### Venous Blood Sample Collection and Processing

Venous blood samples taken for the measurement of PRA and aldosterone were drawn once in the evening following the ventilatory response tests and once in the morning at least 2 h following administration of the prescribed drug. With oral administration of losartan, PRA levels increase (Goldberg et al., 1993) which we utilized to confirm functional blockade of the  $AT_1R$ . Venous blood samples were collected in two 6.0 mL EDTA coated vacutainers. Collected samples were centrifuged at  $4^\circ C$  and separated into 2.0 mL aliquots. The separated plasma was stored in flat top microcentrifuge tubes and stored at  $-80^\circ C$  until later analyzed. PRA and aldosterone were assayed by radioimmunoassay techniques and the PRA assessed by measuring the amounts of ANG-I generated per hour.

### Statistical Analyses

Drug treatment was assigned using an online randomization tool<sup>1</sup>. All statistical analyses were performed in R statistical language (R Foundation for Statistical Computing, Vienna, Austria), lme4 (Bates et al., 2015), lmerTest (Kuznetsova et al., 2017), and emmeans (Lenth, 2019) statistical packages. Mixed effect linear modeling was used to interrogate all defined relationships and the correlation between loop gain and CSA severity, but not AMS responses. Across all models, participant was considered as a random effect, allowing for variable intercepts for each participant. When a significant F-test was achieved, pairwise comparisons were made with a Tukey's *post hoc* analysis to determine differences between the least square means. Statistical significance was set at a level of  $P < 0.05$ . A Mann–Whitney *U*-test was run on the AMS scores to detect differences in responses based on drug. With the exception of AMS scores and ESQ-Cerebral Symptoms Questionnaire (presented as medians and interquartile range [IQR]) data are presented as means  $\pm$  SEM.

<sup>1</sup>www.random.org

## RESULTS

### Participants

Due to technical and logistical limitations, three HCVR tests were excluded from analysis. An additional subject had less than 2 h of recorded sleep on their experimental night. While taking losartan, two participants developed symptoms of AMS, including nausea and vomiting, causing them to withdraw from further study. Both recovered within a few hours. Despite this, the Lake Louise AMS Score (losartan: 2.5 [3.2] (median [IQR], placebo: 3.5 [3.0];  $P = 0.3$ ) and ESQ-Cerebral Symptoms Questionnaire (losartan: 3.5 [13.0], placebo: 5.5 [10.2];  $P = 0.3$ ) did not differ between treatments. **Figure 1** illustrates the flow of participants through the chemosensitivity tests and the sample sizes used for statistical analysis.

All participants showed normal pulmonary function [forced vital capacity (FVC) =  $105 \pm 3\%$  predicted, forced expiratory volume in 1 s (FEV<sub>1</sub>) =  $97 \pm 3\%$  predicted, FEV<sub>1</sub>/FVC =  $93 \pm 1\%$  predicted], had a body mass index of  $24.6 \pm 1.1$  kg/m<sup>2</sup>, and were  $25 \pm 2$  years. Participants did not suffer from undiagnosed sleep disordered breathing (AHI =  $2.0 \pm 0.5$ , ODI =  $0.9 \pm 0.3$  events/hour), were not taking any additional medications and were normotensive (SBP =  $119 \pm 2$ , DBP =  $67 \pm 2$  mmHg).

### Influence of AT<sub>1</sub>R Blockade on Hypoxia-Induced CSA and Loop Gain


Sleep time did not differ between losartan and placebo (losartan:  $363 \pm 26$ ; placebo:  $324 \pm 16$  min,  $P = 0.2$ ). Additionally, we


observed similar ODI (losartan:  $78 \pm 12$ ; placebo:  $81 \pm 14$  events/hour,  $P = 0.7$ ), AHI (losartan:  $75 \pm 14$ ; placebo:  $73 \pm 15$  events/hour,  $P = 0.9$ ), average oxyhemoglobin desaturation (losartan:  $9.5 \pm 0.9$ ; placebo:  $8.9 \pm 1.0\%$ ,  $P = 0.3$ ), and nadir SpO<sub>2</sub> (losartan:  $61.3 \pm 1.2$ ; placebo:  $60.5 \pm 1.7\%$ ,  $P = 0.7$ ) between losartan and placebo. Loop gain was similar between treatments (losartan:  $0.88 \pm 0.05$ , placebo:  $0.88 \pm 0.06$  arbitrary units;  $P = 0.2$ ). There was a strong correlation present between loop gain and AHI ( $r = 0.93$ ,  $P < 0.001$ ; see **Figure 2**) while loop gain was moderately correlated with ODI ( $r = 0.63$ ,  $P = 0.02$ ).

### Influence of AT<sub>1</sub>R Blockade on Resting Cardiorespiratory Parameters Before and After Normobaric Hypoxia

As anticipated, P<sub>ET</sub>O<sub>2</sub> was higher following sleep in the hypoxic chamber ( $90.7 \pm 1.1$  and  $97.3 \pm 1.1$  mmHg,  $P < 0.001$ ) while P<sub>ET</sub>CO<sub>2</sub> fell ( $40.4 \pm 0.6$  and  $36.3 \pm 0.6$  mmHg,  $P < 0.001$ ). Alveolar ventilation was elevated in the post-test compared with the pre-test ( $5.4 \pm 0.3$  and  $6.4 \pm 0.3$  l/min,  $P < 0.05$ ) but was not influenced by drug (losartan:  $5.9 \pm 0.3$ , placebo:  $6.1 \pm 0.6$ ;  $P = 0.8$ ). Both P<sub>ET</sub>O<sub>2</sub> (losartan:  $93.4 \pm 1.0$ , placebo:  $94.6 \pm 1.0$  mmHg,  $P = 0.03$ ) and P<sub>ET</sub>CO<sub>2</sub> (losartan:  $38.0 \pm 0.5$ , placebo:  $38.7 \pm 0.5$  mmHg,  $P = 0.05$ ) were similar between drug treatments.

Although we observed a drug by condition interaction for SBP ( $P = 0.02$ ), *post hoc* analysis did not reveal any pairwise differences. Losartan did not influence DBP (losartan:  $63 \pm 1$ , placebo:  $65 \pm 1$  mmHg), mean arterial pressure (MAP) (losartan:





81 ± 1, placebo: 83 ± 2 mmHg), or HR (losartan: 63 ± 2, placebo: 63 ± 2 bpm;  $P > 0.05$  for all comparisons). Similarly, DBP (63 ± 2, post: 65 ± 2 mmHg), MAP (pre: 81 ± 2, post: 83 ± 2 mmHg), and HR (pre: 62 ± 2, post: 63 ± 2) did not differ between the pre- and post-tests ( $P > 0.1$  for all conditions).

### Influence of AT<sub>1</sub>R Blockade and Normobaric Hypoxia on the HCVR Test

**Figure 3** shows mean data, isometabolic hyperbola, and mixed effect linear models illustrating the hyperoxic and hypoxic HCVR before and after 8 h of nocturnal hypoxia for losartan and placebo. As expected, the HCVR was significantly greater in hypoxia (5.3 ± 1.1 l/min/mmHg) compared with hyperoxia (3.6 ± 0.6 l/min/mmHg;  $P = 0.02$ ). There was no significant difference to the HCVR following 8 h of sleep in normobaric hypoxia (5.0 ± 0.5 and 6.2 ± 0.5 l/min/mmHg,  $P = 0.7$ ). Detailed results for these tests are presented in the following two sections.

#### Hyperoxic HCVR Test

The hyperoxic HCVR (**Figure 3A**) was similar in both losartan and placebo conditions during the pre- (3.9 ± 1.1 and 3.5 ± 0.5 l/min, respectively;  $P = 0.9$ ) and post-tests (3.4 ± 0.5 and 3.6 ± 1.2, respectively;  $P = 0.5$ ), and did not differ following poikilocapnic normobaric hypoxia ( $P = 0.8$ ). **Table 1** summarizes cardiorespiratory parameters throughout the hyperoxic HCVR. There was no significant main effect of condition (pre vs. post), or a condition-by-drug-by-P<sub>ET</sub>CO<sub>2</sub> stage interaction across all cardiorespiratory variables. P<sub>ET</sub>O<sub>2</sub> and P<sub>ET</sub>CO<sub>2</sub> were significantly increased from baseline across all stages of the test. A main effect for stage was observed for minute ventilation, which was significantly greater than baseline at all P<sub>ET</sub>CO<sub>2</sub> stages (see **Table 1**). MAP was not affected by drug or condition and increased with each P<sub>ET</sub>CO<sub>2</sub> level (**Table 1**). The HR response was similar between losartan and placebo and increased with P<sub>ET</sub>CO<sub>2</sub> level.

#### Hypoxic HCVR Test

The hypoxic HCVR (**Figure 3B**) was similar in both losartan and placebo conditions during the pre- (losartan: 5.6 ± 1.5, placebo: 5.4 ± 0.6 l/min;  $P = 1.0$ ) and post-tests (losartan: 5.6 ± 1.5, placebo: 6.4 ± 1.2 l/min/mmHg;  $P = 0.9$ ), and did not differ following poikilocapnic normobaric hypoxia ( $P = 0.7$ ). **Table 2** summarizes the cardiorespiratory parameters throughout the hypoxic HCVR test. There was no significant main effect of condition (pre vs. post), or a condition-by-drug-by-P<sub>ET</sub>CO<sub>2</sub> stage interaction across all cardiorespiratory variables. P<sub>ET</sub>O<sub>2</sub> was slightly lower on losartan (59.7 ± 0.3 and 60.6 ± 0.3 mmHg;  $P = 0.03$ ) and was reduced from baseline across all levels of P<sub>ET</sub>CO<sub>2</sub> by study design. P<sub>ET</sub>CO<sub>2</sub> was increased across each stage of the test, but was slightly lower overall on losartan compared with placebo (41.0 ± 0.8 and 42.2 ± 0.8 mmHg;  $P = 0.001$ ). A main effect for stage was observed for minute ventilation, which was significantly greater than baseline at all P<sub>ET</sub>CO<sub>2</sub> stages (**Table 2**). Neither MAP nor HR were affected by drug or condition, but both increased with each P<sub>ET</sub>CO<sub>2</sub> level (**Table 2**).

### Influence of AT<sub>1</sub>R Blockade and Normobaric Hypoxia on Hypoxic Apnea Response

#### Ventilatory Response


During the HAR tests SpO<sub>2</sub> fell to 87 ± 1% and was similar across conditions ( $P = 0.10$ ) and drug ( $P = 0.07$ ). **Figure 4** shows the mean breath-by-breath change in ventilatory variables, ensemble averaged across all 6 hypoxic apneas. All ventilatory parameters were significantly elevated from baseline during the first breath following apnea cessation with the exception of breathing frequency which was significantly higher following exposure to normobaric hypoxia (20.9 ± 1.5 and 18.7 ± 1.4 bpm;  $P = 0.03$ ). Losartan attenuated breathing frequency following normobaric hypoxia during the post-test (11.3 ± 1.2 and 12.4 ± 1.2 bpm;  $P = 0.03$ ) but not during the pre-test (12.0 ± 1.1 and 11.9 ± 1.2 bpm;  $P = 0.5$ ). Tidal volume and breathing frequency were similar between conditions.

#### Cardiovascular Response

**Figure 5** displays the change in HR and MAP averaged over six apneas. All parameters increased over the duration of the end expiratory breath hold. The 20 s apnea elicited a similar increase in both HR and MAP between both losartan and placebo conditions. Both drug and condition did not have a significant impact on changes in SBP, DBP, or MAP with respect to percent desaturation as summarized in **Table 3**. There was a trend toward a heightened HR response following sleep in the hypoxic chamber ( $P = 0.06$ ).

### Influence of AT<sub>1</sub>R Blockade on Markers of the RAS

Angiotensin-I was significantly higher following losartan compared with placebo (8.92 ± 1.32 and 2.94 ± 1.30 ng/mL,  $P < 0.001$ ). Angiotensin-I was also greater following normobaric hypoxia compared with baseline (7.61 ± 1.32 and 4.25 ± 1.30 ng/mL,  $P = 0.02$ ). PRA was elevated by losartan compared with



**TABLE 1 |** Cardiorespiratory parameters measured during the hyperoxic HCVR test.


	Drug	Baseline $P_{ET}CO_2$		+2 mmHg $P_{ET}CO_2$		+4 mmHg $P_{ET}CO_2$		+6 mmHg $P_{ET}CO_2$	
		Pre	Post	Pre	Post	Pre	Post	Pre	Post
$\dot{V}_I$ (l/min)	Losartan	16 $\pm$ 2	17 $\pm$ 2	23 $\pm$ 2	29 $\pm$ 4	34 $\pm$ 3	36 $\pm$ 4	42 $\pm$ 5	45 $\pm$ 4
	Placebo	14 $\pm$ 1	18 $\pm$ 1	20 $\pm$ 2	26 $\pm$ 2	29 $\pm$ 3	35 $\pm$ 3	35 $\pm$ 4	42 $\pm$ 3
LSM		18 $\pm$ 2		27 $\pm$ 2*		37 $\pm$ 2*†		45 $\pm$ 2*††	
				Drug: $P = 0.8$		Stage: $P < 0.001$		Drug*Stage: $P = 1.0$	
$P_{ET}O_2$ (mmHg)	Losartan	91 $\pm$ 1	96 $\pm$ 2	343 $\pm$ 7	353 $\pm$ 13	337 $\pm$ 3	348 $\pm$ 15	339 $\pm$ 3	347 $\pm$ 16
	Placebo	91 $\pm$ 1	99 $\pm$ 1	339 $\pm$ 4	339 $\pm$ 7	346 $\pm$ 3	346 $\pm$ 2	347 $\pm$ 3	345 $\pm$ 3
LSM		94 $\pm$ 3		343 $\pm$ 3*		344 $\pm$ 3*		344 $\pm$ 3*	
				Drug: $P = 0.6$		Stage: $P < 0.001$		Drug*Stage: $P = 0.3$	
$P_{ET}CO_2$ (mmHg)	Losartan	40 $\pm$ 1	36 $\pm$ 1	42 $\pm$ 1	38 $\pm$ 1	44 $\pm$ 1	41 $\pm$ 1	46 $\pm$ 1	42 $\pm$ 1
	Placebo	41 $\pm$ 1	37 $\pm$ 1	43 $\pm$ 1	39 $\pm$ 1	45 $\pm$ 1	41 $\pm$ 1	47 $\pm$ 1	43 $\pm$ 1
LSM		38 $\pm$ 1		41 $\pm$ 1*		43 $\pm$ 1*†		45 $\pm$ 1*††	
				Drug: $P = 0.09$		Stage: $P < 0.001$		Drug*Stage: $P = 1.0$	
MAP (mmHg)	Losartan	80 $\pm$ 1	81 $\pm$ 2	84 $\pm$ 2	84 $\pm$ 2	87 $\pm$ 2	86 $\pm$ 2	91 $\pm$ 2	89 $\pm$ 2
	Placebo	83 $\pm$ 2	84 $\pm$ 2	86 $\pm$ 3	87 $\pm$ 2	87 $\pm$ 3	88 $\pm$ 2	90 $\pm$ 3	92 $\pm$ 3
LSM		82 $\pm$ 1		85 $\pm$ 1*		87 $\pm$ 1*		91 $\pm$ 1*††	
				Drug: $P = 0.6$		Stage: $P < 0.001$		Drug*Stage: $P = 0.9$	
HR (bpm)	Losartan	62 $\pm$ 2	64 $\pm$ 3	61 $\pm$ 2	63 $\pm$ 3	64 $\pm$ 2	64 $\pm$ 3	68 $\pm$ 2	70 $\pm$ 3
	Placebo	63 $\pm$ 2	63 $\pm$ 2	61 $\pm$ 2	60 $\pm$ 2	63 $\pm$ 2	63 $\pm$ 2	66 $\pm$ 2	65 $\pm$ 2
LSM		63 $\pm$ 1		61 $\pm$ 1		64 $\pm$ 1†		67 $\pm$ 1*††	
				Drug: $P = 0.4$		Stage: $P < 0.001$		Drug*Stage: $P = 0.4$	

All values are mean  $\pm$  SEM. \*Indicates a significant difference from baseline ( $p < 0.05$ ); †significant difference from +2 mmHg ( $p < 0.05$ ); ††significant difference from +4 mmHg ( $p < 0.05$ ). LSM, least square marginal means for stage;  $\dot{V}_I$ , minute ventilation;  $P_{ET}O_2$ , end-tidal oxygen;  $P_{ET}CO_2$ , end-tidal carbon dioxide; MAP, mean arterial pressure; HR, heart rate. Bolded values indicate significant main effects.


**TABLE 2** | Ventilatory parameters measured during the hypoxic HCVR test.

	Drug	Baseline $P_{ET}CO_2$		+2 mmHg $P_{ET}CO_2$		+4 mmHg $P_{ET}CO_2$		+6 mmHg $P_{ET}CO_2$	
		Pre	Post	Pre	Post	Pre	Post	Pre	Post
$\dot{V}_I$ (l/min)	Losartan	16 ± 2	17 ± 1	37 ± 5	50 ± 4	45 ± 4	59 ± 4	50 ± 5	61 ± 3
	Placebo	15 ± 1	18 ± 1	41 ± 5	52 ± 4	47 ± 5	60 ± 5	54 ± 5	62 ± 4
	LSM	18 ± 2		47 ± 2*		55 ± 2*†		59 ± 3*†‡	
				Drug: $P = 0.03$		Stage: $P < 0.001$		Drug*Stage: $P = 0.6$	
$P_{ET}O_2$ (mmHg)	Losartan	92 ± 2	96 ± 1	49 ± 1	50 ± 0	49 ± 0	50 ± 0	49 ± 1	49 ± 0
	Placebo	92 ± 2	98 ± 1	50 ± 1	51 ± 1	50 ± 1	50 ± 1	50 ± 1	50 ± 1
	LSM	92 ± 1		50 ± 1*		50 ± 0*		49 ± 1*	
				Drug: $P = 0.03$		Stage: $P < 0.001$		Drug*Stage: $P = 1.0$	
$P_{ET}CO_2$ (mmHg)	Losartan	40 ± 1	35 ± 0	42 ± 1	37 ± 1	44 ± 1	39 ± 1	46 ± 1	41 ± 1
	Placebo	41 ± 1	37 ± 1	43 ± 1	39 ± 1	45 ± 1	41 ± 1	47 ± 1	43 ± 1
	LSM	39 ± 1		41 ± 1*		43 ± 1*†		45 ± 1*†	
				Drug: $P = 0.03$		Stage: $P < 0.001$		Drug*Stage: $P = 0.6$	
MAP (mmHg)	Losartan	82 ± 2	86 ± 3	87 ± 3	94 ± 3	90 ± 3	96 ± 3	91 ± 3	99 ± 4
	Placebo	87 ± 3	88 ± 2	94 ± 3	98 ± 2	98 ± 3	99 ± 3	102 ± 4	101 ± 3
	LSM	86 ± 1		94 ± 2*		97 ± 2*		99 ± 2*†	
				Drug: $P = 0.3$		Stage: $P < 0.001$		Drug*Stage: $P = 0.9$	
HR (bpm)	Losartan	63 ± 2	60 ± 2	77 ± 2	77 ± 2	80 ± 2	82 ± 2	82 ± 2	82 ± 2
	Placebo	63 ± 2	60 ± 3	79 ± 2	82 ± 2	82 ± 2	82 ± 2	86 ± 2	88 ± 2
	LSM	61 ± 1		78 ± 2*		81 ± 2*		83 ± 2*†	
				Drug: $P = 0.7$		Stage: $P < 0.001$		Drug*Stage: $P = 0.2$	

All values are mean ± SEM. \*Indicates a significant difference from baseline ( $p < 0.05$ ); †Significant difference from +2 mmHg ( $p < 0.05$ ); ‡significant difference from +4 mmHg ( $p < 0.05$ ). LSM, least mean squares;  $\dot{V}_I$ , minute ventilation;  $P_{ET}O_2$ , end-tidal oxygen;  $P_{ET}CO_2$ , end-tidal carbon dioxide; MAP, mean arterial pressure; HR, heart rate. Bolded values indicate significant main effects.



**FIGURE 4** | Ensemble averaged breath-by-breath trace of hypoxic apnoea response test before (A) and after (B) normobaric hypoxia. Breath-by-breath trace beginning 5 breaths prior to apnoea start. Ventilation is increased prior to the apnoea, as participants were asked to take large breaths during nitrogen administration. Values are mean ± SEM. \* $P < 0.05$  compared with respective baseline.  $\dot{V}_I$ , minute ventilation;  $V_T$ , tidal volume; Fb, breathing frequency.



**FIGURE 5 |** Beat-by-beat ensemble averaged trace of hypoxic apnea response test before (A) and after (B) normobaric hypoxia. Beat-by-beat trace beginning 30 s prior to apnea start. Values were signal averaged between participants and across the 6 apneas. Values shown are the mean  $\pm$  SEM. HR, heart rate; MAP, mean arterial pressure.

**TABLE 3 |** Effect of AT<sub>1</sub>R blockade on cardiovascular sensitivity to hypoxic apnea before (Pre) and after (Post) 8 h of nocturnal hypoxia.

	Drug	Pre	Post	Drug	Condition	Interaction
$\Delta$ SBP/ $\Delta$ SpO <sub>2</sub> (mmHg/%desaturation)	Losartan	3.1 $\pm$ 0.6	3.6 $\pm$ 0.7	$P = 0.22$	$P = 0.90$	$P = 0.28$
	Placebo	3.1 $\pm$ 0.6	2.4 $\pm$ 0.6			
$\Delta$ DBP/ $\Delta$ SpO <sub>2</sub> (mmHg/%desaturation)	Losartan	2.3 $\pm$ 0.4	2.9 $\pm$ 0.5	$P = 0.60$	$P = 0.30$	$P = 0.23$
	Placebo	2.5 $\pm$ 0.4	2.2 $\pm$ 0.5			
$\Delta$ MAP/ $\Delta$ SpO <sub>2</sub> (mmHg/%desaturation)	Losartan	2.5 $\pm$ 0.5	3.1 $\pm$ 0.7	$P = 0.43$	$P = 0.98$	$P = 0.24$
	Placebo	2.7 $\pm$ 0.5	2.1 $\pm$ 0.4			
$\Delta$ HR/ $\Delta$ SpO <sub>2</sub> (bpm/%desaturation)	Losartan	1.6 $\pm$ 0.3	2.5 $\pm$ 0.4	$P = 0.42$	$P = 0.06$	$P = 0.14$
	Placebo	1.8 $\pm$ 0.3	1.9 $\pm$ 0.3			

All values are mean  $\pm$  SEM.  $\Delta$ SpO<sub>2</sub>, % of oxyhemoglobin desaturation;  $\Delta$ SBP, change in systolic blood pressure;  $\Delta$ DBP, change in diastolic blood pressure;  $\Delta$ MAP, change in mean arterial pressure;  $\Delta$ HR, change in heart rate.

placebo ( $0.83 \pm 0.12$  and  $0.27 \pm 0.12$  ng/l/s,  $P < 0.001$ ) and following the sleep in the hypoxic chamber ( $0.70 \pm 0.12$  and  $0.39 \pm 0.12$  ng/l/s, respectively,  $P = 0.01$ ). Although aldosterone was higher following normobaric hypoxia compared with baseline ( $135.8 \pm 14.6$  and  $78.1 \pm 14.3$  pmol/l, respectively;  $P < 0.01$ ), it was reduced by losartan compared with placebo ( $76.3 \pm 14.6$  and  $137.5 \pm 14.3$  pmol/l,  $P < 0.01$ ).

## DISCUSSION

The purpose of this study was to determine if AT<sub>1</sub>R blockade would attenuate (1) the chemoreceptor reflex to CO<sub>2</sub> and voluntary hypoxic apnea before or after 8 h of nocturnal hypoxia, and (2) the severity of hypoxia-induced CSA through reductions in loop gain. We observed similar ventilatory response to hypercapnia and cardiorespiratory responses to voluntary

hypoxic apneas between losartan and placebo before and after nocturnal hypoxia. Additionally, the severity of hypoxia-induced CSA and loop gain was unaffected by AT<sub>1</sub>R blockade. Our data suggests that in healthy, young males the chemoreceptor reflex and the severity of hypoxia-induced CSA occurs through a pathway independent of the AT<sub>1</sub>R and local RAS. Upregulation of the AT<sub>1</sub>R through pathology (e.g., heart failure) or chronic hypoxia may be required before functional changes in the chemoreceptor reflex are observed following AT<sub>1</sub>R blockade.

### Influence of AT<sub>1</sub>R Blockade on the Hypercapnic and Voluntary Hypoxic Apnea Chemoreceptor Reflex Before and Following Nocturnal Hypoxia

The AT<sub>1</sub>R is highly expressed in animal carotid body glomus cells and brain regions involved in cardiorespiratory control

including the nucleus tractus solitarius, subfornical organ, median preoptic nucleus, paraventricular nucleus, and rostral ventrolateral medulla (Dampney et al., 2002; Gao et al., 2005; Saxena et al., 2015; Wang et al., 2016; Shell et al., 2019). Activation of AT<sub>1</sub>R by ANG-II in these regions are implicated in the long-term facilitation of the carotid body and the sympathetic nervous system following exposure to intermittent hypoxia (Roy et al., 2018). Additionally, carotid body sensitivity is severely attenuated by losartan in the isolated carotid body preparation (Roy et al., 2018). However, we found the chemoreceptor reflex to hypercapnia and hypoxic hypercapnia to be similar between placebo and losartan which may reflect minimal AT<sub>1</sub>R expression and involvement in chemoreflex modulation in the healthy human. Indeed, our results corroborate those of Foster et al. (2010) who observed no effect of losartan on ventilatory and MAP responses to hypoxia and those of Solaiman et al. (2014) who found no effect of ANG-II infusion in potentiating the ventilatory response to hypoxia or hypercapnia in healthy humans.

Hypoxia activates the carotid body leading to greater ventilatory drive and chemoreceptor reflex sensitivity (Allen, 1998; Marcus et al., 2010). The mechanisms responsible for hypoxic mediated plasticity of the chemoreceptor reflex are not well understood. In rats, carotid body AT<sub>1</sub>R expression was doubled following 4 weeks of chronic hypoxia (10% O<sub>2</sub>) and the carotid body's sensitivity to ANG-II was enhanced but could be blocked by losartan (Leung et al., 2000). These data support a role of the AT<sub>1</sub>R in mediating carotid body neuroplasticity in response to hypoxia. In addition, carotid body angiotensin converting enzyme activity is doubled following 7 days of hypoxia supporting the presence of a local RAS (Lam et al., 2004). In contrast, we exposed healthy humans to 8 h of continuous hypoxia during sleep which led to minimal chemoreceptor reflex alteration as evidenced by the insignificant change in the hypoxic and hyperoxic HCVR. We observed evidence of augmented chemoreflex drive following 8 h of hypoxia including greater alveolar ventilation, P<sub>ET</sub>O<sub>2</sub>, and reduced P<sub>ET</sub>CO<sub>2</sub>. However, hyperoxic and hypoxic chemoreceptor reflex sensitivity were similar following 8 h of hypoxia and losartan did not modulate chemoreflex sensitivity indicating that AT<sub>1</sub>R may not contribute to changes in chemoreflex sensitivity, in healthy young males, following a single night of hypoxia.

Losartan is able to effectively attenuate the increase in sympathetic nerve activity and oxidative stress following both sustained and intermittent hypoxia (Leung et al., 2000; Marcus et al., 2010; Pialoux et al., 2011; Jouett et al., 2017). In rats exposed to 28 days of chronic intermittent hypoxia, the lumbar sympathetic nerve activity response to 20 s apneas was attenuated by losartan (Marcus et al., 2010). The exaggerated lumbar sympathetic and MAP responses to breath hold following chronic intermittent hypoxia were attenuated by losartan, an effect likely mediated by blunted upregulation of AT<sub>1</sub>R expression on the carotid body (Marcus et al., 2010). In contrast, losartan does not influence the ventilatory response to step changes in normocapnic hypoxia in patients with OSA (Morgan et al.,

2018). In agreement with our findings, this indicates an alternative mechanism involved in chemoreceptor sensitization that is independent of the AT<sub>1</sub>R. For example, heart failure models suggest a reduction in carotid body blood flow is necessary to reduce neural nitric oxide synthase expression while elevating carotid body AT<sub>1</sub>R expression and ANG-II concentration (Li and Schultz, 2006). Until upregulation of the AT<sub>1</sub>R occurs in animal models of congestive heart failure, losartan has no effect on chemosensitivity (Li et al., 2006). Once AT<sub>1</sub>Rs are upregulated, losartan is able to abolish the ventilatory and renal sympathetic responses to graded levels of hypoxia (Ding et al., 2011). Therefore, the AT<sub>1</sub>R may determine hypoxic chemoreceptor sensitivity in pathological states such as heart failure rather than in healthy humans acclimating to acute hypoxia.

There is limited data regarding chemosensitive signaling mechanisms within the human carotid body. In a previous human study, ANG-II infusion did not potentiate the ventilatory response to hypoxia or hypercapnia further suggesting that the AT<sub>1</sub>R may not play a role in chemosensitivity in healthy subjects (Solaiman et al., 2014). Supporting this, existing data suggests some similarities as well as species differences with respect to carotid body oxygen sensing and reactive oxygen species (ROS) generation. Similar to animal models, the human carotid body releases acetylcholine and ATP in response to hypoxia and expresses haemoxygenase-2, NADPH oxidase (NOX-2), AMP activated protein kinase (AMPK), and oxygen sensitive K<sup>+</sup> channels (Fagerlund et al., 2010; Mkrtchian et al., 2012; Kählin et al., 2014). However, there are also important differences from mice such as the presence of hydrogen sulfide (H<sub>2</sub>S) synthesizing enzyme cystathionine- $\gamma$ -lyase and absence of TASK-3 channels in human carotid bodies (Mkrtchian et al., 2012). Despite these known species differences, there remains no evidence within the literature for the local expression of AT<sub>1</sub>R or the components of a RAS in the human carotid body.

## Influence of AT<sub>1</sub>R Blockade on Loop Gain and the Severity of Hypoxia Induced CSA

Elevated loop gain contributes to OSA severity and a high controller gain (mostly measured during wakefulness) is associated with CSA in healthy humans at high altitude, in idiopathic CSA (Ainslie et al., 2013; Terrill et al., 2015), and in patients with heart failure (Solin et al., 2000; Javaheri, 2006). Notably, loop gain during sleep, measured using dynamic application of hypoxic-hypercapnia is remarkably similar to values during wakefulness (Messineo et al., 2018). Thus, individuals with high chemoreflex sensitivity likely suffer from worse CSA severity (Messineo et al., 2018). Our findings support this idea, demonstrating a strong relationship between dynamic loop gain measured during sleep and hypoxia-induced CSA severity ( $r = 0.93$ ). Likewise, interventions that reduce loop gain (acetazolamide, supplemental oxygen) are associated with attenuated central and obstructive sleep apnea severity (Javaheri, 2006; Edwards et al., 2012). In animals, losartan attenuates

chemosensitivity, a key contributor to loop gain, which we postulated would reduce CSA severity (Marcus et al., 2010; Ding et al., 2011; Roy et al., 2018). However, we did not observe differences between placebo and losartan for any indices of CSA severity. Similarly, Lombardi et al. (2013) did not find an effect of telmisartan, another AT<sub>1</sub>R antagonist, on the severity of high-altitude CSA during sojourn to 5400 m. This effect is most likely explained by the lack of drug effect on chemosensitivity.

## LIMITATIONS AND CONCLUSION

### Limitations

The AT<sub>1</sub>R block may have had a limited effect at baseline due to relatively low RAS activity in our participants. Jones et al. (2007) found that baseline PRA levels are correlated with changes in PRA 24 h following administration of an AT<sub>1</sub>R blockade in healthy volunteers. Participants with the lowest resting PRA levels experienced the smallest increases in PRA following oral administration of the drugs, indicating that AT<sub>1</sub>R blockade has a smaller effect in those that do not have an already upregulated RAS (Jones et al., 2007). However, we found no relationship between changes in PRA and the HCVR. It may be that an upregulated RAS is first necessary for an AT<sub>1</sub>R blockade to have a potent effect on chemosensitivity and sympathetic activity (Li et al., 2006).

A larger dose or dosing period of losartan may have led to different results. At baseline, participants had already received two 50 mg doses of losartan, 10 and 2 h prior to experimentation. We observed significant increases in PRA suggesting functional AT<sub>1</sub>R blockade, although this response was variable amongst participants. Previously, Foster et al. (2010) found that 4 days of losartan (100 mg) blocked the hypertensive response induced by 6 h of intermittent hypoxia. Another human study investigating long-term facilitation of sympathetic nerve activity induced by intermittent hypoxia found that a single 100 mg dose of losartan 1 h prior to experimentation was sufficient to abolish this response. This effect is also believed to be a function of AT<sub>1</sub>R activation within the carotid body. Considering these two extremes, the dosing protocol used in this study should have been adequate to detect any significant physiological outcomes.

We may not have seen a potentiation in chemosensitivity following nocturnal hypoxia because of the time of day the chemoreceptor reflex was measured. Chemoreceptor reflex sensitivity is affected by circadian rhythms independent of changes in metabolic rate (Stephenson et al., 2000). Although basal ventilation remains constant, the chemoreceptor reflex response to hypercapnia is heightened at night and attenuated during the day. This may explain why we did not see a greater difference in the chemoreflex response following exposure to nocturnal hypoxia. Since all participants performed both placebo and treatment arms of this study, diurnal variations in chemoreceptor sensitivity would not have influenced the effect of losartan on ventilatory response to hypercapnia or apnea.

## Conclusion

The current study examined whether AT<sub>1</sub>R blockade attenuates chemoreceptor reflex sensitivity, loop gain, and CSA severity in healthy, young males using a model of nocturnal hypoxia. We found that losartan did not influence chemoreceptor reflex sensitivity, loop gain, or the severity of CSA. Interestingly, we observed a strong relationship between dynamic loop gain measured during sleep and the severity of hypoxia-induced CSA, consistent with the view that the manifest pattern was indeed chemoreflex driven as intended. Overall, our data show that activation of the AT<sub>1</sub>R does not contribute to the chemoreceptor response to hypercapnia including its central and peripheral contributions, before or after a single night of sustained hypoxia in healthy males. It remains feasible that the AT<sub>1</sub>R may contribute to chemoreceptor reflex sensitivity in pathological states such as heart failure, obstructive sleep apnea, or following chronic hypoxia.

## DATA AVAILABILITY STATEMENT

The datasets generated for this study are available on request to the corresponding author.

## ETHICS STATEMENT

This study involving human participants was reviewed and approved by Clinical Research Ethics Board, University of British Columbia. The participants provided their written informed consent to participate in this study.

## AUTHOR CONTRIBUTIONS

CB, RW, SS, and GF: study design. CB, LB, TV, and GF: data collection. CB, LB, SS, and GF: data analysis. All authors: Interpretation, drafting and approval of the manuscript.

## FUNDING

This work was supported by the Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant (GF), Heart and Stroke Foundation of Canada Grant-in-aid (GF, NA, and JF), Canada Foundation for Innovation (GF), Canadian Institute of Health Research (RW). GF was a Michael Smith Foundation for Health Research Scholar. LB was funded by a Canada Graduate Scholarship - Doctoral from NSERC. SS was supported by the American Heart Association (15SDG25890059).

## ACKNOWLEDGMENTS

Analytical assistance by Amanda Vallance is gratefully acknowledged.

## REFERENCES

- Aaron, E. A., and Powell, F. L. (1993). Effect of chronic hypoxia on hypoxic ventilatory response in awake rats. *J. Appl. Physiol.* 74, 1635–1640. doi: 10.1152/jappl.1993.74.4.1635
- Ainslie, P. N., Lucas, S. J. E., and Burgess, K. R. (2013). Breathing and sleep at high altitude. *Respir. Physiol. Neurobiol.* 188, 233–256. doi: 10.1016/j.resp.2013.05.020
- Allen, A. M. (1998). Angiotensin AT1 receptor-mediated excitation of rat carotid body chemoreceptor afferent activity. *J. Physiol.* 510, 773–781. doi: 10.1111/j.1469-7793.1998.773bj.x
- Bates, D., Mächler, M., Bolker, B. M., and Walker, S. C. (2015). Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* 67, 1–48. doi: 10.18637/jss.v067.i01
- Behan, M., Zabka, A. G., Thomas, C. F., and Mitchell, G. S. (2003). Sex steroid hormones and the neural control of breathing. *Respir. Physiol. Neurobiol.* 136, 249–263.
- Beidleman, B. A., Muza, S. R., Fulco, C. S., Rock, P. B., and Cymerman, A. (2007). Validation of a shortened electronic version of the environmental symptoms questionnaire. *High Alt. Med. Biol.* 8, 192–199. doi: 10.1089/ham.2007.1016
- Berry, R. B., Budhiraja, R., Gottlieb, D. J., Gozal, D., Iber, C., Kapur, V. K., et al. (2012). Rules for scoring respiratory events in sleep: update of the 2007 AASM manual for the scoring of sleep and associated events. *J. Clin. Sleep Med.* 8, 597–619. doi: 10.5664/jcs.m.2172
- Dampney, R. A. L., Fontes, M. A. P., Hirooka, Y., Horiuchi, J., Potts, P. D., and Tagawa, T. (2002). Role of angiotensin II receptors in the regulation of vasomotor neurons in the ventrolateral medulla. *Clin. Exp. Pharmacol. Physiol.* 29, 467–472. doi: 10.1046/j.1440-1681.2002.03658.x
- Ding, Y., Li, Y. L., and Schultz, H. D. (2011). Role of blood flow in carotid body chemoreflex function in heart failure. *J. Physiol.* 589, 245–258. doi: 10.1113/jphysiol.2010.200584
- Duffin, J. (2007). Measuring the ventilatory response to hypoxia. *J. Physiol.* 584, 285–293. doi: 10.1113/jphysiol.2007.138883
- Edwards, B. A., Sands, S. A., Eckert, D. J., White, D. P., Butler, J. P., Owens, R. L., et al. (2012). Acetazolamide improves loop gain but not the other physiological traits causing obstructive sleep apnoea. *J. Physiol.* 590, 1199–1211. doi: 10.1113/jphysiol.2011.223925
- Fagerlund, M. J., Kählin, J., Ebberyd, A., Schulte, G., Mkrtchian, S., and Eriksson, L. I. (2010). The human carotid body: expression of oxygen sensing and signaling genes of relevance for anesthesia. *Anesthesiology* 113, 1270–1279. doi: 10.1097/ALN.0b013e318fac061
- Foster, G. E., Hanly, P. J., Ahmed, S. B., Beaudin, A. E., Pialoux, V., and Poulin, M. J. (2010). Intermittent hypoxia increases arterial blood pressure in humans through a renin-angiotensin system-dependent mechanism. *Hypertension* 56, 369–377. doi: 10.1161/HYPERTENSIONAHA.110.152108
- Gao, L., Wang, W., Li, Y.-L., Schultz, H. D., Liu, D., Cornish, K. G., et al. (2005). Sympathoexcitation by central ANG II: roles for AT 1 receptor upregulation and NAD(P)H oxidase in RVLM. *Am. J. Physiol. Heart Circ. Physiol.* 288, H2271–H2279. doi: 10.1152/ajpheart.00949.2004
- Goldberg, M. R., Tanaka, W., Barchowsky, A., Bradstreet, T. E., McCrea, J., Lo, M. W., et al. (1993). Effects of losartan on blood pressure, plasma renin activity, and angiotensin II in volunteers. *Hypertension* 21, 704–713. doi: 10.1161/01.HYP.21.5.704
- Hart, E. C. J., and Charkoudian, N. (2014). Sympathetic neural regulation of blood pressure: influences of sex and aging. *Physiology* 29, 8–15. doi: 10.1152/physiol.00031.2013
- Javaheri, S. (2006). Acetazolamide improves central sleep apnea in heart failure: a double-blind, prospective study. *Am. J. Respir. Crit. Care Med.* 173, 234–237. doi: 10.1164/rccm.200507-1035OC
- Jones, M. R., Sealey, J. E., and Laragh, J. H. (2007). Effects of angiotensin receptor blockers on ambulatory plasma renin activity in healthy, normal subjects during unrestricted sodium intake. *Am. J. Hypertens.* 20, 907–916. doi: 10.1016/j.amjhyper.2007.04.009
- Jouett, N. P., Morales, G., Raven, P. B., and Smith, M. L. (2017). Losartan reduces the immediate and sustained increases in muscle sympathetic nerve activity after hyperacute intermittent hypoxia. *J. Appl. Physiol.* 122, 884–892. doi: 10.1152/japplphysiol.00683.2016
- Kählin, J., Mkrtchian, S., Ebberyd, A., Hammarstedt-Nordenvall, L., Nordlander, B., Yoshitake, T., et al. (2014). The human carotid body releases acetylcholine, ATP and cytokines during hypoxia. *Exp. Physiol.* 99, 1089–1098. doi: 10.1113/expphysiol.2014.078873
- Kuznetsova, A., Brockhoff, P. B., and Christensen, R. H. B. (2017). lmerTest package: tests in linear mixed effects models. *J. Stat. Softw.* 82, 1–26. doi: 10.18637/jss.v082.i13
- Lam, S. Y., Fung, M. L., and Leung, P. S. (2004). Regulation of the angiotensin-converting enzyme activity by a time-course hypoxia in the carotid body. *J. Appl. Physiol.* 96, 809–813. doi: 10.1152/japplphysiol.00684.2003
- Lam, S. Y., and Leung, P. S. (2003). Chronic hypoxia activates a local angiotensin-generating system in rat carotid body. *Mol. Cell. Endocrinol.* 203, 147–153. doi: 10.1016/S0303-7207(03)00087-X
- Lenth, R. (2019). *emmeans: Estimated Marginal Means, aka Least-Squares Means, R package version 1.4.3.01*. Available online at: <https://CRAN.R-project.org/package=emmeans>
- Leung, P. S., Lam, S. Y., and Fung, M. L. (2000). Chronic hypoxia upregulates the expression and function of AT(1) receptor in rat carotid body. *J. Endocrinol.* 167, 517–524.
- Li, Y., Xia, X., Zheng, H., Gao, L., Li, Y., and Liu, D. (2006). Angiotensin II enhances carotid body chemoreflex control of sympathetic outflow in chronic heart failure rabbits. *Cardiovasc. Res.* 71, 129–138. doi: 10.1016/j.cardiores.2006.03.017
- Li, Y.-L., and Schultz, H. D. (2006). Enhanced sensitivity of Kv channels to hypoxia in the rabbit carotid body in heart failure: role of angiotensin II. *J. Physiol.* 575, 215–227. doi: 10.1113/jphysiol.2006.110700
- Lombardi, C., Meriggi, P., Agostoni, P., Faini, A., Bilo, G., Revera, M., et al. (2013). High-altitude hypoxia and periodic breathing during sleep: gender-related differences. *J. Sleep Res.* 22, 322–330. doi: 10.1111/jsr.12012
- Marcus, N. J., Li, Y.-L., Bird, C. E., Schultz, H. D., and Morgan, B. J. (2010). Chronic intermittent hypoxia augments chemoreflex control of sympathetic activity: role of the angiotensin II type 1 receptor. *Respir. Physiol. Neurobiol.* 171, 36–45. doi: 10.1016/j.resp.2010.02.003
- Messineo, L., Taranto-Montemurro, L., Azarbarzin, A., Oliveira Marques, M. D., Calianese, N., White, D. P., et al. (2018). Breath-holding as a means to estimate the loop gain contribution to obstructive sleep apnoea. *J. Physiol.* 596, 4043–4056. doi: 10.1113/JP276206
- Mkrtchian, S., Kählin, J., Ebberyd, A., Gonzalez, C., Sanchez, D., Balbir, A., et al. (2012). The human carotid body transcriptome with focus on oxygen sensing and inflammation – a comparative analysis. *J. Physiol.* 590, 3807–3819. doi: 10.1113/jphysiol.2012.231084
- Morgan, B. J., Teodorescu, M., Pegelow, D. F., Jackson, E. R., Schneider, D. L., Plante, D. T., et al. (2018). Effects of losartan and allopurinol on cardiorespiratory regulation in obstructive sleep apnoea. *Exp. Physiol.* 103, 941–955. doi: 10.1113/EP087006
- Ohtawa, M., Takayama, F., Saitoh, K., Yoshinaga, T., and Nakashima, M. (1993). Pharmacokinetics and biochemical efficacy after single and multiple oral administration of losartan, an orally active nonpeptide Angiotensin-II receptor antagonist, in humans. *Br. J. Clin. Pharmacol.* 35, 290–297. doi: 10.1111/j.1365-2125.1993.tb05696.x
- Pialoux, V., Foster, G. E., Ahmed, S. B., Beaudin, A. E., Hanly, P. J., and Poulin, M. J. (2011). Losartan abolishes oxidative stress induced by intermittent hypoxia in humans. *J. Physiol.* 589, 5529–5537. doi: 10.1113/jphysiol.2011.218156
- Richard, N. A., Sahota, I. S., Widmer, N., Ferguson, S., Sheel, A. W., and Koehle, M. S. (2014). Acute mountain sickness, chemosensitivity, and cardiorespiratory responses in humans exposed to hypobaric and normobaric hypoxia. *J. Appl. Physiol.* 116, 945–952. doi: 10.1152/japplphysiol.00319.2013
- Roach, R. C., Bartsch, P., Oelz, O., and Hackett, P. H. (1993). “The Lake Louise Acute Mountain Sickness Scoring System,” in *Hypoxia and Molecular Medicine*, eds J. R. Sutton, C. S. Houston, and G. COates (Burlington, VT: Queen City Press), 272–274.
- Roy, A., Farnham, M. M. J., Derakhshan, F., Pilowsky, P. M., and Wilson, R. J. A. (2018). Acute intermittent hypoxia with concurrent hypercapnia evokes P2X and TRPV1 receptor dependent sensory long-term facilitation in naïve carotid bodies. *J. Physiol.* 596, 3149–3169. doi: 10.1113/JP275001
- Saxena, A., Little, J. T., Nedungadi, T. P., and Cunningham, J. T. (2015). Angiotensin II type 1a receptors in subfornical organ contribute towards chronic intermittent hypoxia-associated sustained increase in mean arterial

- pressure. *Am. J. Physiol. Heart Circ. Physiol.* 308, H435–H446. doi: 10.1152/ajpheart.00747.2014
- Shell, B., Farmer, G. E., Nedungadi, T. P., Wang, L. A., and Marciante, A. B. (2019). Angiotensin type 1a receptors in the median preoptic nucleus support intermittent hypoxia-induced hypertension. *Am. J. Physiol. Integr. Comp. Physiol.* 316, R651–R665.
- Sin, D. D., Fitzgerald, F., Parker, J. D., Newton, G., Floras, J. S., and Bradley, T. D. (1999). Risk factors for central and obstructive sleep apnea in 450 men and women with congestive heart failure. *Am. J. Respir. Crit. Care Med.* 160, 1101–1106. doi: 10.1164/ajrccm.160.4.9903020
- Solaiman, A. Z., Feehan, R. P., Chabitnoy, A. M., Leuenberger, U. A., and Monahan, K. D. (2014). Ventilatory responses to chemoreflex stimulation are not enhanced by angiotensin II in healthy humans. *Auton. Neurosci.* 183, 72–79. doi: 10.1016/j.autneu.2014.01.010
- Solin, P., Roebuck, T., Johns, D. P., Walters, H. E., and Naughton, M. T. (2000). Peripheral and central ventilatory responses in central sleep apnea with and without congestive heart failure. *Am. J. Respir. Crit. Care Med.* 162, 2194–2200. doi: 10.1164/ajrccm.162.6.2002024
- Stephenson, R., Mohan, R. M., Duffin, J., Jarsky, T. I. M. M., Mohan, R. M., Duffin, J., et al. (2000). Circadian rhythms in the chemoreflex control of breathing. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 278, 282–286. doi: 10.1152/ajpregu.2000.278.1R282
- Terrill, P. I., Edwards, B. A., Nemati, S., Butler, J. P., Owens, R. L., Eckert, D. J., et al. (2015). Quantifying the ventilatory control contribution to sleep apnoea using polysomnography. *Eur. Respir. J.* 45, 408–418. doi: 10.1183/09031936.00062914
- Tymko, M. M., Ainslie, P. N., MacLeod, D. B., Willie, C. K., and Foster, G. E. (2015). End tidal-to-arterial CO<sub>2</sub> and O<sub>2</sub> gas gradients at low- and high-altitude during dynamic end-tidal forcing. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 308, R895–R906. doi: 10.1152/ajpregu.00425.2014
- Tymko, M. M., Hoiland, R. L., Kuca, T., Boulet, L. M., Tremblay, J. C., Pinsky, B. K., et al. (2016). Measuring the human ventilatory and cerebral blood flow response to CO<sub>2</sub>: a technical consideration for the end-tidal-to-arterial gas gradient. *J. Appl. Physiol.* 120, 282–296. doi: 10.1152/jappphysiol.00787.2015
- Wang, H. W., Huang, B. S., White, R. A., Chen, A., Ahmad, M., and Leenen, F. H. H. (2016). Mineralocorticoid and angiotensin II type 1 receptors in the subfornical organ mediate angiotensin II – induced hypothalamic reactive oxygen species and hypertension. *Neuroscience* 329, 112–121. doi: 10.1016/j.neuroscience.2016.04.050
- White, D. P. (2005). Pathogenesis of obstructive and central sleep apnea. *Am. J. Respir. Crit. Care Med.* 172, 1363–1370. doi: 10.1164/rccm.200412-1631SO

**Conflict of Interest:** SS has worked as a consultant for Nox Medical, Merck, and Apnimed; he also received grant support from Apnimed and ProSomnus.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Brown, Boulet, Vermeulen, Sands, Wilson, Ayas, Floras and Foster. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.