



# A Cross-Sectional Study of the Distribution Patterns and Potential Determinants in Plasma Selenium Status Among Chinese Adults With Hypertension

Zhuo Wang<sup>1</sup>, Tengfei Lin<sup>1,2</sup>, Yaping Wei<sup>1</sup>, Yun Song<sup>1,3</sup>, Lishun Liu<sup>1</sup>, Ziyi Zhou<sup>1</sup>, Xiao Huang<sup>4</sup>, Ping Chen<sup>5</sup>, Chengzhang Liu<sup>3</sup>, Youbao Li<sup>6</sup>, Binyan Wang<sup>3,6,7</sup>, Jianping Li<sup>8</sup>, Yan Zhang<sup>8</sup>, Yong Huo<sup>8</sup>, Hao Zhang<sup>1</sup>, Xiping Xu<sup>1,3,6</sup>, Xianhui Qin<sup>3,6\*</sup> and Huiyuan Guo<sup>1\*</sup>

## OPEN ACCESS

### Edited by:

Manfred Eggersdorfer,  
University of Groningen, Netherlands

### Reviewed by:

Yoshiro Saito,  
Tohoku University, Japan  
Tommaso Filippini,  
University of Modena and Reggio  
Emilia, Italy

### \*Correspondence:

Xianhui Qin  
pharmaqin@126.com  
Huiyuan Guo  
guohuiyuan@cau.edu.cn

### Specialty section:

This article was submitted to  
Nutritional Epidemiology,  
a section of the journal  
Frontiers in Nutrition

**Received:** 23 February 2022

**Accepted:** 19 April 2022

**Published:** 17 May 2022

### Citation:

Wang Z, Lin T, Wei Y, Song Y,  
Liu L, Zhou Z, Huang X, Chen P,  
Liu C, Li Y, Wang B, Li J, Zhang Y,  
Huo Y, Zhang H, Xu X, Qin X and  
Guo H (2022) A Cross-Sectional  
Study of the Distribution Patterns  
and Potential Determinants in Plasma  
Selenium Status Among Chinese  
Adults With Hypertension.  
*Front. Nutr.* 9:882309.  
doi: 10.3389/fnut.2022.882309

<sup>1</sup> Key Laboratory of Precision Nutrition and Food Quality, Ministry of Education, Department of Nutrition and Health, China Agricultural University, Beijing, China, <sup>2</sup> Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen, China, <sup>3</sup> Institute of Biomedicine, Anhui Medical University, Hefei, China, <sup>4</sup> Department of Cardiology, The Second Affiliated Hospital of Nanchang University, Nanchang, China, <sup>5</sup> College of Pharmacy, Jinan University, Guangzhou, China, <sup>6</sup> National Clinical Research Center for Kidney Disease, State Key Laboratory for Organ Failure Research, Division of Nephrology, Nanfang Hospital, Southern Medical University, Guangzhou, China, <sup>7</sup> Shenzhen Evergreen Medical Institute, Shenzhen, China, <sup>8</sup> Department of Cardiology, Peking University First Hospital, Beijing, China

Selenium (Se) is an essential trace element in selenoproteins biosynthesis for the human body and plays an important role in the prevention and control of subsequent cardiovascular disease in adults with hypertension. However, reports on Se status and its potential determinants in populations from different regions of China are limited, especially data on adults with hypertension, a high-risk group more vulnerable to oxidative stress. Thus, we conducted a cross-sectional study from February 2017 to May 2018 of 2,599 participants (1,389 men and 1,210 women) on middle-aged to elderly adults with hypertension with a mean age of 63.1 years (SD 13.3) from 14 provinces of China and aimed to examine the relationship of plasma Se status with demographic characteristics and lifestyles. Overall, the male participants (mean value 75.0  $\mu\text{g/L}$ ) tended to have higher plasma Se concentrations than the female participants (73.7  $\mu\text{g/L}$ ) when controlling for relevant factors. There were significant differences among regions, and in age and body mass index (BMI) in plasma Se distribution, and plasma Se concentrations were significantly lower among those in the regions with relatively lower Se, aged 60 years or older, and with BMI lower than 28  $\text{kg/m}^2$ . Moreover, a higher frequency of meat consumption (1–2 or  $\geq 3$  times/week vs.  $< 1$  time/week) was significantly associated with higher plasma Se concentrations in men and women, and male alcohol drinkers had significantly higher plasma Se concentrations than non-alcohol drinkers. Adequate consumption of fruits and vegetables (0.5–1.5  $\text{kg/week}$ ) was associated with higher plasma Se concentrations among women, but was associated with relatively lower plasma Se concentrations in men. Our results indicated relatively low plasma Se status in Chinese adults with hypertension from 14 provinces, while

specific factors including geographic, demographic, and lifestyle characteristics and blood pressure were significantly associated with plasma Se status in this hypertensive population. In addition, more studies are required to further evaluate dietary structure and other lifestyle factors that influence circulating Se status.

**Keywords:** selenium, distribution patterns, hypertension, geographical, demographics, lifestyle, nutritional epidemiology

## INTRODUCTION

Selenium (Se) is an essential trace element for human health and plays an important role in the body *via* selenoproteins (1). Se is an integral part of the activation site in the form of selenocysteine (Sec) for glutathione peroxidase (GPX) (2), an enzyme that protects tissues against oxidative stress and catalyzes the reduction of peroxidase following cellular damage. Se can be metabolized to Sec and incorporated into selenoprotein P (SELENOP) (3), which is secreted into the plasma. SELENOP is involved in male fertility, proper brain functioning, and other biological processes. A U-shaped relationship between Se status and chronic disease has been previously reported (4). The different amounts of environmental exposure may have a significant impact on human Se status, especially because dietary Se bioavailability is mostly governed by environmental factors such as Se concentration in the soil, water, and air (5).

A few studies have reported that blood pressure is associated with multiple metal biomarkers (6). However, the relationship between Se and hypertension and subsequent chronic diseases is still inconclusive (7). Our previous findings from the China Stroke Primary Prevention Trial (CSPPT) indicated that there is a lower risk of first stroke in adults with hypertension who have relatively higher plasma Se concentrations, which suggests the importance of maintaining proper nutritional status to prevent cardiovascular diseases (CVDs) (8). Adults with hypertension may be more vulnerable to the effects of oxidative stress, and more sensitive to environmental exposure levels of Se and the antioxidant effects of Se-dependent GPX (9). The prevalence of hypertension in China is high and rising, and a variety of CVDs induced by hypertension have substantial socioeconomic consequences (10). Thus, comprehensively revealing the characteristics and determinants of Se distribution in a population with hypertension, especially in those exposed to various environmental Se levels, will contribute to developing nutrition intervention strategies in populations with hypertension for lowering the disease burden.

China is one of the 40 countries designated by the World Health Organization (WHO) as having low or insufficient levels of Se. Low soil Se levels are widespread, and low dietary sources

of Se and unhealthy dietary habits may raise the potential risk of Se deficiency in the Chinese population. It is especially important that attention be paid to the Se status of Chinese people, given that the soil Se deficiency in China is growing because of climate change (11). People with high blood pressure tend to have lower Se levels than healthy subjects, according to previous studies based on limited sample sizes in China and the former Yugoslavia (12, 13). However, this relationship could differ by location as seen in results from the United States and Denmark where hypertensive individuals were found to have higher Se levels (14, 15). Due to the numerous determinants of Se levels, the Se distribution among adults with hypertension remains unconfirmed. The association of the environment, demographics, and dietary habits with Se status among Chinese adults with hypertension especially needs more evidence. Furthermore, most prior investigations used more conveniently obtained samples to assess Se status, such as nails, hair, or urine (16–18). However, the above indicators do not reflect circulating Se because plasma and serum Se are considered the most suitable indicators in epidemiological studies (19). Although the previous studies have shown adverse effects of Se exposure at low levels, the debate concerning the issue of selenoprotein optimization is still ongoing for the evidence regarding the need to increase Se levels is still controversial (20, 21). In particular, high Se exposure has been shown to be related to an increased risk of type 2 diabetes (22). Thus, more evidence is needed to reveal the optimal Se status for humans. Overall, most of the previous research has indicated a relatively low Se status in Chinese with Keshan disease or Kashin–Beck disease (23). However, there is still a lack of large-scale and representative reports on Se status in Chinese adults with hypertension.

In general, previous studies have not adequately disclosed the relationship between the environment, demographic characteristics, and lifestyle with Se status in Chinese adults with hypertension. Thus, we conducted a cross-sectional survey of randomly selected adults with hypertension in 14 Chinese provinces, using inductively coupled plasma mass spectrometry (ICP–MS) detection methods to characterize plasma Se concentration. By evaluating the distribution of plasma Se status and its potential determinants of Se among adults with hypertension, we aimed to identify populations at high risk for low Se and the potential strategies for the adjustment of Se status.

## MATERIALS AND METHODS

### Study Population

We conducted a multi-centric, non-interventional, observational, real-world study for the identification and

**Abbreviations:**  $\beta$ s, beta coefficients; BMI, body mass index; CHNS, China Health and Nutrition Survey; Cis, confidence intervals; CVD, cardiovascular disease; DBP, diastolic blood pressure; EAR, estimated average requirement; GPX, glutathione peroxidase; ICP–MS, inductively coupled plasma mass spectrometry; NHANES, National Health and Nutrition Examination Survey; PRs, prevalence ratios; ROS, reactive oxygen species; SBP, systolic blood pressure; Se, selenium; SELENOP, selenoprotein P; tHcy, total homocysteine; WHO, World Health Organization; IQR, interquartile range.

registration of a high-risk population with both hypertension and elevated total homocysteine levels (tHcy  $\geq 10 \mu\text{mol/L}$ ) within China, which was initiated in February 2017. Patients were recruited from the community through open recruitment rather than through random selection. The inclusion criteria included the following individuals: (1) Individuals who had hypertension, defined as seated, resting systolic blood pressure (SBP) of 140 mm Hg or higher, or diastolic blood pressure (DBP) of 90 mm Hg or higher at the recruitment visit, or who were taking an antihypertensive medication, according to the diagnostic criteria of the 2010 Chinese guidelines for the management of hypertension (24); (2) Individuals who had elevated total homocysteine (tHcy  $\geq 10 \mu\text{mol/L}$ ); and (3) individuals who gave signed, written, and informed consent.

Of the eligible participants in this observational study, two subsamples were selected, without duplication, at 10 and 16 months after the study began, in December 2017 and June 2018, respectively. First, 900 participants were randomly selected from nine provinces and stratified by province, who were enrolled from June to August 2017 and had complete screening records (physical exam, questionnaire, and biological samples). Second, another 1,709 participants were randomly selected from 14 provinces (including the 9 provinces in the first sampling set plus an additional 5 provinces) and stratified by province, sex, and age groups, who were enrolled from February 2017 to May 2018 and had complete screening records (25). Finally, after combining the two subsamples, a total of 2,599 participants from all 14 provinces without duplication, were included in our current study, excluding those with missing values of Se (Supplementary Figure 1).

The study protocol was approved by the Ethics Committee of Peking University First Hospital, Beijing, China (Approval No.: 20161231), and the study was conducted according to the Declaration of Helsinki. All methods were performed in accordance with the relevant guidelines and regulations. Written, informed consent was obtained from all the participants.

## Blood Sample Collection and Laboratory Assays

A fasting, venous blood sample was obtained from each participant. Plasma samples were separated within 30 min of collection and stored at  $-80^\circ\text{C}$ . Plasma Se concentrations of the randomly selected participants were measured by ICP-MS using Thermo Fisher iCAP Q ICPMS, in a commercial lab (Beijing DIAN Medical Diagnostics Laboratory, Beijing, China). Both intra-assay and inter-assay coefficients of variation (CV) for duplicate samples (randomly placed among the study sample) were calculated. The intra-assay CV of plasma Se ranged from 2.14 to 9.38%, while the inter-assay ranged from 2.41 to 3.31%.

## Major Definitions

The 14 provinces were stratified by soil Se concentrations based on reports from previous studies, and these were then categorized into three levels: Se-marginal (0.125– $<0.175 \text{ mg/kg}$ ),

Se-sufficient (0.175 to  $<0.40 \text{ mg/kg}$ ), and Se-rich (0.40– $3.0 \text{ mg/kg}$ ) (26, 27). The Se-marginal regions in China for this study included four study provinces (Gansu, Heilongjiang, Liaoning, and Shandong), while Se-sufficient regions included eight study provinces (Anhui, Beijing, Hebei, Jiangsu, Ningxia, Yunnan, Shanxi, and Sichuan), and Se-rich regions included two study provinces (Guangxi and Hunan).

## Statistical Analysis

For baseline characteristics and plasma Se concentrations, normally distributed and approximately normally distributed continuous variables were presented as mean  $\pm$  SD and were compared using *t*-tests, and non-normally distributed continuous variables were presented as median (25th–75th percentile) and were compared using the Wilcoxon–Rank–Sum tests (for two factors) or the Kruskal–Wallis tests (for more than two factors). Categorical variables were presented as numbers (percentage) and were compared using the  $\chi^2$  (for two factors) tests or the ANOVA tests (for more than two factors). Pairwise comparisons of average plasma Se concentrations in each group were conducted by least significant difference (LSD) tests.

Adjusted Se values were calculated by general linear regression models after controlling for sex, age, body mass index (BMI), smoking status (never, past, current: defined as having smoked  $\geq 1$  cigarette per day or  $\geq 18$  packs in the past year), alcohol drinking status (never, past, current: defined as drinking alcohol at least 2 times per week in the past year), region (Se-marginal, Se-sufficient, and Se-rich provinces in China), SBP, DBP, history of hypertension (no, yes), antihypertensive drug use (no, yes), multivitamin use (no, yes), meat consumption ( $<1$  time/week, 1–3 times/week, 3–5 or more times/week), and consumption of fruits and vegetables ( $<0.5 \text{ kg/week}$ , 0.5–1.5 kg/week,  $>1.5 \text{ kg/week}$ ). In addition, the linear trends in Se concentration across age, BMI, region, smoking, alcohol drinking, meat consumption, and fruit and vegetable consumption groups were assessed using general linear regression models.

Regression models were estimated *via* ordinary least squares and restricted cubic-spline functions were used to assess non-linear trends with three knots for continuous variables stratified by sex. Beta coefficients ( $\beta$ s) and 95% confidence intervals (CIs) of Se concentrations in association with selected demographic variables were estimated using linear regression models, without or with adjustment for potential confounding factors.

The map of provincial distribution patterns of the prevalence of plasma Se was plotted using R software, version 3.6.1.<sup>1</sup> The SHP file of the China map was obtained from the website of the Resource and Environment Data Cloud Platform.<sup>2</sup>

A two-tailed  $p < 0.05$  was considered statistically significant in all analyses. R soft-ware, version 3.6.1 (see text footnote 1) and Empower version 2.0<sup>3</sup> were used for all statistical analyses and map plotting.

<sup>1</sup><http://www.R-project.org/>

<sup>2</sup><http://www.resdc.cn>

<sup>3</sup><http://www.empowerstats.com>

## RESULTS

### Study Participants and Baseline Characteristics

As illustrated in the flowchart of the study participants (**Supplementary Figure 1**), the current study included a total of 2,599 participants, among which 736 (28.3%) were from Se-marginal regions of China, 1,463 (56.3%) were from Se-sufficient regions of China, and 400 (15.4%) were from Se-rich regions of China (**Table 1**).

Overall, this study included relatively more men (53.4%), elderly participants aged 60 years or above (59.4%), and those with normal BMI (42.8%) among all the participants (**Supplementary Table 1**). Comparisons of the baseline characteristics of the participants randomly selected in the current analysis and those not included, showed no obvious differences, and are presented in **Supplementary Table 2**. The participant characteristics by region are listed in **Table 1**. The participants in Se-sufficient regions and Se-rich regions had lower baseline blood pressure (systolic and diastolic) and fewer were current smokers and alcohol drinkers, who had lower BMI and a higher history of hypertension than those from Se-marginal regions. The participants from Se-rich regions had higher frequencies of meat consumption and fruit and vegetable consumption than those from Se-marginal regions and Se-sufficient regions. Moreover, multivitamin use was low across all regions. Overall, compared with the participants from Se-marginal regions, significantly higher average plasma Se concentrations were found in the participants from Se-sufficient regions and Se-rich regions ( $p < 0.05$ ) after adjustment for the related factors. Moreover, the participant characteristics stratified by the 14 study provinces and sex are listed in **Supplementary Tables 1, 3**. The mean  $\pm$  SD values of adjusted plasma Se concentrations were higher in the male participants ( $75.0 \pm 19.4$ ,  $\mu\text{g/L}$ ) than in the female participants ( $73.7 \pm 19.4$ ,  $\mu\text{g/L}$ ) (**Supplementary Table 3**). Accordingly, as shown in **Supplementary Figure 2**, compared with women, there was a shift toward the right in the distribution of plasma Se concentration in males.

### Demographic and Lifestyle Distribution Patterns of Plasma Selenium Status

Overall, there were significant regional, sex, age, and BMI differences in plasma Se status among the participants with hypertension from 14 provinces of China. The regional distribution pattern of plasma Se status is presented in **Figure 1**. In provinces of Se-marginal regions, including Gansu, Heilongjiang, Liaoning, and Shandong, participants had relatively lower mean values of plasma Se concentrations, ranging from 58.3 (Gansu) to 76.1 (Shandong)  $\mu\text{g/L}$ . For those living in Se-sufficient regions (Anhui, Beijing, Hebei, Jiangsu, Ningxia, Shanxi, Sichuan, and Yunnan), relatively low plasma Se concentrations (mean values ranging from 67.3 for Yunnan to 83.2 for Jiangsu,  $\mu\text{g/L}$ ) were found. In addition, significantly higher plasma Se concentrations were found in Se-rich regions, including Guangxi (mean value, 81.7  $\mu\text{g/L}$ ) and

Hunan (mean value, 88.6  $\mu\text{g/L}$ ) (**Supplementary Table 1**). The results comparing plasma Se concentrations among different sexes, age groups, and BMI groups were also explored in this study. After controlling for several potential determinants, including demographic characteristics and lifestyle, men tended to have relatively higher average plasma Se concentrations than females ( $p = 0.082$ ) (**Supplementary Table 3**). The differences in plasma Se status between men and women by province, age, and BMI is also shown (**Figure 2** and **Supplementary Table 4**). An age pattern in regards to plasma Se distribution, was found with a significant decline in Se concentration by age (men: for trend,  $p < 0.001$ , for non-linearity,  $p = 0.784$ ; women: for trend,  $p < 0.001$ , for non-linearity,  $p = 0.160$ ) (**Figure 2A** and **Table 2**). Moreover, plasma Se concentrations increased with elevated BMI, especially in men (for trend = 0.023, for non-linearity,  $p = 0.209$ ) (**Figure 2B** and **Table 2**).

The lifestyle distribution patterns of plasma Se status among men and women are shown in **Figure 3**. Current male alcohol drinkers had significantly higher average plasma Se concentrations than male non-alcohol drinkers ( $p < 0.05$ ). However, current male smokers tended to have lower average plasma Se concentrations than male non-smokers ( $p = 0.177$ ) (**Figure 3A** and **Supplementary Table 5**). Moreover, the average plasma Se concentrations significantly increased with higher frequency in meat in both men and women (for trend,  $p < 0.05$ ) (**Figure 3B** and **Table 3**).

### Association of Demographic Characteristics and Lifestyle With Plasma Selenium Status

The multivariate regression analysis showed that, for the male participants, after adjusting for relative factors, the lower plasma Se concentrations were significantly associated with age  $\geq 70$  years ( $p < 0.001$ ), being from Se-sufficient regions ( $p < 0.001$ ) and Se-marginal regions ( $p < 0.001$ ), and having BMI  $< 24$   $\text{kg/m}^2$  ( $p = 0.022$ ) (**Table 2**). Moreover, for males, relatively lower plasma Se concentrations were associated with current smoking ( $p = 0.043$ ) and consumption of fruits and vegetables  $> 1.5$   $\text{kg/week}$  ( $p = 0.044$ ), whereas with current alcohol drinking ( $p < 0.001$ ), consuming meat 1–2 times/week ( $p = 0.020$ ) and consuming meat 3–5 or more times/week ( $p < 0.001$ ) were associated with relatively higher plasma Se concentrations (**Table 3**). For the female participants, the lower plasma Se concentrations were associated with age  $\geq 70$  years ( $p < 0.001$ ) and being from Se-sufficient regions ( $p < 0.001$ ) and Se-marginal regions ( $p < 0.001$ ) (**Table 2**). Additionally, relatively higher plasma Se concentrations were associated with consuming meat 1–2 times/week ( $p = 0.017$ ), consuming meat 3–5 or more times/week ( $p = 0.002$ ) and consuming fruits and vegetables 0.5–1.5  $\text{kg/week}$  ( $p = 0.039$ ) (**Table 3**). Furthermore, plasma Se concentrations were inversely associated with baseline SBP (per SD increment,  $p = 0.008$ ; for non-linearity,  $p = 0.312$ ) and baseline DBP (per SD increment,  $p = 0.002$ ; for non-linearity,  $p = 0.757$ ) in the male participants. For the female participants, plasma Se concentrations were inversely associated with baseline DBP

**TABLE 1** | Baseline characteristics of the study participants stratified by region.

Characteristics	Se-marginal region <sup>1</sup>	Se-sufficient region	Se-rich region
<i>N</i>	736	1,463	400
Male, <i>n</i> (%)	383 (52.0)	779 (53.2)	227 (56.8)
<b>Age, years</b>			
<60	306 (41.6)	587 (40.1)	163 (40.8)
60 to <70	178 (24.2)	373 (25.5)	113 (28.2)
≥70	252 (34.2)	503 (34.4)	124 (31.0)
<b>Body mass index, kg/m<sup>2</sup></b>			
<24	242 (32.9)	550 (37.6)*	202 (50.5) ††
24 to <28	317 (43.1)	648 (44.3)	147 (36.8)
≥28	177 (24.0)	265 (18.1)	51 (12.8)
<b>Smoking status, <i>n</i> (%)</b>			
Never	528 (71.7)	1,006 (68.8)**	286 (71.5)
Past	42 (5.7)	202 (13.8)	27 (6.8)
Current	166 (22.6)	255 (17.4)	87 (21.8)
<b>Alcohol drinking status, <i>n</i> (%)</b>			
Never	564 (76.6)	1072 (73.3)*	312 (78.0) †
Past	32 (4.3)	126 (8.6)	33 (8.2)
Current	140 (19.0)	265 (18.1)	55 (13.8)
SBP, mm Hg	147.0 (135.0–156.1)	139.3 (130.0–150.0) **	143.2 (133.0–153.3) †
DBP, mm Hg	90.0 (80.7–97.0)	85.7 (78.7–92.3) **	86.8 (81.3–92.3) †
Antihypertensive drug use, <i>n</i> (%)	557 (75.7)	1,102 (75.3)	99 (24.8) ††
Multivitamin use, <i>n</i> (%)	52 (7.1)	148 (10.1) *	9 (2.2) ††
History of hypertension, <i>n</i> (%)	592 (80.4)	1423 (97.3) **	352 (88.0) †
<b>Meat consumption frequency, <i>n</i> (%)</b>			
<1 time/week	183 (24.9)	326 (22.3)**	84 (21.0) ††
1–2 times/week	273 (37.1)	596 (40.7)	94 (23.5)
3–5 or more times/week	280 (38.0)	541 (37.0)	222 (55.5)
<b>Fruits and vegetables consumption, <i>n</i> (%)</b>			
<0.5 kg/week	183 (24.9)	326 (22.3)	84 (21.0) ††
0.5–1.5 kg/week	273 (37.1)	596 (40.7)	94 (23.5)
>1.5 kg/week	280 (38.0)	541 (37.0)	222 (55.5)
Se, μg/L	69.3 ± 20.7	74.0 ± 19.3 **	85.1 ± 18.5 ††

For continuous variables, the values are presented as median interquartile range (IQR) or mean ± SD.

<sup>1</sup>Se-marginal areas in China include four provinces (Gansu, Heilongjiang, Liaoning, and Shandong), Se-sufficient areas in China include eight provinces (Anhui, Beijing, Hebei, Jiangsu, Ningxia, Shanxi, Sichuan, and Yunnan), and Se-rich areas in China include two provinces (Guangxi and Hunan).

<sup>2</sup>Plasma Se concentrations were adjusted for sex, age, BMI, SBP, DBP, history of hypertension, antihypertensive drug use, multivitamin use, smoking, alcohol drinking, meat consumption, and consumption of fruits and vegetables.

\* $p < 0.05$ , \*\* $p < 0.001$ ; the values significantly differed among the participants from Se-marginal and Se-sufficient areas in China.

† $p < 0.05$ , †† $p < 0.001$ ; the values significantly differed among the participants from Se-marginal and Se-rich areas in China.


(per SD increment,  $p = 0.010$ ; for non-linearity,  $p = 0.428$ ) (Supplementary Table 6).

## DISCUSSION

To our knowledge, our study presents the first nationwide plasma Se distribution obtained using the ICP-MS method in both middle-aged and elderly adults with hypertension in China. Our research provides two new insights. First, in this population, low circulating Se status was remarkably widespread, and plasma Se status varied significantly according to the region, sex, age, and BMI. Second, the lifestyle characteristics including smoking, alcohol drinking, and consumption of meat, fruits, or vegetables all had a close relationship with plasma Se status and may be

potential intervenable factors for improvement of circulating Se status in adults with hypertension.


As Se is a trace element with a narrow normal range, maintaining a dynamic balance of Se distribution is imperative to human antioxidant function (1). Relatively high circulating Se concentrations (>120 μg/L on average) have been found among populations in the United States, Japan, Canada, etc. (28, 29). In contrast, the low circulating Se status and multielement imbalances were found to be associated with Keshan disease or Kashin-Beck disease in the northeast and northwest regions of China (21, 30, 31). However, a study conducted among 10 provinces and municipalities of China assessing hair Se concentrations in Chinese inhabitants across the northeast to southeast China indicated that about 84% of all residents had normal hair Se content. The study (32) also discovered



sex and geographical differences in hair Se status. Moreover, recent evidence based on the 2015 China Health and Nutrition Survey (CHNS) showed that close to 60% of men's daily intake of Se is insufficient [not reaching the recommended amount of the estimated average requirement (EAR)] (33). Other studies have focused on Se status among people with various diseases, such as cancer, CVD, diabetes, etc., revealing the differences in Se distribution compared to healthy people (34–36). Overall, the previous studies have examined Se levels and Se intake of people in specific parts of China (37–39), but these studies are limited by the available materials used to measure Se, such as hair, nails, and urine, which do not

reflect Se circulation status in the body (19). While a few studies on blood Se status have been conducted, they have been limited in terms of representativeness and sample size (28). As far as we know, there has been no report on the status of circulating Se levels in a large and representative Chinese population with hypertension that is exposed to diverse environmental Se levels.

We found relatively low Se levels and significant regional differences in Se levels in Chinese adults with hypertension. Hypertensive patients may encounter oxidative stress more frequently than healthy subjects and may need more support from nutrients, such as alpha tocopherol (Vit E), ascorbic acid



(Vit C), Se, etc., to lower the concentration of reactive oxygen species (ROS). Thus, the relatively low Se levels among these patients may be related to a higher Se demand and consumption based on their special physical conditions (9, 40). Furthermore, the interaction between the origin of foods consumed, dietary habits, and the environment on human Se status could be crucial. The level of Se in food is mostly determined by the amount of Se in the soil. Se cannot be synthesized in the human body. Consequently, the source for Se in the human body almost entirely originates from food intake (41). Cereals, fish, eggs, and meats are well known as the major dietary sources of Se. However, the Se content of the same food was found to be lower in China, compared with that of America, Japan, Canada, etc. (42). Additionally, much of the Chinese population consume more cereals, fruits, or vegetables than they do seafood, fish, eggs, or meat (the main Se-rich food in China) in daily life (43). Thus, we speculated that it was

Se-rich food bioavailability, as indicated by environmental Se exposure, along with diverse dietary habits that led to the varied distribution of human plasma Se regionally. From what we have found, the average plasma Se levels of participants in Se-marginal regions were lower, while those in Se-rich regions were higher. It is worth noting that even in Se-sufficient regions, the Se status among participants was not greatly improved, and the Se insufficiency rate exceeded 50%, indicating that Se levels are not only affected by soil Se. Ecological factors, dietary intake, and differences in body metabolism may also be important influences that jointly determine individual Se levels (44).

In this study, it was found that the plasma Se levels of men were slightly higher than that of women, even when the potential effects of other factors were excluded, which was consistent with some previous studies (45, 46), but not with all of them (47). Sexual dimorphism may partially explain the differences observed among males and females regarding Se status (48). This is of fundamental importance, and sex differences should be considered when analyzing the Se status. Se has indeed been demonstrated to play a direct role in testosterone synthesis and in protecting the testes against toxins (49). It has been consistently shown that Se directly influences testosterone production from early development and it may also indirectly enhance spermatogenesis through its regulation of testosterone production (50). To achieve normal physiological functions, men require more Se than women, especially since the retention rate for Se is prominent in the testes (51). This could be one of the key explanations for the Se disparity between men and women. In addition, even when exposed to similar environmental Se levels, the sex differences in Se levels may also be determined by dietary habits and lifestyles, which deserves more relevant research in the future.

With regard to aging, the decline in Se levels in Chinese adults with hypertension was noticeable. The impact of age on Se status needs to be completely considered from two aspects. Less Se-rich food intake could be the first main reason for the poor Se status among the elderly. A nutritional health survey conducted among the Chinese elderly found that residents living in inland areas consumed saltwater fish less than one time per month (52). A large number of elderly Chinese adults tend to be vegetarian, which is detrimental to adequate Se intake. Moreover, the decrease in absorption and metabolism function could partly explain the poor Se status in elderly participants (53).

Higher BMI may be associated with higher plasma Se levels. The plasma Se concentrations of participants with BMI over 28  $\text{kg/m}^2$  were greater than that of participants with BMI lower than 24  $\text{kg/m}^2$ . The trend remained after adjustment for relevant variables. Previous research findings on the link between BMI and serum Se have been inconsistent. A recent study based on the U.S. Third National Health and Nutrition Examination Survey (NHANES III) found a negative correlation between BMI and serum Se concentrations (46). However, in a nationally representative sample of 1,045 British adults, and in a cohort of 3,387 Danish men aged 53–74, BMI was not associated with circulating Se concentrations (15, 47). The extraordinary effect revealed by BMI in our findings should not

**TABLE 2** | The  $\beta$  and 95% CI of plasma Se concentrations by demographic factors in Chinese middle-aged and elderly adults with hypertension stratified by sex.

Variables	N	Plasma Se, $\mu$ g/L	Crude Model*		Adjusted Model†	
			$\beta$ (95% CI)	p	$\beta$ (95% CI)	p
<b>Male</b>						
<b>Age, years</b>						
<60	597	79.4 $\pm$ 19.0	0.00 (Ref)		0.00 (Ref)	
60 to <70	353	76.1 $\pm$ 20.3	-3.35 (-6.08, -0.62)	0.016	-2.30 (-5.02, 0.42)	0.097
$\geq$ 70	439	71.2 $\pm$ 23.3	-8.17 (-10.73, -5.61)	<0.001	-6.65 (-9.51, -3.78)	<0.001
$\rho$ , for trend				<0.001		<0.001
<b>Region</b>						
Se-rich	227	89.3 $\pm$ 18.5	0.00 (Ref)		0.00 (Ref)	
Se-sufficient	779	75.1 $\pm$ 18.7	-14.15 (-17.12, -11.18)	<0.001	-15.02 (-18.29, -11.75)	<0.001
Se-marginal	383	69.8 $\pm$ 23.4	-19.50 (-22.79, -16.20)	<0.001	-19.82 (-23.44, -16.21)	<0.001
$\rho$ , for trend				<0.001		<0.001
<b>BMI, kg/m<sup>2</sup></b>						
$\geq$ 28	267	78.2 $\pm$ 20.7	0.00 (Ref)		0.00 (Ref)	
24 to <28	624	76.4 $\pm$ 20.2	-1.82 (-4.83, 1.19)	0.237	-2.09 (-4.94, 0.76)	0.150
<24	498	74.2 $\pm$ 22.1	-3.99 (-7.12, -0.87)	0.012	-3.59 (-6.67, -0.50)	0.022
$\rho$ , for trend				0.009		0.023
<b>Female</b>						
<b>Age, years</b>						
<60	459	75.0 $\pm$ 19.9	0.00 (Ref)		0.00 (Ref)	
60 to <70	311	72.9 $\pm$ 22.9	-2.11 (-5.14, 0.91)	0.170	-2.65 (-5.56, 0.25)	0.073
$\geq$ 70	440	69.7 $\pm$ 20.6	-5.33 (-8.07, -2.58)	<0.001	-4.79 (-7.59, -2.00)	<0.001
$\rho$ , for trend				<0.001		<0.001
<b>Region</b>						
Se-rich	173	88.4 $\pm$ 16.3	0.00 (Ref)		0.00 (Ref)	
Se-sufficient	684	72.7 $\pm$ 20.2	-15.71 (-19.01, -12.41)	<0.001	-15.92 (-19.60, -12.24)	<0.001
Se-marginal	353	64.5 $\pm$ 20.4	-23.95 (-27.54, -20.35)	<0.001	-21.77 (-25.81, -17.72)	<0.001
$\rho$ , for trend				<0.001		<0.001
<b>BMI, kg/m<sup>2</sup></b>						
$\geq$ 28	226	74.2 $\pm$ 19.7	0.00 (Ref)		0.00 (Ref)	
24 to <28	488	71.8 $\pm$ 20.5	-2.40 (-5.72, 0.93)	0.237	-2.96 (-6.06, 0.14)	0.061
<24	496	72.6 $\pm$ 22.3	-1.65 (-4.96, 1.67)	0.012	-3.07 (-6.22, 0.08)	0.056
$\rho$ , for trend				0.492		0.093

For continuous variables, the values are presented as mean  $\pm$  SD.


<sup>1</sup>Se-marginal areas in China include four provinces (Gansu, Heilongjiang, Liaoning, and Shandong), Se-sufficient areas in China include eight provinces (Anhui, Beijing, Hebei, Jiangsu, Ningxia, Shanxi, Sichuan, and Yunnan), and Se-rich areas in China include two provinces (Guangxi and Hunan).

\*Values for  $\beta$  and 95% CI are from multilinear regression analysis.

†If not stratified, the model was adjusted for sex, age, region (Se-marginal, Se-sufficient, and Se-rich areas), BMI, SBP, DBP, history of hypertension, antihypertensive drug use, multivitamin use, smoking, alcohol drinking, meat consumption, and consumption of fruits and vegetables.

be ignored. The current population of elderly people in China has suffered through difficult periods during the establishment and development of modern China. In the initial stages, their amount of primary food intake was insufficient, rendering a focus on their nutritional status to be irrelevant. However, after the food scarcity problem was completely resolved, more attention has been drawn to human health, especially on nutrition. Therefore, people with higher BMI tend to have higher Se levels, which may be linked to high-quality food intake and social-economic status. In general, the relationship between BMI and Se status needs to be verified by more large-scale studies, and our study provides a certain extent of evidence for the Se distribution of hypertensive adults with different BMI levels in China.

Although the relationship between Se and blood pressure or hypertension is still inconclusive, we found plasma Se to be inversely associated with baseline SBP and DBP. Our findings were consistent with that of some previous case-control studies (13, 23). However, the results based on cross-sectional studies are still controversial. One study that included 722 Finnish male adults also found a significant inverse relation between Se and SBP (54), while some other studies in 2,638 middle-aged Americans, 3,041 Danish elderly adults, and 680 Indian adults found a positive association between Se and SBP (14, 15, 55). Moreover, most other studies did not find a significant relationship between Se and blood pressure (7). However, more studies are needed to verify our results and to further examine the biological mechanisms underlying the associations.



The close relation between circulating Se status with lifestyle and dietary habits including smoking, alcohol drinking, and meat, fruit, and vegetable consumption, was also found among adults with hypertension. Our finding that male smokers tended to have relatively lower plasma Se levels compared to male non-smokers is consistent with previous results. A study conducted among 44 adults in a seleniferous area of America indicated that smokers had lower tissue Se concentrations compared to non-smokers due, at least in part, to lower Se intake (56). However, there is other evidence that smokers may have reduced food intake and tend to select diets of low nutrient density. Male smokers tended to have a higher frequency of meat and fruit and vegetable consumption and lower BMI than non-smokers in the current study, suggesting that the lower Se concentrations of smokers may be determined by multiple factors. Alcohol intake was found to be associated with elevated serum Se levels in 1,183 Japanese and 124 Americans (57, 58). Similarly, we discovered that male alcohol drinkers had considerably higher

plasma Se levels. The results from NHANES 2003–2008 showed that men consumed an excess of non-alcohol sourced energy on drinking days relative to non-drinking days. Specifically, men consumed more from the meat, poultry, and fish group, and in particular, more meat (59). Accordingly, male alcohol drinkers in the current study, tended to consume more meat, suggesting that the higher Se concentrations observed in male alcohol drinkers may be partly explained by their relatively higher Se-rich food intake (Supplementary Table 5). The typical food sources with the highest Se concentrations (mg/kg) are seafood and organ meats, followed by muscle meats and crops, dairy products and milk, grain and cereals, fruits, and vegetables (42). As expected, in our study, higher circulating Se levels were found in men or women with the greatest consumption of meat. However, greater consumption of fruits and vegetables seemed to be associated with relatively lower plasma Se levels for men, which may relate to the low Se content in fruits and vegetables and personal dietary structures.

**TABLE 3** | The  $\beta$  and 95% CI of plasma Se concentrations by lifestyle and dietary habits in Chinese middle-aged and elderly adults with hypertension stratified by sex<sup>1</sup>.

Variables	N	Plasma Se, $\mu$ g/L	Crude Model *		Adjusted Model <sup>†</sup>	
			$\beta$ (95% CI)	p	$\beta$ (95% CI)	p
<b>Male</b>						
<b>Smoking status</b>						
Never	1820	74.0 $\pm$ 21.1	0.00 (Ref)		0.00 (Ref)	
Past	271	74.7 $\pm$ 22.2	-0.83 (-3.90, 2.23)	0.594	-1.61 (-4.74, 1.52)	0.313
Current	508	75.5 $\pm$ 20.7	0.50 (-1.97, 2.98)	0.690	-2.53 (-4.98, -0.08)	0.043
$\rho$ , for trend				0.726		0.041
<b>Alcohol drinking status</b>						
Never	1948	73.3 $\pm$ 21.3	0.00 (Ref)		0.00 (Ref)	
Past	191	74.8 $\pm$ 19.6	-0.21 (-3.62, 3.20)	0.905	-0.39 (-3.85, 3.08)	0.826
Current	460	78.8 $\pm$ 20.6	4.69 (2.23, 7.15)	<0.001	4.44 (1.97, 6.91)	<0.001
$\rho$ , for trend				<0.001		<0.001
<b>Meat consumption frequency</b>						
<1 time/week	252	70.4 $\pm$ 20.8	0.00 (Ref)		0.00 (Ref)	
1-2times/week	447	73.5 $\pm$ 23.5	3.08 (-0.12, 6.28)	0.059	3.64 (0.55, 6.73)	0.020
3-5 or more times/week	690	79.6 $\pm$ 18.7	9.18 (6.19, 12.17)	<0.001	5.31 (2.29, 8.33)	<0.001
$\rho$ , for trend				<0.001		<0.001
<b>Fruits and vegetables consumption</b>						
<0.5 kg/week	320	76.8 $\pm$ 23.7	0.00 (Ref)		0.00 (Ref)	
0.5-1.5 kg/week	487	74.2 $\pm$ 21.1	-2.63 (-5.60, 0.33)	0.081	-2.29 (-5.12, 0.54)	0.112
>1.5 kg/week	582	77.0 $\pm$ 19.3	0.20 (-2.66, 3.07)	0.890	-2.85 (-5.63, -0.07)	0.044
$\rho$ , for trend				0.583		0.056
<b>Female</b>						
<b>Meat consumption frequency</b>						
<1 time/week	369	67.4 $\pm$ 19.7	0.00 (Ref)		0.00 (Ref)	
1-2times/week	442	71.8 $\pm$ 20.5	4.36 (1.50, 7.21)	0.002	3.35 (0.60, 6.11)	0.017
$\geq$ 3 times/week	399	78.2 $\pm$ 21.8	10.07 (6.22, 13.92)	<0.001	4.74 (1.74, 7.74)	0.002
$\rho$ , for trend				<0.001		0.001
<b>Fruits and vegetables consumption</b>						
<0.5 kg/week	273	69.3 $\pm$ 18.8	0.00 (Ref)		0.00 (Ref)	
0.5-1.5 kg/week	476	72.9 $\pm$ 22.1	3.60 (0.47, 6.73)	0.024	3.17 (0.16, 6.18)	0.039
>1.5 kg/week	461	74.1 $\pm$ 21.1	4.75 (1.60, 7.90)	0.003	1.86 (-1.22, 4.93)	0.237
$\rho$ , for trend				0.004		0.384

For continuous variables, the values are presented as mean  $\pm$  SD.

<sup>1</sup>Se-marginal areas in China include four provinces (Gansu, Heilongjiang, Liaoning, and Shandong), Se-sufficient areas in China include eight provinces (Anhui, Beijing, Hebei, Jiangsu, Ningxia, Shanxi, Sichuan, and Yunnan), and Se-rich areas in China include two provinces (Guangxi and Hunan).

\*Values  $\beta$  and 95% CI are from multilinear regression analysis.

<sup>†</sup>If not stratified, model was adjusted for sex, age, region (Se-marginal, Se-sufficient, and Se-rich areas), BMI, SBP, DBP, history of hypertension, antihypertensive drug use, multivitamin use, smoking, alcohol drinking, meat consumption, and consumption of fruits and vegetables.

Several limitations of this study merit consideration when interpreting our conclusions. First, our study did not cover all 31 provinces throughout the country. However, the study population was from 14 populated provinces in China, each with diverse Se content in the soil. The results should be interpreted with caution due to the small sample size in some categories when the data were stratified by sex and soil Se distribution. Second, we lacked detailed data on potential confounding factors such as estimated dietary Se information, Se concentrations of food groups, information on Se supplementation, etc. Third, the results of this study were based on a population of middle-aged and elderly Chinese adults with hypertension, thus the results might not be representative of other ethnic groups. Fourth, due to limitations of the initial design, we were not

able to test the data of GPX activity and SELENOP, which are important functional biomarkers of Se. Further characterization of these biomarkers will aid in revealing the functional part of Se status in adults with hypertension. Moreover, the type of Se compound is now considered relevant in the assessment of its relation to human health and diseases (60-62). Thus, the determination of different forms of Se, such as inorganic selenium and organic Se, would help to better assess Se status and its association with related diseases in future studies. Owing to these limitations, our findings need to be confirmed in future studies with larger sample sizes and more Se-related biomarkers. In addition, studies are required to further evaluate dietary structures and other lifestyle factors that influence circulating Se status.

## CONCLUSION

In conclusion, using a validated ICP-MS method, we found significant evidence for low plasma Se concentrations and a high prevalence of Se insufficiency among Chinese adults with hypertension. Women, the elderly (age  $\geq 60$  years), those with lower BMI ( $<28 \text{ kg/m}^2$ ), and those living in Se-marginal and Se-sufficient regions of China had significantly lower plasma Se levels. Our findings revealed the importance of environmental, demographic, and lifestyle factors on human Se status. Additionally, we found that proper alcohol intake and increased consumption of Se-rich food (such as meat) may contribute to helping maintain relatively higher circulating Se levels among Chinese middle-aged and elderly patients with hypertension to meet the demands for antioxidation.

## DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

## ETHICS STATEMENT

The studies involving human participants were reviewed and approved by the Ethics Committee of Peking University First Hospital, Beijing, China. The patients/participants provided their written informed consent to participate in this study.

## AUTHOR CONTRIBUTIONS

ZW, XX, XQ, and HG: conceptualization. XX, XQ, CL, and ZW: methodology. LL and CL: software. XQ, CL, and YW: validation. ZW, TL, and YL: formal analysis. ZW, TL, YW, YS, LL, ZZ, XH, PC, YL, BW, JL, YZ, YH, and HZ: investigation. XX, XQ, YL, XH, and HG: resources. LL, TL, and CL: data curation. ZW, XQ, and HG: manuscript (original draft) preparation. XX and YH: review and editing of manuscripts. XX, XQ, and HG: supervision.

## REFERENCES

1. Rayman MP. Selenium and human health. *Lancet*. (2012) 379:1256–68. doi: 10.1016/S0140-6736(11)61452-9
2. Takata N, Myburgh J, Botha A, Nomngongo PN. The importance and status of the micronutrient selenium in South Africa: a review. *Environ Geochem Health*. (2021). doi: 10.1007/s10653-021-01126-3
3. Shetty S, Marsicano JR, Copeland PR. Uptake and utilization of selenium from selenoprotein P. *Biol Trace Elem Res*. (2018) 181:54–61. doi: 10.1007/s12011-017-1044-9
4. Duntas LH, Benavente S. Selenium: an element for life. *Endocrine*. (2015) 48:756–75. doi: 10.1007/s12020-014-0477-6
5. Yu D, Liang D, Lei L, Zhang R, Sun X, Lin Z. Selenium geochemical distribution in the environment and predicted human daily dietary intake in northeastern Qinghai. *China Environ Sci Pollut Res Int*. (2015) 22:11224–35. doi: 10.1007/s11356-015-4310-4
6. Everson TM, Niedzwiecki MM, Toth D, Tellez-Plaza M, Liu H, Barr DB, et al. Metal biomarker mixtures and blood pressure in the United States:

ZW and XX: project administration. XX: funding acquisition. All authors have read the final version of the manuscript and have given final approval for this version of the manuscript to be published.

## FUNDING

This study was supported by funding from the National Key Research and Development Program [2016YFE0205400, 2018ZX09739010 and 2018ZX09301034003]; the Science and Technology Planning Project of Guangzhou, China [201707020010]; the Science, Technology, and Innovation Committee of Shenzhen [GJHS20170314114526143 and JSGG20180703155802047]; the Economic, Trade, and Information Commission of Shenzhen Municipality [20170505161556110, 20170505160926390, and 201705051617070]; the National Natural Science Foundation of China [81730019 and 81973133]; the President Foundation of Nanfang Hospital, Southern Medical University, [2018Z009]; Outstanding Youth Development Scheme of Nanfang Hospital, Southern Medical University, [2019J004]; the National Natural Science Foundation of China, [82000691]; the Science and Technology Planning Project of Guangzhou, China [202102021155]; the 111 Project from the Education Ministry of China [B18053]; the National Natural Science Foundation of China [81960074]; and the Jiangxi Outstanding Person Foundation [20192BCBL23024].

## ACKNOWLEDGMENTS

We acknowledge the contribution of all staff members who participated in this study as well as the study participants.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fnut.2022.882309/full#supplementary-material>

cross-sectional findings from the 1999-2006 national health and nutrition examination survey (NHANES). *Environ Health*. (2021) 20:15. doi: 10.1186/s12940-021-00695-1

7. Kuruppu D, Hendrie HC, Yang L, Gao S. Selenium levels and hypertension: a systematic review of the literature. *Public Health Nutr*. (2014) 17:1342–52. doi: 10.1017/S1368980013000992
8. Wang Z, Ma H, Song Y, Lin T, Liu L, Zhou Z, et al. Plasma selenium and the risk of first stroke in adults with hypertension: a secondary analysis of the China Stroke Primary Prevention Trial. *Am J Clin Nutr*. (2022) 115:222–31. doi: 10.1093/ajcn/nqab320
9. Sinha N, Dabla PK. Oxidative stress and antioxidants in hypertension—a current review. *Curr Hypertens Rev*. (2015) 11:132–42. doi: 10.2174/1573402111666150529130922
10. Lu J, Lu Y, Wang X, Li X, Linderman GC, Wu C, et al. Prevalence, awareness, treatment, and control of hypertension in China: data from 1.7 million adults in a population-based screening study (China PEACE million persons project). *Lancet*. (2017) 390:2549–58. doi: 10.1016/S0140-6736(17)32478-9

11. Jones GD, Droz B, Greve P, Gottschalk P, Poffet D, McGrath SP, et al. Selenium deficiency risk predicted to increase under future climate change. *Proc Natl Acad Sci USA*. (2017) 114:2848–53. doi: 10.1073/pnas.1611576114
12. Li N, Gao Z, Luo D, Tang X, Chen D, Hu Y. Selenium level in the environment and the population of Zhoukoudian area, Beijing, China. *Sci Total Environ*. (2007) 381:105–11. doi: 10.1016/j.scitotenv.2007.03.027
13. Mihailovič MB, Avramovič DM, Jovanovič IB, Pesut OJ, Matić DP, Stojanov VJ. Blood and plasma selenium levels and GSH-Px activities in patients with arterial hypertension and chronic heart disease. *J Environ Pathol Toxicol Oncol*. (1998) 17:285–9.
14. Laclaustra M, Navas-Acien A, Stranges S, Ordovas JM, Guallar E. Serum selenium concentrations and hypertension in the US population. *Circ Cardiovasc Qual Outcomes*. (2009) 2:369–76. doi: 10.1161/CIRCOUTCOMES.108.831552
15. Suadicanì P, Hein HO, Gyntelberg F. Serum selenium concentration and risk of ischaemic heart disease in a prospective cohort study of 3000 males. *Atherosclerosis*. (1992) 96:33–42. doi: 10.1016/0021-9150(92)90035-f
16. Li M, Yun H, Huang J, Wang J, Wu W, Guo R, et al. Hair selenium content in middle-aged and elderly Chinese population. *Biol Trace Elem Res*. (2020) 199:3571–8. doi: 10.1007/s12011-020-02482-4
17. Wu W, Jiang S, Zhao Q, Zhang K, Wei X, Zhou T, et al. Environmental exposure to metals and the risk of hypertension: a cross-sectional study in China. *Environ Pollut*. (2018) 233:670–8. doi: 10.1016/j.envpol.2017.10.111
18. Gao S, Jin Y, Unverzagt FW, Liang C, Hall KS, Cao J, et al. Selenium level and depressive symptoms in a rural elderly Chinese cohort. *BMC Psychiatry*. (2012) 12:72. doi: 10.1186/1471-244X-12-72
19. Combs GF Jr. Biomarkers of selenium status. *Nutrients*. (2015) 7:2209–36. doi: 10.3390/nu7042209
20. Vinceti M, Filippini T, Jablonska E, Saito Y, Wise LA. Safety of selenium exposure and limitations of selenoprotein maximization: molecular and epidemiologic perspectives. *Environ Res*. (2022) 211:113092. doi: 10.1016/j.envres.2022.113092
21. Rayman MP. Selenium intake, status, and health: a complex relationship. *Hormones (Athens)*. (2020) 19:9–14. doi: 10.1007/s42000-019-00125-5
22. Vinceti M, Filippini T, Wise LA, Rothman KJ. A systematic review and dose-response meta-analysis of exposure to environmental selenium and the risk of type 2 diabetes in nonexperimental studies. *Environ Res*. (2021) 197:111210. doi: 10.1016/j.envres.2021.111210
23. Li Q, Liu M, Hou J, Jiang C, Li S, Wang T. The prevalence of Keshan disease in China. *Int J Cardiol*. (2013) 168:1121–6. doi: 10.1016/j.ijcard.2012.11.046
24. Liu LS. Writing group of 2010 Chinese guidelines for the management of hypertension. [2010 Chinese guidelines for the management of hypertension]. *Zhonghua Xin Xue Guan Bing Za Zhi*. (2011) 39:579–615.
25. Chen H, Cui Z, Lu W, Wang P, Wang J, Zhou Z, et al. Geographical, sex, age, and seasonal differences in serum manganese status among Chinese adults with hypertension. *Biol Trace Elem Res*. (2022). doi: 10.1007/s12011-022-03135-4
26. Dinh QT, Cui Z, Huang J, Tran TAT, Wang D, Yang W, et al. Selenium distribution in the Chinese environment and its relationship with human health: a review. *Environ Int*. (2018) 112:294–309. doi: 10.1016/j.envint.2017.12.035
27. Yu T, Yang ZF, Hou QY, Lv YY, Xi XX, Li M. Topsoil selenium distribution in relation to geochemical factors in main agricultural areas of China. In: Bañuelos GS editor. *Global Advances in Selenium Research From Theory to Application*. London: Taylor & Francis Group (2016). doi: 10.1201/b19240-4
28. Combs GF Jr. Selenium in global food systems. *Br J Nutr*. (2001) 85:517–47. doi: 10.1079/bjn2000280
29. Bastola MM, Locatis C, Maisiak R, Fontelo P. Selenium, copper, zinc and hypertension: an analysis of the national health and nutrition examination survey (2011–2016). *BMC Cardiovasc Disord*. (2020) 20:45. doi: 10.1186/s12872-020-01355-x
30. Ning Y, Wang X, Zhang P, Anatoly SV, Prakash NT, Li C, et al. Imbalance of dietary nutrients and the associated differentially expressed genes and pathways may play important roles in juvenile Kashin-Beck disease. *J Trace Elem Med Biol*. (2018) 50:441–60. doi: 10.1016/j.jtemb.2018.01.012
31. Wang X, Ning Y, Yang L, Yu F, Guo X. Zinc: the other suspected environmental factor in kashin-beck disease in addition to selenium. *Biol Trace Elem Res*. (2017) 179:178–84. doi: 10.1007/s12011-017-0964-8
32. Li S, Bañuelos GS, Wu L, Shi W. The changing selenium nutritional status of Chinese residents. *Nutrients*. (2014) 6:1103–14. doi: 10.3390/nu6031103
33. Huang Q, Wang L, Jiang H, Wang H, Zhang B, Zhang J, et al. Intra-individual double burden of malnutrition among adults in China: evidence from the China health and nutrition survey 2015. *Nutrients*. (2020) 12:2811. doi: 10.3390/nu12092811
34. Vinceti M, Filippini T, Del Giovane C, Dennert G, Zwahlen M, Brinkman M, et al. Selenium for preventing cancer. *Cochrane Database Syst Rev*. (2018) 1:CD005195. doi: 10.1002/14651858.CD005195.pub4
35. Steinbrenner H, Duntas LH, Rayman MP. The role of selenium in type-2 diabetes mellitus and its metabolic comorbidities. *Redox Biol*. (2022) 50:102236. doi: 10.1016/j.redox.2022.102236
36. Jenkins DJA, Kitts D, Giovannucci EL, Sahye-Pudaruth S, Paquette M, Blanco Mejia S, et al. Selenium, antioxidants, cardiovascular disease, and all-cause mortality: a systematic review and meta-analysis of randomized controlled trials. *Am J Clin Nutr*. (2020) 112:1642–52. doi: 10.1093/ajcn/nqaa245
37. Sun JW, Shu XO, Li HL, Zhang W, Gao J, Zhao LG, et al. Dietary selenium intake and mortality in two population-based cohort studies of 133 957 Chinese men and women. *Public Health Nutr*. (2016) 19:2991–8. doi: 10.1017/S1368980016001130
38. Inoue Y, Umezaki M, Jiang H, Li D, Du J, Jin Y, et al. Urinary concentrations of toxic and essential trace elements among rural residents in Hainan Island, China. *Int J Environ Res Public Health*. (2014) 11:13047–64. doi: 10.3390/ijerph111213047
39. Su LQ, Jin YL, Unverzagt FW, Cheng YB, Hake AM, Ran L, et al. Nail selenium level and diabetes in older people in rural China. *Biomed Environ Sci*. (2016) 29:818–24. doi: 10.3967/bes2016.109
40. Ahmad KA, Yuan Yuan D, Nawaz W, Ze H, Zhuo CX, Talal B, et al. Antioxidant therapy for management of oxidative stress induced hypertension. *Free Radic Res*. (2017) 51:428–38. doi: 10.1080/10715762.2017.1322205
41. Bodnar M, Szczygłowska M, Konieczka P, Namiesnik J. Methods of selenium supplementation: bioavailability and determination of selenium compounds. *Crit Rev Food Sci Nutr*. (2016) 56:36–55. doi: 10.1080/10408398.2012.709550
42. Ullah H, Liu G, Yousaf B, Ali MU, Abbas Q, Munir MAM, et al. Developmental selenium exposure and health risk in daily foodstuffs: a systematic review and meta-analysis. *Ecotoxicol Environ Saf*. (2018) 149:291–306. doi: 10.1016/j.ecoenv.2017.11.056
43. Wang J, Yang L, Li H, Li Y, Wei B. Dietary selenium intake based on the Chinese Food Pagoda: the influence of dietary patterns on selenium intake. *Nutr J*. (2018) 17:50. doi: 10.1186/s12937-018-0358-6
44. Filippini T, Michalke B, Wise LA, Malagoli C, Malavolti M, Vescovi L, et al. Diet composition and serum levels of selenium species: a cross-sectional study. *Food Chem Toxicol*. (2018) 115:482–90. doi: 10.1016/j.fct.2018.03.048
45. Chawla R, Filippini T, Loomba R, Cilloni S, Dhillion KS, Vinceti M. Exposure to a high selenium environment in Punjab, India: Biomarkers and health conditions. *Sci Total Environ*. (2020) 719:134541. doi: 10.1016/j.scitotenv.2019.134541
46. Zhong Q, Lin R, Nong Q. Adiposity and serum selenium in U.S. *Adults. Nutrients*. (2018) 10:727. doi: 10.3390/nu10060727
47. Spina A, Guallar E, Rayman MP, Tigbe W, Kandala NB, Stranges S. Anthropometric indices and selenium status in British adults: the U.K. national diet and nutrition survey. *Free Radic Biol Med*. (2013) 65:1315–21. doi: 10.1016/j.freeradbiomed.2013.09.025
48. Hybsier S, Schulz T, Wu Z, Demuth I, Minich WB, Renko K, et al. Sex-specific and inter-individual differences in biomarkers of selenium status identified by a calibrated ELISA for selenoprotein P. *Redox Biol*. (2017) 11:403–14. doi: 10.1016/j.redox.2016.12.025
49. Liu P, Yuan G, Zhou Q, Liu Y, He X, Zhang H, et al. The association between metal exposure and semen quality in Chinese males: the mediating effect of androgens. *Environ Pollut*. (2020) 264:113975. doi: 10.1016/j.envpol.2020.113975
50. Qazi IH, Angel C, Yang H, Zoidis E, Pan B, Wu Z, et al. Role of selenium and seleno-proteins in male reproductive function: a review of past and present evidences. *Antioxidants (Basel)*. (2019) 8:268. doi: 10.3390/antiox8080268
51. Mirnamihi M, Faroughi F, Tahmasbpour E, Ebrahimi P, Beigi Harchegani A. An overview on role of some trace elements in human reproductive

- health, sperm function and fertilization process. *Rev Environ Health*. (2019) 34:339–48. doi: 10.1515/reveh-2019-0008
52. Liu Z, Zhao L, Man Q, Wang J, Zhao W, Zhang J. Dietary micronutrients intake status among chinese elderly people living at home: data from CNNHS 2010-2012. *Nutrients*. (2019) 11:1787. doi: 10.3390/nu11081787
  53. Cai Z, Zhang J, Li H. Selenium, aging and aging-related diseases. *Aging Clin Exp Res*. (2019) 31:1035–47. doi: 10.1007/s40520-018-1086-7
  54. Salonen JT, Salonen R, Ihanainen M, Parviainen M, Seppänen R, Kantola M, et al. Blood pressure, dietary fats, and antioxidants. *Am J Clin Nutr*. (1988) 48:1226–32. doi: 10.1093/ajcn/48.5.1226
  55. Vinceti M, Chawla R, Filippini T, Dutt C, Cilloni S, Loomba R, et al. Blood pressure levels and hypertension prevalence in a high selenium environment: results from a cross-sectional study. *Nutr Metab Cardiovasc Dis*. (2019) 29:398–408. doi: 10.1016/j.numecd.2019.01.004
  56. Swanson CA, Longnecker MP, Veillon C, Howe M, Levander OA, Taylor PR, et al. Selenium intake, age, gender, and smoking in relation to indices of selenium status of adults residing in a seleniferous area. *Am J Clin Nutr*. (1990) 52:858–62. doi: 10.1093/ajcn/52.5.858
  57. Isobe Y, Asakura H, Tsujiguchi H, Kannon T, Takayama H, Takeshita Y, et al. alcohol intake is associated with elevated serum levels of selenium and selenoprotein P in humans. *Front Nutr*. (2021) 8:633703. doi: 10.3389/fnut.2021.633703
  58. Snook JT. Effect of ethanol use and other lifestyle variables on measures of selenium status. *Alcohol*. (1991) 8:13–6. doi: 10.1016/0741-8329(91)91136-p
  59. Breslow RA, Chen CM, Graubard BI, Jacobovits T, Kant AK. Diets of drinkers on drinking and nondrinking days: NHANES 2003-2008. *Am J Clin Nutr*. (2013) 97:1068–75. doi: 10.3945/ajcn.112.050161
  60. Vinceti M, Grill P, Malagoli C, Filippini T, Storani S, Malavolti M, et al. Selenium speciation in human serum and its implications for epidemiologic research: a cross-sectional study. *J Trace Elem Med Biol*. (2015) 31:1–10. doi: 10.1016/j.jtemb.2015.02.001
  61. Maass F, Michalke B, Willkommen D, Schulte C, Tönges L, Boerger M, et al. Selenium speciation analysis in the cerebrospinal fluid of patients with Parkinson's disease. *J Trace Elem Med Biol*. (2020) 57:126412. doi: 10.1016/j.jtemb.2019.126412
  62. Solovyev N, Vanhaecke F, Michalke B. Selenium and iodine in diabetes mellitus with a focus on the interplay and speciation of the elements. *J Trace Elem Med Biol*. (2019) 56:69–80. doi: 10.1016/j.jtemb.2019.07.005

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Wang, Lin, Wei, Song, Liu, Zhou, Huang, Chen, Liu, Li, Wang, Li, Zhang, Huo, Zhang, Xu, Qin and Guo. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.