Check for updates

#### **OPEN ACCESS**

EDITED BY José María Huerta, Carlos III Health Institute (ISCIII), Spain

#### REVIEWED BY

Estêvão Rios Monteiro, University Center Augusto Motta, Brazil Dana Mihaela Ciobanu, University of Medicine and Pharmacy Iuliu Hatieganu, Romania

\*CORRESPONDENCE

Maria Del Carmen Bisi Molina ⊠ mdcarmen2007@gmail.com Taísa Sabrina Silva Pereira ⊠ taisa.silva@udlap.mx

RECEIVED 27 September 2024 ACCEPTED 04 April 2025 PUBLISHED 02 May 2025 CORRECTED 18 August 2025

#### CITATION

Mercadante Urquía YJ, Martins HX, Silva Pereira TS, Batista de Azevedo L, Lopes Júnior LC and Molina MDCB (2025) Physical activity and screen time behavior, and non-alcoholic beverage consumption during the COVID-19 pandemic in the longitudinal study of adult health (ELSA-Brasil). *Front. Nutr.* 12:1503010. doi: 10.3389/fnut.2025.1503010

#### COPYRIGHT

© 2025 Mercadante Urquía, Martins, Silva Pereira, Batista de Azevedo, Lopes Júnior and Molina. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

# Physical activity and screen time behavior, and non-alcoholic beverage consumption during the COVID-19 pandemic in the longitudinal study of adult health (ELSA-Brasil)

Yazareni José Mercadante Urquía<sup>1</sup>, Haysla Xavier Martins<sup>1</sup>, Taísa Sabrina Silva Pereira<sup>2</sup>\*, Letícia Batista de Azevedo<sup>3</sup>, Luís Carlos Lopes Júnior<sup>1</sup> and Maria Del Carmen Bisi Molina<sup>1,4</sup>\*

<sup>1</sup>Post-graduate Program in Collective Health, Universidade Federal do Espírito Santo, Vitória, Brazil, <sup>2</sup>Department of Health Sciences, Universidad de las Américas Puebla, Puebla, Mexico, <sup>3</sup>Post-graduate Program in Nutrition and Health, Universidade Federal do Espírito Santo, Vitória, Brazil, <sup>4</sup>Postgraduate Program in Nutrition and Longevity, Universidade Federal de Alfenas, Alfenas, Brazil

**Background:** The COVID-19 pandemic affected various dietary and lifestyle behaviors. Understanding the relationship between physical activity, screen time, and beverage consumption is critical to identify at-risk populations, particularly individuals with chronic non-communicable diseases (NCDs), for targeted intervention strategies.

**Objective:** This study aimed to evaluate the relationship between physical activity, screen time behavior patterns, and non-alcoholic beverage consumption among participants in ELSA-Brasil during the COVID supplementary study.

**Methods:** The study was conducted in five of the six ELSA-Brasil research centers, from July 2020 to March 2021. Sociodemographic variables, physical activity, screen time, non-alcoholic beverage consumption, and lifestyle factors were analyzed using bivariate analyses and binary logistic regression models.

**Results:** The total sample consisted of 4,442 participants with a mean age of  $62.0 \pm 8.6$  years. Significant gender differences were found in sociodemographic and lifestyle variables, influencing beverage consumption patterns. Participants with excessive screen time and physical inactivity were more likely to consume sugar-sweetened beverages. Specifically, men exhibited a significant association with an odds ratio of 2.44 (95%CI: 1.54-3.84), while women had an odds ratio of 1.72 (95% CI: 1.02-2.91). Men with appropriate screen time but physically inactive behavior had nearly double the likelihood of consuming sugar-sweetened artificial beverages. Additionally, men with excessive screen time and physical inactivity had higher consumption of industrialized juices (OR: 1.93; 95% CI: 1.22-3.06), while women were more likely to consume soft drinks (OR: 2.28; 95% CI : 1.10-4.72).

**Conclusion:** These findings underscore the importance of addressing both screen time and physical activity in public health initiatives, through gender-specific approaches that account for socioeconomic disparities when prioritizing interventions.

#### KEYWORDS

screen time (ST), physical activity, carbonated beverages, sugar-sweetened beverages (SSB), COVID-19, health behavior, lifestyle

### **1** Introduction

The social restriction measures resulting from the COVID-19 pandemic have led to modifications in various aspects of life across different populations (1), with significant distinctions based on pre-existing health and socioeconomic conditions (2). Among the changes observed, there was a substantial increase in the time spent on screen devices, such as smartphones, video games, computers, and televisions (3), and, conversely, a reduction in physical activity (4). In Brazil, a cross-sectional study conducted by Malta et al. (5) evidenced an increase in screen time during the COVID-19 pandemic compared to the pre-pandemic period, with an average increase of 1 h and 45 min for TV and 1 h and 30 min for computer or tablet use. Regarding physical inactivity, a 26% increase was observed in Brazilian adults (6). Moreover, in a sex-stratified analysis using data from the Brazilian longitudinal study of adult health (ELSA-Brasil), Molina et al. (7) observed a reduction in physical activity levels during the COVID-19 pandemic, with declines of 195.5 (standard deviation [SD]  $\pm$  1,146.4) and 240.5 (SD  $\pm$  1,474.2) metabolic equivalents/week among women and men, respectively.

Social distancing measures also significantly impacted dietary consumption (8, 9). A study conducted with adults from southeastern Brazil evidenced an increase in snacking between meals, higher consumption of morning snacks, and a decrease in breakfast consumption (10). These changes are partly attributed to the repercussions of the pandemic on individuals' mental health (9, 11), where coping with feelings such as anxiety or even boredom often involves eating (12, 13).

Regarding the consumption of non-alcoholic beverages during the pandemic, a cross-sectional study with 4,022 Australian adults identified that most did not change their consumption of sugary drinks. Among those who increased their frequency of intake, the most likely were men and those with children at home, while those with higher education levels showed a reduction in frequency (14). These results align with those of Park and collaborators (2023) in a sample of 4,034 adults from the United States, where being male, having lower education, and having children were associated with a higher likelihood of being a high consumer ( $\geq 2$  times/day) of sugary drinks (15). A general increase in coffee consumption was observed during the pandemic (16, 17), with remote work being identified as a driver of this global demand increase (18).

A comprehensive understanding of the relationship between physical activity, screen time, and non-alcoholic beverage consumption is critical, as these represent modifiable risk factors for non-communicable diseases (NCDs). Their interplay during the pandemic may have exacerbated health vulnerabilities and disparities. The COVID-19 pandemic created a unique opportunity to examine how abrupt lifestyle changes influenced health behaviors.

By examining these relationships in ELSA-Brasil participants – a cohort with detailed pre-pandemic data – we can isolate pandemic-specific effects and identify high-risk groups requiring targeted interventions. Although the COVID supplementary study is

cross-sectional, the availability of baseline data from ELSA-Brasil enables a more nuanced interpretation of pandemic-related behavioral changes. This investigation underscores the urgency of examining these combined behaviors to mitigate long-term health consequences. The ELSA-Brasil cohort, with its robust pre-pandemic data and diverse adult population, provides an exceptional platform to assess these relationships with greater validity than conventional cross-sectional studies, offering insights particularly valuable for public health strategies targeting similar populations during future health crises.

Engaging in unhealthy behavior makes individuals susceptible to adopting other deleterious habits (16). Furthermore, the negative modifications resulting from COVID-19, such as physical inactivity (17), sedentary behavior – represented by increased screen time – (18), and excessive consumption of certain non-alcoholic beverages (19, 20), amplify the burden of NCDs. These changes contribute as risk factors for the occurrence of obesity and other NCDs. Therefore, it is essential to evaluate these lifestyle habits in a combined manner.

We hypothesized that lower physical activity levels and higher screen time would be associated with greater consumption of sugarsweetened beverages during the pandemic period. This hypothesis was grounded in previous evidence showing that sedentary behaviors and unhealthy dietary patterns frequently co-occurred during the pandemic, as observed in Latin American populations (21) and Brazilian adolescents (22), while stress-related coping mechanisms during lockdowns drove parallel increases in both screen time (23, 24) and consumption of energy-dense foods like sugary drinks through emotional distress pathways (12, 13). The ELSA-Brasil design permitted examination of these associations while controlling for relevant confounders.

This study addresses critical gaps in understanding the interplay between physical activity, screen time, and non-alcoholic beverage consumption during the COVID-19 pandemic. By focusing on Brazilian public servants, both active and retired, it provides unique insights into pandemic-induced behavioral changes. The originality of this research lies in its comprehensive evaluation of these combined lifestyle factors and its detailed analysis by type of beverage, a novel approach not extensively explored in previous studies among adults. In the post-pandemic context, our study offers crucial empirical evidence on the relationships between these factors. The findings contribute to a broader understanding of the health risks associated with pandemic-induced lifestyle changes, with the potential to inform the development of targeted interventions for at-risk groups. These insights strengthen the theoretical foundation on the adverse health effects of such lifestyle changes in future similar scenarios.

Given the above, this study aims to evaluate the relationship between physical activity and screen time behavior patterns and the consumption of non-alcoholic beverages among public servants, both active and retired, participating in ELSA-Brasil during the COVID-19 pandemic.

# 2 Materials and methods

### 2.1 Study design and population

The present study is a cross-sectional quantitative analysis that utilizes data from a supplementary study conducted during the COVID-19 pandemic (2020-2021). This supplementary study is part of a larger multicenter cohort, ELSA-Brasil, which began its baseline period in 2008-2010 with 15,105 participants, both active and retired public servants, aged 35 to 74 at the time of recruitment. ELSA-Brasil aims to investigate associations between chronic diseases, primarily cardiovascular diseases and diabetes, and biological, behavioral, environmental, occupational, and social factors (25). The study is conducted at six research centers located in five public universities and one research institute of the Brazilian Ministry of Health (26). The supplementary study, named the COVID wave, was implemented to assess the pandemic's impact on the health of the cohort participants. Both ELSA-Brasil and the supplementary study received approval from the Ethics Committees of the participating research centers, and all participants provided informed consent, ensuring data confidentiality and ethical treatment (27).

In the supplementary COVID study (2020–2021), participants from five of the six research centers were invited from three regions of the country, including the Northeast, South, and Southeast (Federal Universities of Bahia, Rio Grande do Sul, Espírito Santo and Minas Gerais; and Oswaldo Cruz Foundation [FIOCRUZ], located in Rio de Janeiro). A total of 5,544 individuals agreed to respond to the questionnaires administered during this phase of the study. Participants without data on screen time (n = 573), beverage consumption (n = 18), and physical activity (n = 511) were excluded from the analysis due to non-response. The total sample size included in the analysis was 4,442 participants. Supplementary Figure 1 illustrates the flowchart of cohort participants and their follow-up over the study periods.

Outliers were adjusted to the 99th percentile value of the respective variable (screen time > 16 h/day and physical activity > 645 min/week) to minimize the impact of extreme values on the analysis.

This study adhered to the Declaration of Helsinki guidelines and received approval from ELSA-Brasil and the Research Ethics Committees of all participating institutions, with the following approval numbers: 343/06 at the FIOCRUZ-RJ Institute (COVID Study – N° 4.063.982), 041/06 at UFES (COVID Study - N° 4.090.940), 186/06 at UFMG (COVID Study - N° 4.082.055), 194/06 at UFRGS (COVID Study - N° 4.023.601), and 027/06 at UFBA (COVID Study - N° 4.067.184).

### 2.2 Data collection and measures

Data were collected during the COVID wave, from July 2020 to March 2021, conducted online using a specific platform and through telephone contact due to social distancing measures implemented in various Brazilian states during this period. Sociodemographic, lifestyle, and food and beverage consumption data were collected.

### 2.2.1 Sociodemographic data

From the data of Wave 3 (2017–2019) of ELSA-Brasil, the participants' age was calculated by adding an average of 3 years, and

the *per capita* income was treated as a continuous variable. Additionally, data collected during the COVID wave included sex (male or female), self-reported race/ethnicity (categorized into two categories: White, and Black, Brown, Yellow, and Indigenous), according to the classification of the *Instituto Brasileiro de Geografia e Estatística* (28); current employment status (active or retired), and whether participants worked remotely during the pandemic (yes or no).

# 2.2.2 Exposure: physical activity and screen time behavior

To characterize physical activity and screen time behavior, a composite variable was created combining both physical activity behavior and screen time. In the COVID wave of the ELSA-Brasil study, physical activity (PA) was assessed using the short version of the International Physical Activity Questionnaire (IPAQ), adapted and validated for the Brazilian population (29). The questionnaire included the following questions: "Since the beginning of social distancing, on average how many days a week do you take walks in your free time?"; "Since the beginning of social distancing, how many days a week do you do vigorous physical activities in your free time?"; and "Since the beginning of social distancing, how many days a week do you do moderate physical activities?." PA was reported in minutes per week by multiplying the weekly frequency by the duration of each activity. Only PA sessions lasting at least 10 consecutive minutes were considered. Vigorous PA was defined as activities requiring significant physical effort and causing much faster breathing than usual, while moderate PA required moderate effort and caused slightly faster breathing than usual. Participants were considered physically active if they engaged in ≥150 min/week of moderate-to-vigorous PA and physically inactive if they engaged in <150 min/week of moderate-tovigorous PA (30).

Screen time was estimated from participants' self-reported time spent in front of screen devices during social distancing, using the question "How much time are you now spending in front of any screen (including cell phone, computer, TV, laptops, or others)?." The data were obtained in hours per day and categorized using the sample's median screen time hours as the cutoff point, adjusted for outliers based on the 99th percentile (16 h/day), into two categories:  $\leq 6$  h/day (Appropriate Screen Time) and >6 h/day (Excessive Screen Time). The cutoff point was based on the median screen time of the study sample, due to the lack of consensus in the current literature regarding recommended screen time for adults. Additionally, this cutoff coincides with that used in another significant study on the Brazilian population, which considered screen time  $\geq 6$  h/day as high (31).

Based on these variables, a recategorization was performed, creating four categories of the combined Physical Activity and Screen Time Behavior variable: "Appropriate Screen Time/Physically Active"; "Appropriate Screen Time/Physically Inactive"; "Excessive Screen Time/Physically Active"; and "Excessive Screen Time/ Physically Inactive."

# 2.2.3 Outcome: non-alcoholic beverage consumption on the previous day

Non-alcoholic beverage consumption was assessed through a 24-h dietary recall with the question "Look at this list of drinks and check all that you consumed yesterday (from when you woke up until you went to sleep)," and data were obtained on the consumption of

soft drink or soda, industrialized juices (boxed, bottled, concentrated, or powdered), sugar-sweetened artificial beverages (soda + industrialized juices), and coffee with and without sugar. Participants indicated whether they consumed each of these beverages on the previous day. The 24-h dietary recall was applied twice during the COVID wave, with a minimum interval of 1 month between the first and second application.

### 2.2.4 Lifestyle covariates

Data on smoking were collected with the question "Do you currently smoke cigarettes?" with three response options (Never smoked, Former smoker, and Current smoker). Alcohol consumption data were collected with the question "Since the beginning of social distancing, have you consumed any type of alcoholic beverage?" with a dichotomous response (Yes or No).

### 2.3 Statistical analysis

Analyses were stratified by sex due to the interaction in the association between exposure and outcome, as well as the different patterns of PA/screen time behavior and beverage consumption between men and women. For evaluating differences between groups, Student's T-test and ANOVA (Supplementary Table 1) were applied for continuous variables, and the Chi-Square test was used for categorical variables. Bivariate analyses were performed to assess the association between beverage consumption and sociodemographic, lifestyle, and PA/screen time behavior variables. Binary logistic regression analysis was performed using an initial unadjusted model, which was subsequently adjusted based on a theoretical model with two hierarchical levels (32): adjusted model 1 included sociodemographic variables (age, per capita income, remote work), and adjusted model 2 included sociodemographic and lifestyle variables (smoking and alcohol consumption). Additionally, a test was conducted to rule out multicollinearity. Variables remained in the model if they reached a significance level of 20% in the bivariate analyses. Descriptive and exploratory analyses were performed using SPSS version 23.0 with a significance level of 5%.

### **3 Results**

The total study sample consisted of 4,442 participants with a mean age of  $62.0 \pm 8.6$  years. Significant differences were found between genders in relation to sociodemographic and lifestyle variables (Table 1). Most of the sample consisted of women (58.2%) and individuals who were working (59.1%), with appropriate screen time and physical inactivity (45.4%), non-smokers (65.5%), and alcohol consumers during the pandemic (62.0%). Additionally, men exhibited a higher average screen time (p = 0.026).

Beverage consumption varied according to sociodemographic and lifestyle variables (Table 2). The majority of soft drink or soda consumers were women (p < 0.001), working (p = 0.003), non-smokers (p < 0.001), and working remotely during the pandemic (p = 0.012). They were also physically inactive (p < 0.001) and predominantly exhibited combinations of appropriate or inappropriate screen time with physical inactivity (p < 0.001). A similar pattern was

observed among consumers of other types of sugar-sweetened artificial beverages.

Individuals who consumed coffee, both with and without sugar or sweeteners, were predominantly white (p < 0.001), employed, working remotely during the pandemic (p < 0.001), non-smokers, and alcohol consumers during the pandemic. Consumers of both types of beverages were mostly physically inactive (p < 0.001), and the combined behaviors of physical activity and screen time showed predominant patterns of "Appropriate screen time/Physically inactive" (p < 0.001).

The mean per capita income was significantly lower among those consuming soft drinks or soda, sugar-sweetened artificial beverages, and coffee with sugar, and significantly higher among those consuming coffee without sugar (data not shown in the table).

The results of the binary logistic regression assessing the association between physical activity/screen time behavior and the consumption of non-alcoholic beverages are presented in Figure 1 and Supplementary Table 2. After adjusting for potentially confounding sociodemographic and lifestyle variables (Adjusted Model 2), it was identified that the behavior category of Excessive screen time/Physically inactive was associated with higher odds of consuming sugar-sweetened artificial beverages among both men and women.

Among men, the behavior of "Appropriate screen time/Physically inactive" was significantly associated with a 1.93-fold increase in the likelihood of consuming sugar-sweetened artificial beverages, whereas presenting a behavior of "Excessive screen time/Physically inactive" increased the likelihood by 2.44 times among men (95% CI 1.54–3.84) and 1.72 times among women (95% CI 1.02–2.91). Additionally, among men, the behavior of "Excessive screen time/Physically inactive" also increased the likelihood of consuming industrialized juices by 2.02 times (95% CI 1.15–3.53), and among women, it increased the likelihood of consuming soft drinks or soda by 2.28 times (95% CI 1.10–4.72). No associations were found between physical activity and screen time behavior and the consumption of coffee with or without sugar.

### 4 Discussion

In this study, the "Excessive Screen Time/Physically Inactive" behavior pattern during the COVID-19 pandemic among participants of the ELSA-Brasil cohort was associated with a 144% increase in the odds of consuming sugar-sweetened artificial beverages among men and a 72% increase among women. This pattern also correlated with a 128% higher probability of soda consumption in women and a 102% higher likelihood of consuming industrialized juices in men. Additionally, the "Adequate Screen Time/Physically Inactive" pattern increased the likelihood of consuming sugar-sweetened artificial beverages by 93% among men.

Despite the fact that no associations were found in the present study between physical activity/screen time behavior and the consumption of coffee with or without sugar, similar associations between *per capita* income and coffee consumption patterns identified in the present study have been found in previous studies conducted in the USA (33) and in Brazilian studies (34, 35). These studies found that individuals with higher income levels tended to consume less sweetened coffee (35). Additionally, as in the present study, the TABLE 1 Sociodemographic characteristics and lifestyle among participants of the COVID supplementary study in ELSA-Brasil by gender (2020–2021).

Variables	Total (n = 4 442)	Sé	n valueª						
	10(0(()) 1)(12)	Males (n = 1857)	Females $(n = 2.585)$						
	n (%)	nuces (17 – 1007)	n (%)						
A = (1 + 346)	(20 + 8 (	(21) 80	(10+84	0.504					
Age (years) $(n - 4.318)$	62.0 ± 8.6	62.1 ± 8.9	61.9 ± 8.4	0.504					
Race (II - 4,010)									
White	2,477 (57.4)	1,070 (59.1)	1,407 (56.1)	0.057					
Black, Brown, Yellow, and Indigenous	1841 (42.6)	742 (40.9)	1,099 (43.9)						
Per capita income (US\$ - monthly)	811,6 ± 587,4	786,4 ± 551,7	829,9 ± 611,3	0.016*					
Employment status <sup>(n = 4,416)</sup>									
Employed/working	2,612 (59.1)	1,269 (69.0)	1,343 (52.1)	<0.001*					
Retired	1804 (40.9)	571 (31.0)	1,233 (47.9)						
Remote work <sup>(n = 2,608)</sup>									
Yes	2022 (77.5)	997 (78.6)	1,025 (76.5)	0.205					
No	586 (22.5)	271 (21.4)	315 (23.5)						
Screen time behavior									
Below the median ( $\leq 6$ h/d)	2,494 (56.1)	1,024 (55.1)	1,470 (56.9)	0.257					
Above the median (> 6 h/d)	1948 (43.9)	833 (44.9)	1,115 (43.1)						
Screen Time (hours)	6.6 ± 3.6	6.7 ± 3.6	6.6 ± 3.5	0.026*					
Physical activity behavior									
Physically inactive (<150 min/week)	3,569 (80.3)	1,433 (77.2)	2,136 (82.6)	<0.001*					
Physically active (≥150 min/week)	873 (19.7)	424 (22.8)	449 (17.4)						
Physical activity and screen time (ST) behavior	r								
Appropriate ST/Physically Active	478 (10.8)	233 (12.5)	245 (9.5)	<0.001*					
Appropriate ST/Physically Inactive	2016 (45.4)	791 (42.6)	1,225 (47.4)						
Excessive ST/Physically Active	395 (8.9)	191 (10.3)	204 (7.9)						
Excessive ST/Physically Inactive	1,553 (35.0)	642 (34.6)	911 (35.2)						
Smoking									
Non-smoker	2,909 (65.5)	1,116 (60.1)	1793 (69.4)	<0.001*					
Former smoker	1,251 (28.2)	617 (33.2)	634 (24.5)						
Current smoker	282 (6.3)	124 (6.7)	158 (6.1)						
Alcoholic beverage consumption									
Yes	2,756 (62.0)	1,324 (71.3)	1,432 (55.4)	<0.001*					
No	1,686 (38.0)	533 (28.7)	1,153 (44.6)						

<sup>a</sup>Chi-square test for categorical variables and Student's *t*-test for continuous variables.

\*p values <0.05, highlighted in bold, were considered statistically significant.

consumption of sugary beverages was also associated with lower *per capita* income among Brazilian adults (34). This can be explained by the lower cost per calorie of low-nutrient foods often chosen by individuals with lower *per capita* income, while those from higher socioeconomic backgrounds have greater nutritional knowledge and prefer healthier food choices (36).

These findings align with the existing literature highlighting the negative impacts of excessive screen time and physical inactivity, especially during periods of social restriction such as the COVID-19 pandemic (37). Excessive screen time has been correlated with higher consumption of unhealthy foods and beverages among adults, children, and adolescents in Brazil and other countries worldwide (21, 22, 38, 39).

Several physiological and behavioral pathways explain this association. Prolonged screen time promotes sedentary behavior, which disrupts glucose metabolism and increases cravings for highcalorie foods, including sugar-sweetened beverages (40). Concurrently, excessive screen exposure elevates stress and anxiety, triggering emotional eating as a coping mechanism, often manifesting as a preference for sugary beverages (41). These mechanisms could synergistically exacerbate sugar-sweetened beverages consumption during pandemic-related restrictions.

The patterns observed in the present study reflect the impact of screen time and physical inactivity on beverage consumption behaviors. During the COVID-19 pandemic, imposed restrictions led to increased time spent at home and remote work, contributing

(Continued)

10.3389/fnut.2025.1503010

Variables	Beverage consumption on the previous day										
	Soft drink or soda		Sugar-sweetened artificial beverages		Industrialized juices		Coffee with sugar/ sweetener		Coffee without sugar		
	Yes n (%)	No <i>n</i> (%)	Yes n (%)	No <i>n</i> (%)	Yes n (%)	No n (%)	Yes n (%)	No n (%)	Yes n (%)	No n (%)	
Gender <sup>(n = 4,375)</sup>											
Male	371 (47.1)	1,486 (40.7)	660 (47.1)	1,197 (39.4)	360 (48.2)	1,497 (40.5)	1,092 (42.3)	765 (41.1)	509 (39.5)	1,348 (42.7)	
Female	417 (52.9)	2,168 (59.3)	741 (52.9)	1844 (60.6)	387 (51.8)	2,198 (59.5)	1,487 (57.7)	1,098 (58.9)	778 (60.5)	1807 (57.3)	
<i>p</i> value <sup>ª</sup>	p = 0	0.001*	p = <	0.001*	p = <0	0.001*	<i>p</i> = 0.405		<i>p</i> = 0.052		
Age (years) (n = 4,357)							,				
	$60.9 \pm 8.7$	$62.3\pm8.6$	$61.2 \pm 8.7$	$62.4 \pm 8.5$	$61.5 \pm 8.8$	62.1 ± 8.6	$62.5\pm8.7$	$61.4 \pm 8.4$	$62.0 \pm 8.4$	$62.0 \pm 8.7$	
<i>p</i> value <sup>a</sup>	p = <0.001*		p = <0.001*		<i>p</i> = 0.059		p = <0.001*		<i>p</i> = 0.770		
Race (n = 4,318)											
White	444 (58.1)	2033 (57.2)	782 (57.5)	1,695 (57.3)	414 (56.9)	2063 (57.4)	1,308 (52.2)	1,169 (64.6)	856 (68.4)	1,621 (52.9)	
Black, Brown, Yellow, and Indigenous	320 (41.9)	1,521 (42.8)	577 (42.5)	1,264 (42.7)	313 (43.1)	1,528 (42.6)	1,200 (47.8)	641 (35.4)	395 (31.6)	1,446 (47.1)	
<i>p</i> value <sup>a</sup>	<i>p</i> =	0.658	<i>p</i> =	0.895	<i>p</i> = 0.805		p < 0.001*		p < 0.001*		
Per capita income (US\$) <sup>(n = 4,346)</sup>											
	739,6 ± 532,8	827,1 ± 597,4	766,5 ± 565,6	832,3 ± 596,0	791,0 ± 595,2	815,8 ± 585,8	715,7 ± 530,7	943,4 ± 634,2	1,003,2 ± 660,3	733,0 ± 535,3	
<i>p</i> value <sup>a</sup>	p = <	0.001*	p = 0	0.001*	<i>p</i> = 0	).299	p < 0	.001*	p < 0.001*		
Employment status (n = 4,416)											
Employed/ Working	501 (63.8)	2,111 (58.1)	877 (62.8)	1735 (57.5)	462 (62.0)	2,150 (58.6)	1,430 (55.8)	1,182 (63.8)	802 (62.8)	1810 (57.7)	
Retired	284 (36.2)	1,520 (41.9)	519 (28.8)	1,285 (42.5)	283 (38.0)	1,521 (41.4)	1,134 (44.2)	670 (36.2)	476 (37.2)	1,328 (42.3)	
<i>p</i> value <sup>a</sup>	p = 0	0.003*	p = 0	0.001*	<i>p</i> = 0	0.086	p < 0	p < 0.001*		p = 0.002*	
Remote work <sup>(n = 2,608)</sup>											
Yes	367 (73.3)	1,655 (78.5)	643 (73.5)	1,379 (79.6)	339 (73.7)	1,683 (78.4)	1,022 (71.7)	1,000 (84.6)	705 (87.9)	1,317 (72.9)	
No	134 (26.7)	452 (21.5)	232 (26.5)	354 (20.4)	121 (26.3)	465 (21.6)	404 (28.3)	182 (15.4)	97 (12.1)	489 (27.1)	
<i>p</i> value <sup>a</sup>	p = 0	0.012*	p = 0	0.001*	<b>p</b> = 0	.031*	p < 0.001*		p < 0.001*		
Screen time behavior											
Below the median ( $\leq 6 \text{ h/d}$ )	402 (51.0)	2092 (57.3)	733 (52.3)	1761 (57.9)	394 (52.7)	2,100 (56.8)	1,559 (60.4)	935 (50.2)	645 (50.1)	1849 (58.6)	
Above the median (> 6 h/d)	386 (49.0)	1,562 (42.7)	668 (47.7)	1,280 (42.1)	353 (47.3)	1,595 (43.2)	1,020 (39.6)	928 (49.8)	642 (49.9)	1,306 (41.4)	
<i>p</i> value <sup>a</sup>	p = 0.002*		p < 0.001*		p = 0.043*		p < 0.001*		p < 0.001*		

TABLE 2 Sociodemographic characteristics and lifestyle among participants of the COVID supplementary study in ELSA-Brasil by beverage consumption (2020–2021).

TABLE 2 (Continued
--------------------

Variables	Beverage consumption on the previous day										
	Soft drink or soda		Sugar-sweetened artificial beverages		Industrialized juices		Coffee with sugar/ sweetener		Coffee without sugar		
	Yes n (%)	No n (%)	Yes n (%)	No n (%)	Yes <i>n</i> (%)	No n (%)	Yes <i>n</i> (%)	No n (%)	Yes <i>n</i> (%)	No <i>n</i> (%)	
Physical activity behavior											
Physically inactive (<150 min/week)	686 (87.1)	2,883 (78.9)	1,190 (84.9)	2,379 (78.2)	621 (83.1)	2,948 (79.8)	2,147 (83.2)	1,422 (76.3)	971 (75.4)	2,598 (82.3)	
Physically active (≥150 min/week)	102 (12.9)	771 (21.1)	211 (15.1)	662 (21.8)	126 (16.9)	747 (20.2)	432 (16.8)	441 (23.7)	316 (24.6)	557 (17.7)	
<i>p</i> value <sup>a</sup>	p < 0.	.001*	p < 0	.001*	<b>p</b> = 0	.038*	p < 0	p < 0.001* p ·		< 0.001*	
Physical activity and screen time (ST) behavior											
Appropriate ST/Physically Active	50 (6.3)	428 (11.7)	105 (7.5)	373 (12.3)	64 (8.6)	414 (11.2)	252 (9.8)	226 (12.1)	167 (13.0)	311 (9.9)	
Appropriate ST/Physically Inactive	352 (44.7)	1,664 (45.5)	628 (44.8)	1,388 (45.6)	330 (44.2)	1,686 (45.6)	1,307 (50.7)	709 (38.1)	478 (37.1)	1,538 (48.7)	
Excessive ST/Physically Active	52 (6.6)	343 (9.4)	106 (7.6)	289 (9.5)	62 (8.3)	333 (9.0)	180 (7.0)	215 (11.5)	149 (11.6)	246 (7.8)	
Excessive ST/Physically Inactive	334 (42.4)	1,219 (33.4)	562 (40.1)	991 (32.6)	291 (39.0)	1,262 (34.2)	840 (32.6)	713 (38.3)	493 (38.3)	1,060 (33.6)	
<i>p</i> value <sup>a</sup>	p < 0.001*		p < 0.001*		p = 0.033*		p < 0.001*		p < 0.001*		
Smoking											
Non-smoker	460 (58.4)	2,449 (67.0)	850 (60.7)	2059 (67.7)	467 (62.5)	2,442 (66.1)	1,578 (61.2)	1,331 (71.4)	869 (67.5)	2040 (64.7)	
Former smoker	241 (30.6)	1,010 (27.6)	419 (29.9)	832 (27.4)	221 (29.6)	1,030 (27.9)	798 (30.9)	453 (24.3)	359 (27.9)	892 (28.3)	
Current	87 (11.0)	195 (5.3)	132 (9.4)	150 (4.9)	59 (7.9)	223 (6.0)	203 (7.9)	79 (4.2)	59 (4.6)	223 (7.1)	
p value <sup>a</sup>	p < 0.	.001*	p < 0.001*		p = 0.072		p < 0.001*		p = 0.006*		
Alcoholic beverage consumption											
Yes	493 (62.9)	2,263 (61.9)	876 (62.5)	1880 (61.8)	456 (61.0)	2,300 (62.2)	1,511 (58.6)	1,245 (66.8)	935 (72.6)	1821 (57.7)	
No	295 (37.4)	1,391 (38.1)	525 (37.5)	1,161 (38.2)	291 (39.0)	1,395 (37.8)	1,068 (41.4)	618 (33.2)	352 (27.4)	1,334 (42.3)	
<i>p</i> value <sup>a</sup>	<i>p</i> = 0.746		<i>p</i> = 0.665		<i>p</i> = 0.563		p < 0.001*		p < 0.001*		

07

Frontiers in Nutrition

<sup>a</sup>Chi-square test for categorical variables and Student's *t*-test for continuous variables.

\**p* values <0.05, highlighted in bold, were considered statistically significant.



#### FIGURE 1

Adjusted logistic regression model: odds ratios (95% CI) for soft drinks, sugar-sweetened artificial beverages, and industrialized juices consumption by physical activity and screen time behavior in the ELSA-Brasil COVID-19 supplementary study (2020–2021).

to increased screen time and sedentary behaviors (23). Even before the pandemic, a study analyzing screen time trends during leisure time among Brazilian adults from 2016 to 2021 found that the average time spent on devices such as mobile phones, computers, and tablets increased significantly, while TV viewing time remained stable (42).

Furthermore, during the pandemic, a cross-sectional study with ELSA-Brasil participants on remote work showed that working from home was associated with a significant increase in screen time and cumulative sitting time of more than 8 h per day, as well as higher domestic physical activity but not leisure-time physical activity (43). These results reflect the impact of lockdown measures on reducing physical activity and increasing sedentary behavior. Similarly, another study with ELSA-Brasil participants indicated that individuals with diabetes experienced a significant increase in screen time (1.3 h/day)

and sitting or reclining time (1.4 h/day) during the pandemic, while physical activity decreased notably (44).

Several mechanisms may explain the observed associations. Screen time often displaces physical activity, leading to prolonged periods of sedentary behavior, which is independently associated with poor dietary habits, including increased consumption of sugary drinks (45). Additionally, a study with young Colombian adults observed a mild negative correlation between daily physical activity and the frequency of sugary drink consumption, particularly in men, with higher weekly vigorous physical activity being associated with lower intake of these drinks (46). Multiple mechanisms have been suggested to explain the relationship between screen time and unhealthy food consumption, including exposure to advertising and "mindless eating" (38, 47). A sedentary lifestyle, combined with frequent consumption of sugary drinks, creates a synergistic effect, contributing to adverse health outcomes (48–51). Studies have shown that physical inactivity can influence the consumption of sugary drinks (46, 52).

Moreover, stress and the disruption of normal routines during the pandemic may have intensified these behaviors: increased consumption of unhealthy foods, physical inactivity, and sedentary behavior (23). A systematic review identified physical inactivity and increased screen time as significant factors associated with weight gain during the COVID-19 pandemic (23).

In this context, the pandemic has been associated with increased screen time as a coping mechanism, which in turn can lead to higher consumption of unhealthy foods and sugary drinks (23, 24). The relationship between physical inactivity and sugary drink consumption may be mediated by brain reward responses to food cues, which are inversely associated with physical activity (53, 54), as well as the availability of time and ease of access to these drinks during sedentary activities that increased during the pandemic.

In pre-pandemic studies with participants from the ELSA-Brasil Wave 1, it was identified that most individuals engaged in low physical activity (76.9%), similar to the current study where most were physically inactive (80.3%) (55). Additionally, in the baseline (Wave 1) of the study, those who consumed sweetened beverages predominantly had a low level of physical activity, while those who consumed unsweetened and artificially sweetened beverages generally participated in moderate to intense physical activity. Furthermore, in that wave, individuals with a normal nutritional status reported higher consumption of sugary drinks, while those with obesity consumed more sugar-sweetened artificial beverages (55).

A study conducted with adults and adolescents from eight Latin American countries, including Brazil, found that 30.49% of the sample exhibited a combination of unhealthy behaviors: spending a lot of time in front of a screen and consuming sugary drinks (21). In Brazil, Chile, Peru, and Ecuador, the combination of an unhealthy diet or sugary drink consumption along with excessive screen use predominated (21). The authors concluded that these unhealthy behaviors were highly prevalent, highlighting the association between sedentary behavior and sugary drink consumption (21).

The WHO defines physical activity as any bodily movement produced by skeletal muscles which requires energy expenditure. Sedentary screen time refers to the time spent viewing screen-based entertainment (TV, computer, mobile devices), excluding screenbased games that require physical activity or movement (30). Screen time is a proxy variable for sedentary behavior. There is robust evidence establishing a relationship between sedentary behavior and all-cause mortality, as well as cardiovascular diseases, type 2 diabetes, and metabolic syndrome. Additionally, there is moderate evidence suggesting an association with the incidence rates of ovarian, colon, and endometrial cancers (48). A Brazilian cohort study identified a higher risk of mortality among adults who spend more time watching television, regardless of physical activity and other variables (49).

On the other hand, research indicates that high consumption of sugary drinks is linked to adverse metabolic outcomes, including an increased risk of obesity, type 2 diabetes, and metabolic syndrome (50, 51, 56). Due to their ability to rapidly and significantly raise blood glucose and insulin levels, sugary drinks contribute to a high dietary glycemic load. Consuming a diet with a high glycemic load is known to promote glucose intolerance and insulin resistance, as well as increased fat accumulation, particularly in the visceral regions, exacerbating the risk of metabolic diseases (50). Within the ELSA-Brasil cohort in Waves 1 and 2, higher consumption of sugary drinks

was found to significantly increase the relative risk of metabolic syndrome, elevated fasting glucose levels, and high blood pressure. Additionally, moderate consumption of these drinks also increased the relative risk of elevated waist circumference in Brazilian adults (57). Similarly, soda and dietary fructose consumption were positively associated with a higher likelihood of hyperuricemia and elevated uric acid levels in Brazilian adults (58, 59).

Our findings underscore the importance of integrated public health strategies that address both physical activity and screen time to mitigate the consumption of unhealthy beverages. Additionally, considering that studies conducted within the ELSA-Brasil cohort showed that remaining physically active was associated with a 43% reduction in the risk of SARS-CoV-2 infection among those who also adopted specific COVID-19 protective practices (60), especially when combined with COVID-19 protective measures, public health interventions should promote active lifestyles and mindful eating habits, particularly during periods of social restriction. Given the significant health risks associated with high consumption of sugary drinks, policies aimed at reducing their consumption, such as taxation and public awareness campaigns, are crucial.

Among the limitations of the present study is its cross-sectional design, which precludes causal associations. To address this inherent limitation, we implemented statistical adjustments for key sociodemographic and lifestyle confounders, including age, income, smoking status, and alcohol consumption, using a hierarchical theoretical modeling approach. However, prospective cohort studies remain necessary to establish temporal relationships between physical activity, screen time behaviors, and beverage consumption patterns. Additionally, due to the restrictions and social distancing measures established during the supplementary COVID phase study in ELSA-Brasil during the pandemic, it was not possible to collect anthropometric data for the inclusion of body composition adjustment variables in the statistical models. To partially mitigate this constraint, we incorporated pre-pandemic data from ELSA-Brasil's Wave 3 for core sociodemographic variables and included remote work status as a proxy for pandemic-specific behavioral modifications. Future investigations should integrate anthropometric data from the next ELSA-Brasil study waves to enhance model precision.

Moreover, no quantitative data on food and beverage consumption were collected to estimate the specific quantities consumed of each type of drink and food, and thus caloric intake adjustments were not possible, complicating deeper analyses on possible dose-response effects of physical activity and screen time behavior on beverage consumption. Recently, a diet quality index was developed and validated to compare dietary consumption data across different waves of ELSA-Brasil, including the COVID wave (61), which is a potential line of analysis to further investigate the findings of the present study regarding the effects of physical activity and screen time behavior on diet quality. Screen time was a variable implemented in the supplementary COVID study, so longitudinal analyses to assess changes over time in screen time behavior across different waves are not possible. Nevertheless, our median-based cutoff (≤6 vs. >6 h/day) for screen time demonstrated conceptual alignment with previous Brazilian adult population studies, enhancing cross-study comparability. The composite physical activity/screen time variable enabled examination of synergistic behavioral effects, representing a methodological advancement in this research domain.

Despite these limitations, the present study provides valuable information on the combined behaviors of physical activity and screen

time and their relationship with sugary drink consumption in a cohort of Brazilian adults, given that most studies published to date on this topic focus on the child and adolescent population. Additionally, a separate analysis by beverage type was conducted, which is scarce in the current literature and allows for the identification of possible differentiated behaviors in beverage consumption habits. The findings highlight the need for public health strategies that promote active lifestyles and reduce screen time to mitigate the consumption of unhealthy beverages and their metabolic consequences. Moreover, the robust methodology and large sample size of the ELSA-Brasil study provide reliable data from a segment of the Brazilian population.

# 4.1 Practical applications and clinical relevance

The observed associations between the "Excessive Screen Time/ Physically Inactive" behavioral pattern and beverage consumption patterns during the COVID-19 pandemic highlight the importance of monitoring these behavioral interactions in similar public health crises. Our findings, documented under specific pandemic conditions including social distancing and remote work, provide evidence to guide surveillance strategies during periods of prolonged confinement or restricted mobility.

In clinical practice, the results suggest that during extended periods of enforced sedentary behavior, routine assessment of both screen time and physical activity levels could serve as indicators for identifying adults at higher risk of unhealthy beverage consumption patterns, potentially informing Health and Food and Nutritional Education strategies. The sex-specific differences we observed in beverage consumption associations indicate that gender-specific monitoring approaches may be warranted when evaluating behavioral changes during lockdowns or restricted mobility periods. These findings contribute empirical evidence to understanding how sudden lifestyle disruptions may influence dietary behaviors in adult populations.

## **5** Conclusion

Significant differences were observed between genders regarding sociodemographic and lifestyle variables. Beverage consumption patterns varied significantly according to these variables. Participants who consumed soft drinks or soda, sugar-sweetened beverages, and coffee with sugar generally had lower per capita incomes, higher screen time, and were physically inactive. Conversely, those consuming coffee without sugar had higher per capita incomes and tended to have higher screen time and physical inactivity.

Excessive screen time combined with physical inactivity was significantly associated with higher odds of consuming sugarsweetened artificial beverages among both men and women. Men with appropriate screen time but physically inactive behavior had nearly double the likelihood of consuming sugar-sweetened artificial beverages, while excessive screen time combined with physical inactivity significantly increased the likelihood of consuming these beverages among both men and women.

Additionally, excessive screen time and physical inactivity among men were associated with higher consumption of industrialized juices, whereas among women, this behavior increased the likelihood of consuming soft drinks or soda. No significant associations were found between physical activity and screen time behavior and the consumption of coffee, whether with or without sugar.

The demonstrated associations suggest public health strategies should prioritize integrated behavioral interventions that simultaneously target screen time reduction and physical activity promotion. Gender-specific approaches appear warranted, particularly programs addressing industrialized juice consumption in men and soda intake in women. For lower-income populations, where unhealthy beverage consumption was most prevalent, targeted nutritional support could help mitigate these patterns. The identified threshold of >6 h/day screen time combined with physical inactivity provides a clear benchmark for identifying high-risk groups needing intervention.

These findings offer evidence-based parameters for developing effective strategies to improve beverage consumption patterns. The study specifically highlights how addressing screen time and physical activity as interconnected behaviors, while considering gender and socioeconomic factors, could lead to more impactful public health interventions. The results provide measurable targets for program development and evaluation in community and clinical settings.

## Data availability statement

The original contributions presented in the study are included in the article/Supplementary material, further inquiries can be directed to the corresponding author.

## **Ethics statement**

The studies involving humans were approved by Research Ethics Committees of the Oswaldo Cruz Foundation (FIOCRUZ-RJ), Federal University of Espírito Santo (UFES), Federal University of Minas Gerais (UFMG), Federal University of Rio Grande do Sul (UFRGS), and Federal University of Bahia (UFBA). The studies were conducted in accordance with the local legislation and institutional requirements. The participants provided their written informed consent to participate in this study.

## Author contributions

YM: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Visualization, Writing – original draft, Writing – review & editing. HM: Investigation, Writing – original draft. TS: Conceptualization, Methodology, Supervision, Writing – review & editing. LB: Visualization, Writing – review & editing. LL: Writing – review & editing. MM: Conceptualization, Methodology, Supervision, Writing – review & editing.

# Funding

The author(s) declare that financial support was received for the research and/or publication of this article. This research was supported by the Ministry of Health (Science and Technology Department - DCIT) and the Ministry of Science, Technology, and Innovation in

Brazil (Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq), grant numbers 0106 0212.00 and 405551/2015-0 BA; 01 06 0300.00 and 405543/2015-8 ES; 0106 0278.00 and 405552/2015-7 MG; and 01 06 0071.00 and 405544/2015-4 RJ; 01 06 0010.00 and 405545/2015-0 RS; 01 06 0115.00 and 405547/2015-3 SP. Additionally, it was funded by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) through a doctoral scholarship awarded to the researcher YM, and by a Level 1C Research Productivity Fellowship from the CNPq awarded to the researcher MM. The publication of this article was funded by Fundação de Amparo à Pesquisa e Inovação do Espírito Santo (FAPES) and Universidad de las Américas Puebla.

### Acknowledgments

The authors thank the participants, researchers, and all collaborators of the ELSA-Brasil study for their invaluable contributions.

## **Conflict of interest**

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

### References

 Smirnova D, Syunyakov T, Pavlichenko A, Bragin D, Fedotov I, Filatova V, et al. Interactions between anxiety levels and life habits changes in general population during the pandemic lockdown: decreased physical activity, falling asleep late and internet browsing about COVID-19 are risk factors for anxiety, whereas social media use is. *Psychiatr Danub*. (2021) 33:119–29.

2. Kelley M, Ferrand RA, Muraya K, Chigudu S, Molyneux S, Pai M, et al. An appeal for practical social justice in the COVID-19 global response in low-income and middle-income countries. *Lancet Glob Heal*. (2020) 8:e888–9. doi: 10.1016/S2214-109X(20)30249-7

3. Bhutani S, Cooper JA. COVID-19–related home confinement in adults: weight gain risks and opportunities. *Obesity.* (2020) 28:1576–7. doi: 10.1002/oby.22904

4. Caputo EL, Reichert FF. Studies of physical activity and COVID-19 during the pandemic: A scoping review. J Phys Act Heal. (2020) 17:1275–84. doi: 10.1123/jpah.2020-0406

5. Malta DC, Szwarcwald CL, Barros MB de A, Gomes CS, Machado ÍE, de SJPRB, et al. A pandemia da COVID-19 e as mudanças no estilo de vida dos brasileiros adultos: um estudo transversal, 2020. *Epidemiol Serv Saúde*. (2020) 29:e2020407. doi: 10.1590/s1679-49742020000400026

6. Da Silva DRP, Werneck AO, Malta DC, De Souza Júnior PRB, Azevedo LO, De Azevedo Barros MB, et al. Changes in the prevalence of physical inactivity and sedentary behavior during COVID-19 pandemic: A survey with 39, 693 Brazilian adults. *Cad Saude Publica*. (2021) 37:e00221920. doi: 10.1590/0102-311X00221920

7. Molina MDCB, Aprelini CM de O, Meireles AL, Griep RH, Giatti L, Almeida M, et al. Impact of socioeconomic factors on lifestyle changes among employees of public higher education institutions in ELSA-Brasil during COVID-19 pandemic. *CAD Saude Publica*. (2023) 39:e00047123. doi: 10.1590/0102-311xen047123

8. Alamri ES. Effects of covid-19 home confinement on eating behavior: A review. J Public Health Res. (2021) 10:2088. doi: 10.4081/jphr.2021.2088

9. Herle M, Smith AD, Bu F, Steptoe A, Fancourt D. Trajectories of eating behavior during COVID-19 lockdown: longitudinal analyses of 22,374 adults. *Clin Nutr ESPEN*. (2021) 42:158–65. doi: 10.1016/j.clnesp.2021.01.046

10. Urquía YJM, Pereira TSS, Martinez OGE, Silva DA, Martins MCT, Molina M. Alimentación, estilo de vida y distanciamiento social: un estudio en el sureste brasilero durante la pandemia de COVID-19. *Población Salud Mesoam*. (2022) 20:49840. doi: 10.15517/psm.v20i1.49840

11. Giordani RCF, da SJAC, Da Silva JA. Editorial: eating behavior and mental health during the COVID-19 pandemic. *Front. Nutr.* (2023) 10:1295557. doi: 10.3389/fnut.2023.1295557

12. Di Renzo L, Gualtieri P, Pivari F, Soldati L, Attinà A, Cinelli G, et al. Eating habits and lifestyle changes during COVID-19 lockdown: an Italian survey. *J Transl Med.* (2020) 18:1–15. doi: 10.1186/s12967-020-02399-5

### **Correction note**

A correction has been made to this article. Details can be found at: 10.3389/fnut.2025.1674419.

### **Generative AI statement**

The authors declare that no Gen AI was used in the creation of this manuscript.

### Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

### Supplementary material

The Supplementary material for this article can be found online at: https://www.frontiersin.org/articles/10.3389/fnut.2025.1503010/ full#supplementary-material

13. Salazar-Fernández C, Palet D, Haeger PA, Mella FR. The perceived impact of COVID-19 on comfort food consumption over time: the mediational role of emotional distress. *Nutrients*. (2021) 13:1–12. doi: 10.3390/nu13061910

14. Booth L, Santos JA, Hoek AC, Webster J, Pettigrew S. Population subgroups at risk of unhealthy changes in food and beverage consumption during COVID-19 lockdowns. *Int J Behav Med.* (2024) 31:163–8. doi: 10.1007/s12529-023-10165-2

15. Park S, Lee SH, Blanck HM. Characteristics associated with being a high consumer of sweet foods and sugar-sweetened beverages among US adults during the COVID-19 pandemic, 2021. *Nutrients.* (2023) 15:1–11. doi: 10.3390/nu15102363

16. Prendergast KB, Mackay LM, Schofield GM. The clustering of lifestyle Behaviours in New Zealand and their relationship with optimal wellbeing. *Int J Behav Med.* (2016) 23:571–9. doi: 10.1007/s12529-016-9552-0

17. Santos AC, Willumsen J, Meheus F, Ilbawi A, Bull FC. The cost of inaction on physical inactivity to public health-care systems: a population-attributable fraction analysis. *Lancet Glob Heal*. (2023) 11:e32–9. doi: 10.1016/S2214-109X(22)00464-8

18. Santos RMS, Mendes CG, Sen Bressani GY, de Alcantara VS, de Almeida Nogueira YJ, de Miranda DM, et al. The associations between screen time and mental health in adolescents: a systematic review. *BMC Psychol.* (2023) 11:127. doi: 10.1186/s40359-023-01166-7

19. Nieber K. The impact of coffee on health. *Planta Med.* (2017) 83:1256-63. doi: 10.1055/s-0043-115007

20. Li B, Yan N, Jiang H, Cui M, Wu M, Wang L, et al. Consumption of sugar sweetened beverages, artificially sweetened beverages and fruit juices and risk of type 2 diabetes, hypertension, cardiovascular disease, and mortality: A meta-analysis. *Front Nutr.* (2023) 10:1019534. doi: 10.3389/fnut.2023.1019534

21. Leme ACB, Ferrari G, Fisberg RM, Kovalskys I, Gómez G, Cortes LY, et al. Cooccurrence and clustering of sedentary behaviors, diet, sugar-sweetened beverages, and alcohol intake among adolescents and adults: the Latin American nutrition and health study (ELANS). *Nutrients*. (2021) 13:1809. doi: 10.3390/nu13061809

22. Rocha LL, Pessoa MC, Gratão LHA, do Carmo AS, de Freitas Cunha C, TRPR d O, et al. Health behavior patterns of sugar-sweetened beverage consumption among Brazilian adolescents in a nationally representative school-based study. *PLoS One.* (2021) 16:e0245203. doi: 10.1371/journal.pone.0245203

23. Nour TY, Altintaş KH. Effect of the COVID-19 pandemic on obesity and it is risk factors: a systematic review. *BMC Public Health.* (2023) 23:1–24. doi: 10.1186/s12889-023-15833-2

24. Salman A, Sigodo KO, Al-Ghadban F, Al-Lahou B, Alnashmi M, Hermassi S, et al. Effects of COVID-19 lockdown on physical activity and dietary behaviors in Kuwait: A cross-sectional study. *Nutrients*. (2021) 13:2252. doi: 10.3390/nu13072252

25. Schmidt MI, Duncan BB, Mill JG, Lotufo PA, Chor D, Barreto SM, et al. Cohort profile: longitudinal study of adult health (ELSA-Brasil). *Int J Epidemiol.* (2015) 44:68–75. doi: 10.1093/ije/dyu027

26. Aquino EML, Barreto SM, Bensenor IM, Carvalho MS, Chor D, Duncan BB, et al. Brazilian longitudinal study of adult health (ELSA-Brasil): objectives and design. *Am J Epidemiol [Internet]*. (2012) 175:315–24. doi: 10.1093/aje/kwr294

27. Aquino EML, Vasconcellos-Silva PR, Coeli CM, Araújo MJ, Santos SM, de Figueiredo RC, et al. Aspectos éticos em estudos longitudinais: o caso do ELSA-Brasil. *Rev Saude Publ.* (2013) 47:19–26. doi: 10.1590/S0034-8910.2013047003804

28. Petruccelli JL, Saboia AL. Características Étnico-raciais da População: Classificações e Identidades [Internet]. Características étnico-raciais da população. Clasificações e Identidades. Rio de Janeiro: Instituto Brasileiro de Geografia e Estatística - IBGE (2013).

29. Matsudo S, Araújo T, Matsudo V, Andrade D, Andrade E, Oliveira LC, et al. Questionário Internacional de Atividade Física (Ipaq): Estudo de validade e reprodutibilidade no Brasil. *Rev Bras Atividade Física Saúde*. (2001) 6:5–18. doi: 10.12820/rbafs.v.6n2p5-18

30. WHO. World Health Organization. WHO guidelines on physical activity and sedentary behaviour. Geneva: WHO (2020).

31. Silva DR, Collings P, Araujo RHO, Barboza LL, Szwarcwald CL, Werneck AO. Correlates of screen-based behaviors among adults from the 2019 Brazilian National Health Survey. *BMC Public Health.* (2021) 21:2289–9. doi: 10.1186/s12889-021-12340-0

32. Victora CG, Huttly SR, Fuchs SC, Olinto MTA. The role of conceptual frameworks in epidemiological analysis: a hierarchical approach. *Int J Epidemiol*. (1997) 26:224–7. doi: 10.1093/ije/26.1.224

33. Rehm CD, Ratliff JC, Riedt CS, Drewnowski A. Coffee consumption among adults in the United States by demographic variables and purchase location: analyses of NHANES 2011-2016 data. *Nutrients*. (2020) 12:1–13. doi: 10.3390/nu12082463

34. Fontes AS, Pallottini AC, Vieira DADS, de M FM, Marchioni DM, CLG C, et al. Demographic, socioeconomic and lifestyle factors associated with sugar-sweetened beverage intake: a population-based study. *Rev Bras Epidemiol.* (2020) 23:e200003. doi: 10.1590/1980-549720200003

35. Pereira RA, Souza AM, Duffey KJ, Sichieri R, Popkin BM. Beverage consumption in Brazil: results from the first National Dietary Survey. *Public Health Nutr.* (2015) 18:1164–72. doi: 10.1017/S1368980014001657

36. Darmon N, Drewnowski A. Contribution of food prices and diet cost to socioeconomic disparities in diet quality and health: a systematic review and analysis. *Nutr Rev.* (2015) 73:643–60. doi: 10.1093/nutrit/nuv027

37. Silva DTC, Prado WL, Cucato GG, Correia MA, Ritti-Dias RM, Lofrano-Prado MC, et al. Impact of COVID-19 pandemic on physical activity level and screen time is associated with decreased mental health in Brazillian adults: A cross-sectional epidemiological study. *Psychiatry Res.* (2022) 314:114657. doi: 10.1016/j.psychres.2022.114657

38. McNaughton SA, Pendergast FJ, Worsley A, Leech RM. Eating occasion situational factors and sugar-sweetened beverage consumption in young adults. *Int J Behav Nutr Phys Act.* (2020) 17:1–12. doi: 10.1186/s12966-020-00975-y

39. Moradell A, Santaliestra-Pasías AM, Aparicio-Ugarriza R, Huybrechts I, Bertalanné Szommer A, Forsner M, et al. Are physical activity and sedentary screen time levels associated with food consumption in European adolescents? The HELENA study. *J Am Nutr Assoc.* (2023) 42:55–66. doi: 10.1080/07315724.2021.1978900

40. Nagata JM, Magid HSA, Gabriel KP. Screen time for children and adolescents during the coronavirus disease 2019 pandemic. *Obesity*. (2020) 28:1582–3. doi: 10.1002/oby.22917

41. Tebar WR, Christofaro DGD, Diniz TA, Lofrano-Prado MC, Botero JP, Correia M, et al. Increased screen time is associated with alcohol desire and sweetened foods consumption during the COVID-19 pandemic. *Front Nutr.* (2021) 8:630586. doi: 10.3389/fnut.2021.630586

42. Cardoso PC, Caldeira TCM, de STM, Claro RM. Changes in screen time in Brazil: A time-series analysis 2016-2021. *Am J Health Promot.* (2023) 37:681–4. doi: 10.1177/08901171231152147

43. de O da S SP, Griep RH, FJG P, Barreto SM, SMA M, de JM da FM. Work from home and the association with sedentary behaviors, leisure-time and domestic physical activity in the ELSA-Brasil study. *BMC Public Health.* (2023) 23:305–10. doi: 10.1186/s12889-023-15167-z

44. Dartora WJ, Schmidt MI, Griep RH, Duncan BB. Changes in lifestyle habits in individuals with diabetes during the COVID-19 pandemic: the ELSA-Brasil cohort study. *COVID*. (2023) 3:1601–11. doi: 10.3390/covid3100109

45. Vizcaino M, Buman M, Desroches T, Wharton C. From TVs to tablets: the relation between device-specific screen time and health-related behaviors and characteristics. *BMC Public Health.* (2020) 20:1–10. doi: 10.1186/s12889-020-09410-0

46. García-Laguna G, Martínez-Rodríguez TY, Ramos-Caballero DM, Bernal-Gómez SJ. La actividad física regula la ingesta de bebidas azucaradas y la velocidad al comer en adultos jóvenes de una institución de educación superior en Bogotá-Colombia. *Nutr Clínica Dietética Hosp.* (2020) 41:171–7. doi: 10.12873/413martinez

47. Cleland VJ, Patterson K, Breslin M, Schmidt MD, Dwyer T, Venn AJ. Longitudinal associations between TV viewing and BMI not explained by the "mindless eating" or "physical activity displacement" hypotheses among adults. *BMC Public Health*. (2018) 18:1–12. doi: 10.1186/s12889-018-5674-4

48. De Rezende LFM, Lopes MR, Rey-Lopez JP, Matsudo VKR, Luiz ODC. Sedentary behavior and health outcomes: an overview of systematic reviews. *PLoS One.* (2014) 9:e105620. doi: 10.1371/journal.pone.0105620

49. Turi BC, Monteiro HL, Lemes R, Codogno JS, Lynch KR, Asahi Mesquita CA, et al. TV viewing time is associated with increased all-cause mortality in Brazilian adults independent of physical activity. *Scand J Med Sci Sports.* (2018) 28:596–603. doi: 10.1111/sms.12882

50. Malik VS, Popkin BM, Bray GA, Després JP, Willett WC, Hu FB. Sugar-sweetened beverages and risk of metabolic syndrome and type 2 diabetes: a meta-analysis. *Diabetes Care.* (2010) 33:2477–83. doi: 10.2337/dc10-1079

51. Drouin-Chartier JP, Zheng Y, Li Y, Malik V, Pan A, Bhupathiraju SN, et al. Changes in consumption of sugary beverages and artificially sweetened beverages and subsequent risk of type 2 diabetes: results from three large prospective U.S. *Cohorts Women Men Diabetes Care.* (2019) 42:2181–9. doi: 10.2337/dc19-0734

52. Kristal RB, Blank AE, Wylie-Rosett J, Selwyn PA. Factors associated with daily consumption of sugar-sweetened beverages among adult patients at four federally qualified health centers, Bronx, New York, 2013. *Prev Chronic Dis.* (2014) 11:342. doi: 10.5888/pcd11.140342

53. Drummen M, Dorenbos E, Vreugdenhil ACE, Raben A, Westerterp-Plantenga MS, Adam TC. Insulin resistance, weight, and behavioral variables as determinants of brain reactivity to food cues: a prevention of diabetes through lifestyle intervention and population studies in Europe and around the world – a PREVIEW study. *Am J Clin Nutr.* (2019) 109:315–21. doi: 10.1093/ajcn/nqy252

54. Sigala DM, Stanhope KL. An exploration of the role of sugar-sweetened beverage in promoting obesity and health disparities. Curr Obes Rep. (2021) 10:39–52. doi: 10.1007/s13679-020-00421-x

55. Siqueira JH, Santana NMT, Pereira TSS, Moreira AD, Benseñor IM, Barreto SM, et al. Consumo de bebidas alcoólicas e não alcoólicas: Resultados do ELSA-Brasil. *Cien Saude Colet.* (2021) 26:3825–37. Available from:. doi: 10.1590/1413-81232021269.2.30682019

56. Malik VS, Hu FB. Sugar-sweetened beverages and Cardiometabolic health: an update of the evidence. *Nutrients*. (2019) 11:1840. doi: 10.3390/nu11081840

57. Siqueira JH, Pereira TSS, Moreira AD, Diniz MFHS, Velasquez-Melendez G, Fonseca MJM, et al. Consumption of sugar-sweetened soft drinks and risk of metabolic syndrome and its components: results of the ELSA-Brasil study (2008–2010 and 2012–2014). *J Endocrinol Invest.* (2023) 46:159–71. doi: 10.1007/s40618-022-01895-3

58. Siqueira JH, Mill JG, Velasquez-Melendez G, Moreira AD, Barreto SM, Benseñor IM, et al. Sugar-sweetened soft drinks and fructose consumption are associated with hyperuricemia: cross-sectional analysis from the Brazilian longitudinal study of adult health (ELSA-Brasil). *Nutrients.* (2018) 10:981. doi: 10.3390/nu10080981

59. Siqueira JH, Pereira TSS, Velasquez-Melendez G, Barreto SM, Benseñor IM, Mill JG, et al. Sugar-sweetened soft drinks consumption and risk of hyperuricemia: results of the ELSA-Brasil study. *Nutr Metab Cardiovasc Dis.* (2021) 31:2004–13. doi: 10.1016/j.numecd.2021.04.008

60. Pitanga FJG, da C AM, Duncan BB, Mill JG, Giatti L, del CB MM, et al. Leisure time physical activity and SARS-CoV-2 infection among ELSA-Brasil participants. *Int J Environ Res Public Health*. (2022) 19:1–9. doi: 10.3390/ijerph192114155

61. Oliveira-Aprelini CM, Herzog-Siqueira J, Xavier-Martins H, Silva-Pereira TS, Mercadante-Urquía YJ, Bisi-Molina M. Desarrollo y validación de la puntuación de calidad alimentaria en el ELSA-Brasil. *Nutr Hosp.* (2024) 41:175–85. doi: 10.20960/ nh.04598