### Check for updates

### **OPEN ACCESS**

APPROVED BY Frontiers Editorial Office, Frontiers Media SA, Switzerland

\*CORRESPONDENCE Frontiers Production Office production.office@frontiersin.org

RECEIVED 23 June 2025 ACCEPTED 24 June 2025 PUBLISHED 08 July 2025

#### CITATION

Frontiers Production Office (2025) Correction: Brown-shell eggs shows high incidence of blood and meat spots accompanied by unique microbial distribution patterns. *Front. Nutr.* 12:1652275. doi: 10.3389/fnut.2025.1652275

### COPYRIGHT

© 2025 Frontiers Production Office. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

# Correction: Brown-shell eggs shows high incidence of blood and meat spots accompanied by unique microbial distribution patterns

## Frontiers Production Office\*

Frontiers Media SA, Lausanne, Switzerland

### KEYWORDS

chicken eggs, egg quality, Egg contents, blood and meat spot, microbiota

## A Correction on

Brown-shell eggs shows high incidence of blood and meat spots accompanied by unique microbial distribution patterns

by Wu, J., Yan, Y., Chen, J., Li, J., Li, G., Wu, G., Wang, B., Zheng, G., Yang, Y., Du, Y., and Lian, L. (2025). *Front. Nutr.* 12:1561194. doi: 10.3389/fnut.2025.1561194

In the published article, there was an error in Figure 1 as published. The position of labels "WL\_CON (n = 124)" and "RIR\_CASE (n = 51)" in Figure 1 were erroneously reversed. The corrected Figure 1 and its caption appear below.

The original version of this article has been updated.

