



BRCA1 Versus BRCA2 and PARP Inhibitors Efficacy in Solid Tumors: A Meta-Analysis of Randomized Controlled Trials

Shan Li^{1†}, Li Tao^{2†}, Haiyun Dai³, Xue Gong⁴, Yuguo Zhuo¹, Hui Xiang⁵, Yueyang Zhao⁶, Qing Gao^{1*‡} and Liang Deng^{1*‡}

OPEN ACCESS

Edited by:

Alice Chen,
National Cancer Institute (NCI),
United States

Reviewed by:

Luis E. Arias-Romero,
National Autonomous University of
Mexico, Mexico
Yogesh Chawla,
Mayo Clinic, United States

*Correspondence:

Liang Deng
dengliang@cqmu.edu.cn
Qing Gao
981051852@qq.com

[†]These authors have contributed
equally to this work and
share first authorship

[‡]These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Cancer Molecular Targets
and Therapeutics,
a section of the journal
Frontiers in Oncology

Received: 01 June 2021

Accepted: 12 October 2021

Published: 28 October 2021

Citation:

Li S, Tao L, Dai H, Gong X, Zhuo Y,
Xiang H, Zhao Y, Gao Q and Deng L
(2021) BRCA1 Versus BRCA2 and
PARP Inhibitors Efficacy in Solid
Tumors: A Meta-Analysis of
Randomized Controlled Trials.
Front. Oncol. 11:718871.
doi: 10.3389/fonc.2021.718871

¹ Department of Gastroenterology, The First Affiliated Hospital of Chongqing Medical University, Chongqing, China,

² Department of Radiology, The First Affiliated Hospital of Chongqing Medical University, Chongqing, China, ³ Department of Respiratory and Critical Care Medicine, The First Affiliated Hospital of Chongqing Medical University, Chongqing, China,

⁴ Department of Gynecology, The First Affiliated Hospital of Chongqing Medical University, Chongqing, China, ⁵ Center for Disease Control and Prevention of Nan'an District, the Department for Infectious Disease Prevention and Control, Chongqing, China, ⁶ Neuroscience Research Center, Chongqing Medical University, Chongqing, China

Background: BRCA2 mutation has a more substantial impact on the homologous recombination and superior therapeutic response to platinum-based chemotherapy than BRCA1 mutation. Whether BRCA2-mutated patients could benefit more from PARPi than BRCA1-mutated patients remains unclear. We performed a meta-analysis to assess the efficacy difference of PARPi between BRCA1 mutation carriers and BRCA2 mutation carriers.

Methods: Pubmed, Embase, and Cochrane Library were comprehensively searched for randomized controlled trials (RCTs) of PARPi that had available hazard ratios (HRs) of progression-free survival (PFS) in both BRCA1-mutated population and BRCA2-mutated population. We calculated the pooled PFS HRs and 95%CI using randomized-effect models, and the difference between the two estimates was compared by interaction test.

Results: A total of 11 eligible RCTs of high quality were identified through search. Overall, 1544 BRCA1 mutation carriers and 1191 BRCA2 mutation carriers were included in the final analysis. The pooled PFS HR was 0.42 (95% CI: 0.35-0.50) in BRCA1-mutated patients who were treated with PARPi compared with patients in the control group. In BRCA2-mutated patients treated with PARPi, the pooled PFS HR compared with the control groups was 0.35 (95% CI: 0.24-0.51). The difference in efficacy of PARPi was not significant between the two subgroups ($P_{\text{heterogeneity}} = 0.40$, for interaction).

Conclusion: BRCA1-mutated patients and BRCA2-mutated patients could benefit from PARPi, and the efficacy is comparable. Currently, there is no evidence that BRCA2-mutated patients would benefit more from PARPi than BRCA1-mutated patients.

Systematic Review Registration: <https://www.crd.york.ac.uk/PROSPERO/>, identifier CRD42020214582.

Keywords: PARPi, BRCA1, BRCA2, meta-analysis, progression-free survival

INTRODUCTION

BRCA1 and BRCA2 are two critical genes involved in DNA repair *via* homologous recombination (1, 2). Cells with mutations in BRCA1 and BRCA2 are unable to activate error-free homologous recombination to repair DNA double-strand breaks, as this can result in genomic instability and even a predisposition to malignant transformation (3, 4). Many previous studies reported BRCA mutations significantly increased the susceptibility to various cancer, including breast cancer, ovarian cancer, prostate cancer, pancreatic cancer, and colon cancer (5–8).

Interestingly, BRCA-mutated cancer cells' ability to repair DNA damage would also be impaired by the same defect, as they would become hyperdependent on the remaining repair pathway. Therefore, BRCA-mutated individuals were more sensitive to treatments relying on the induction of DNA damage, such as platinum-based chemotherapy and radiotherapy. Clinical effects of BRCA1 and BRCA2 mutations have been analyzed commonly together, but many studies have found that compared with BRCA1 mutation carriers, BRCA2 mutation carriers were associated with improved platinum-based chemotherapy response and longer progression-free duration (18.0 months for BRCA2 vs. 12.5 months for BRCA1, $P=0.04$) (9–12). Therefore, it is becoming increasingly apparent that these mutations do not have the same effects, as some studies proved that BRCA2 might have a stronger association with homologous recombination and a hypermutator phenotype (9, 11).

Targeting DNA repair in cancer, poly (ADP-ribose) polymerase (PARP) inhibitors (PARPi) were the first clinically approved target therapy, and these promising drugs revolutionarily changed the therapeutic strategies in various cancers, including breast cancer, ovarian cancer, prostate cancer, pancreatic cancer (13–24). Similarly, accumulated evidence indicated that malignancies arising in patients with BRCA mutations are sensitive to PARPi because they have DNA repair defects, as mentioned before (25). Interestingly, in a recent trial (POLO), a favorable progression-free survival (PFS) hazard ratio (HR: 0.40, 95%CI 0.20–0.85) was seen in the BRCA1 mutated patients treated with olaparib for pancreatic cancer compared with placebo, but in BRCA2 mutated patients, the benefit was not significant (HR: 0.63, 95% CI 0.39–1.02) (22). By contrast, in another trial (Profound), the efficacy of PARPi was not significant in BRCA1 mutated carriers (HR: 0.41, 95% CI 0.13–1.39), while in BRCA2 mutated patients, the benefit was significant (HR: 0.21, 95% CI 0.13–0.32) (24). Therefore, whether patients with BRCA1 mutation and patients with BRCA2 mutation can both benefit from PARPi and whether these mutations have similar prognostic effects in the use of PARPi remained unclear and controversial.

Considering the currently high costs of PARPi (\$250K per PFS life-year), it is of great importance for physicians to decide which subgroup would potentially benefit more from PARPi (26). In this scenario, we performed a meta-analysis to evaluate and compare the efficacy of treatments with PARPi in patients with BRCA1 mutation and patients with BRCA2 mutation.

METHOD

We followed the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines to conduct our meta-analysis (27). Our meta-analysis has been registered on PROSPERO, and the CRD code was CRD42020214582.

Search Strategy and Selection Criteria

We comprehensively searched PubMed, Embase, and Cochrane Library for phase 2 and 3 randomized controlled trials published from the inception of each database to October 2020 with no language restrictions. The following keywords were used: niraparib, rucaparib, talazoparib, olaparib, veliparib, (poly(ADP-ribose) polymerase inhibitors, PARP Inhibitors, and randomized controlled trial (see **Supplementary Materials, Table 1**).

To be eligible, randomized controlled trials (RCT) had to fulfill all following criteria: (1) phase 2 or 3 randomized controlled trials; (2) trials evaluating the relative efficacy of a PARPi alone or with other regimens compared with regimens that did not include a PARPi in patients with cancer; (3) studies reporting HR and 95% credible interval (CI) of PFS in both BRCA1-mutated and BRCA2-mutated subgroups, regardless of mutation types (germline or somatic). We excluded trials that: (1) were single-arm or non-randomized trials; (2) were retrospective or phase 1 trials; (3) did not present HR of PFS in both subgroups of patients with BRCA1 mutation and patients with BRCA2 mutation; (4) presented HR of PFS in two subgroups, including data from patients with both BRCA1 and BRCA2 mutations in the analysis.

Moreover, in case of trials that did not include survival subgroup analysis according to BRCA mutation status in the main text, each study's supplement would be carefully reviewed in the process of at the full-text screening stage. Additionally, if the same trial appeared in different publications, only the most complete or updated one would be included. The above screening and selection work would be done by two independent investigators. Subsequently, any discrepancies in the study selection process were discussed and resolved by a consensus formed by all investigators involved.

Risk of Bias Assessment and Data Extraction

Two investigators would assess the quality of potentially eligible studies according to The Cochrane Risk of Bias Tool (27). Every potentially included study would be evaluated by the following criteria: (1) randomized sequence generation; (2) allocation concealment; (3) blinding of participants, personnel; (4) blinding of outcome assessment; (5) incomplete outcome data; (6) selective outcome reporting, and (7) other sources of bias. Based on this method, each risk of bias was described as low risk, high risk, or unclear risk.

From each included publications, name of study, study phase, trial design, underlying malignancy, number of patients, line of therapy, and HR for progression according to patients' BRCA mutation types would be collated. Again, any disagreements were discussed and resolved by the consensus formed by all investigators.

Statistical Analysis

The primary endpoint was the progression-free survival in patients with BRCA1 mutation and patients with BRCA2 mutation measured by hazard ratios. Accordingly, we derived the hazard ratios for progression and their 95% confidence intervals from each study, separately for BRCA1-mutated patients and BRCA2-mutated patients. We assessed the heterogeneity between different trials by the Cochrane's Q statistic and I^2 statistic. Because of the clinical heterogeneity inherent in the data, we utilized random-effect models to calculate the pooled HR of PFS in all analyses. Finally, differences in pooled HRs between the two subgroups were calculated by interaction test, described by Altman and Bland (28). Accordingly, to perform the comparison of the two estimated quantities, the two estimates should be independent, not obtained from the same individuals (28). Therefore, patients with both BRCA1 and BRCA2 mutations were excluded from the analysis. Additionally, considering the relatively small sample of BRCA1 mutation carriers included in PROfound and POLO trials, within-trial interaction (deft approach) alone might not be appropriate (22, 24, 29). Therefore, we combined both across-trial and within-trial interactions to improve the power of the interaction test (29).

Subgroup analysis was also conducted to investigate the variation of the effect of BRCA mutation status on the PARPi efficacy. We only considered subgroups including more than two studies, as different cancer types, lines of therapy, and study methodology were selected. Additionally, since the number of patients in PROfound and POLO was limited, we performed another subgroup analysis, pooling the other 9 trials (22, 24). Besides, publication bias was assessed by Egger's test and Begg's test (30–32). Three investigators performed the statistical analyses by STATA 12.0 and R studio (metafor package), and all the data were expressed as the combination of HR and 95% CI. Moreover, a two-tailed $P < .05$ was regarded as statistically significant in the two-tailed test.

RESULTS

Search Results and Patient Characteristics

Initially, 3422 publications were identified through the database search. After duplication, title and abstract screen, and full-text review, a total of 11 RCT were included in the final analysis (14–24, 33) (see **Figure 1**). Additionally, we excluded Study19 to assess within-trial interaction because this trial only provided HR of PFS for patients with BRCA1 mutation while the data for BRCA2 was not available (34). At last, only one study was a phase 2 study among eligible studies, while the remaining ten trials were designed as phase 3 RCTs. All current clinical proved PARPi were included in this meta-analysis (six in olaparib, two in veliparib, and one each in talazoparib, niraparib, and rucaparib). Six trials were done in patients with ovarian cancer, three trials in patients with breast cancer, one each in pancreatic adenocarcinoma and prostate cancer.

Table 1 listed the main characteristics of the 11 included RCTs. Overall, 2747 BRCA-deficient patients were included, of which 1544 (56.2%) were BRCA1 mutation carriers, 1191 (43.4%) were BRCA2 mutation carriers, and 12 (0.4%) were patients with both mutations. The BRCA1/2 mutations were confirmed using BRCAAnalysis CDx (Myriad) in ten trials, and five patients in SOLO1 with BRCA mutations which BGI confirmed, were excluded in the analysis of PFS in the SOLO1 trial (14–23). Besides, in the PROfound trial, FoundationOne CDx was utilized instead, as both BRCAAnalysis CDx and FoundationOne CDx were approved by FDA for BRCA mutation testing (24, 35, 36).

Risk of Bias

The method quality of the included trials was generally moderate to good (see **Supplementary Material, Table 2**). Randomized treatment allocation sequences were generated in all trials. The main issue affecting quality was the lack of blinding because three trials (EMBRACA, OlympiAD, and PROfound) were open-labeled (15, 21, 24).

Efficacy of PARP Inhibitors and BRCA Mutation Status

Overall, BRCA1-mutated patients who were treated with PARPi had a significantly reduced risk of progression (HR: 0.42, 95% CI: 0.35–0.50) compared with patients in the control group (see **Figure 2**). Similarly, BRCA2-mutated patients treated with PARPi were also associated with better progression-free survival than controls (HR: 0.35, 95% CI: 0.24–0.51). It should be noted that the difference in efficacy of PARPi was not significant between the two subgroups ($Z=0.84$, $P_{\text{heterogeneity}} = 0.40$, for interaction). Notably, patients with both BRCA1 and BRCA2 mutations were excluded in all statistical analyses to make the two estimates from BRCA1 and BRCA2 subgroups independent for the interaction test.


Subgroup Analysis

Subgroup analyses were performed according to cancer types, study methodology, and lines of the therapy. Again, no statistically significant differences in the efficacy of PARPi were found between BRCA1-mutated patients and BRCA2-mutated patients in any of these analyses (**Figure 3**). Both BRCA1 mutation carriers and BRCA2 mutation carriers could significantly benefit from PARPi regardless of cancer types and different therapeutic lines. In another subgroup analysis including solely breast cancer and ovarian cancer, excluding pancreatic and prostate cancer, a similar outcome was seen.

Interestingly, compared with BRCA2-mutated patients who were treated with chemotherapy or other standard regimens, the benefit of the addition of PARPi seemed marginal for BRCA2-mutated patients (HR: 0.51, 95%CI: 0.25–1.05). The possible explanation might be the limited number of patients involved.

Publication Bias

The Begg's test ($P=0.34$) and Egger's test ($P=0.21$) showed no significant publication bias in the meta-analysis.



DISCUSSION

To our knowledge, this study is the first meta-analysis focus on the efficacy difference of PARPi between BRCA1-mutated patients and BRCA2-mutated patients. With published data from 11 high-quality RCTs for more than 2700 patients, our pooled analysis indicated that both BRCA1 and BRCA2 mutated patients with solid tumors could benefit from PARPi, and the efficacies were comparable.

Previous studies reported that BRCA2 mutation was associated with better platinum-based chemotherapy response and a significantly longer survival time than BRCA1 mutation (10–12, 37, 38). Interestingly, these researchers believed that this divergence of therapy response might relate to the different nature of these two gene mutations. BRCA2 was directly responsible for regulating RAD51 protein, which was essential in the double-strand break repair by homologous recombination (9, 11). In contrast, besides DNA damage repair, BRCA1 was deemed as a scaffold protein and played more diverse roles, including checkpoint control and mitotic spindle assembly, as the dysfunction of BRCA1 might not necessarily affect homologous recombination (11, 39, 40). From there, the BRCA2-deficient cancer cells might be more vulnerable to DNA-damaging agents such as platinum-based chemotherapy and thereby revealing improved clinical outcomes.

Consistent with the findings from platinum-based chemotherapy, some preclinical trials studying PARP inhibition also demonstrated similar results. One preclinical study demonstrated that, although BRCA1-deficient and BRCA2-deficient cells were both sensitive to PARP inhibition, compared with wild-type cells, BRCA1-deficient cells showed a 57-fold increase in the sensitivity, while BRCA2-deficient cells indicated a 133-fold enhanced sensitivity (41).

Interestingly, our study showed a different result, in a clinical setting, that BRCA1-mutated patients and BRCA2-mutated patients could equally benefit from the treatments with PARPi. There could be several explanations. Firstly, the clinicopathological features of the two subgroups might not be the same. For instance, some studies found that among patients with ovarian cancer, compared with BRCA2 mutation carriers, BRCA1-mutated patients were diagnosed at an earlier age (11, 12). Besides, some other researches focusing on breast cancer indicated that BRCA2-associated breast cancer showed more malignant features on imaging but BRCA1-mutated counterpart showed more aggressive pathological features such as high grade (42, 43). Whether these clinical features could be the contributing factors to the efficacy of the PARPi remains unknown. Therefore, a future individual patient data meta-analysis would be highly valuable. Secondly, not all BRCA1 mutations were equal, and some of the

TABLE 1 | The basic characteristics and main outcomes of the 11 included randomized controlled trials.

Study ID	Phase	Treatment groups		Tumor types	‡Number of patients (Exp/Con)			Line of therapy	Progression-free Survival	
		Intervention	Control		BRCA1	BRCA2	Both		HR (95% CI) for BRCA1	HR (95% CI) for BRCA2
SOLO1	3	Olaparib	Placebo	ovarian cancer	191/91	66/40	3/0	Maintenance after first line	0.40 (0.29–0.56)	0.20 (0.10–0.38)
EMBRACA	3	Talazoparib	*Physician's Choice Chemotherapy	breast cancer	133/63	154/81	0/0	≥1	0.59 (0.39–0.90)	0.47 (0.32–0.70)
VELIA/ GOG-3005	3	Veliparib plus #Chemotherapy followed by Veliparib	Placebo plus #Chemotherapy followed by placebo	ovarian cancer	78/59	30/31	0/2	First-line + Maintenance	0.38 (0.23–0.63)	0.64 (0.27–1.56)
BROCADE	2	Veliparib plus #Chemotherapy	Placebo plus #Chemotherapy	breast cancer	51/53	44/46	0/0	≥1	0.745 (0.454–1.224)	0.783 (0.433–1.417)
NOVA	3	Niraparib	Placebo	ovarian cancer	84/43	50/18	1/0	Maintenance after ≥ 2 lines	0.38 (0.23–0.61)	0.12 (0.04–0.31)
SOLO2	3	Olaparib	Placebo	ovarian cancer	132/61	58/35	0/0	Maintenance after ≥ 2 lines	0.30 (0.21–0.43)	0.36 (0.22–0.62)
ARIEL3	3	Rucaparib	Placebo	ovarian cancer	80/37	50/29	0/0	Maintenance after ≥ 2 lines	0.32 (0.19–0.53)	0.12 (0.06–0.26)
OlympiAD	3	Olaparib	*Physician's Choice Chemotherapy	breast cancer	117/51	84/46	4/0	≥1	0.54 (0.37–0.79)	0.68 (0.45–1.07)
POLO	3	Olaparib	Placebo	pancreatic cancer	29/16	62/46	1/0	Maintenance after first line	0.40 (0.20–0.85)	0.63 (0.39–1.02)
PAOLA-1	3	Olaparib plus bevacizumab	Placebo plus bevacizumab	ovarian cancer	111/49	45/31	1/0	Maintenance after first line	0.29 (0.176–0.470)	0.23 (0.090–0.541)
PROfound	3	Olaparib	Physician's choice (enzalutamide or abiraterone)	prostate cancer	10/5	92/53	0/0	≥2	0.41 (0.13–1.39)	0.21 (0.13–0.32)

*Physician's Choice Chemotherapy: (Capecitabine, Eribulin, Gemcitabine, or Vinorelbine) in EMBRACA, and (Capecitabine, Vinorelbine, or Eribulin) in OlympiAD; #Chemotherapy: (Carboplatin and Paclitaxel) in VELIA and BROCADE; EXP, interventional arm; CON, control arm; HR, hazard ratios; CI, confidence interval; ‡Number of patients: patient with both BRCA1 and BRCA2 mutations included in "Both" group, but excluded in "BRCA1" or "BRCA2" arms and excluded in the later pooled analysis.

mutation locations might result in hypomorphic BRCA1 isoforms, which could affect the efficacy of DNA-damaging agents. For example, cancer cells carrying the BRCA1(C61G) mutation or BRCA1 exon 11 respond poorly to platinum drugs and olaparib (44). On the other hand, the genetic interactions between BRCA1 and BRCA2 should also be noted (45). Some studies claimed that BRCA1 and BRCA2 might also have complementary functions, as a specific BRCA1 mutation (mutations on the C-terminal region of BRCA1) could affect the function of BRCA2 through PALB2 and result in similar clinical outcomes like BRCA2 mutation (12, 46, 47). Notably, the above evidence was all from preclinical trials. However, there could be a potential pitfall for therapeutic stratification of PARPi while all BRCA1-mutated patients continue to be considered a single clinical entity. Whether patients with such BRCA1 mutation locations could also benefit from PARPi in a clinical setting was still unclear. Hence, future studies with more detailed stratifications within the BRCA1-mutated population would be urgently needed.

Another explanation might be related to other therapeutic actions of PARPi and other functions of BRCA1 beyond DNA damage. In one recent meta-analysis studying the efficacy of PARPi in newly diagnosed ovarian cancer, a significant clinical benefit of PARPi has been obtained both in Non-BRCA mutated patients and in patients even without homologous recombination deficiency (48). In fact, besides DNA damage-induced cell death, PARP inhibitors, by inhibiting the PARP-1, can alter cellular energy metabolism and redox balance, leading to cancer cell apoptosis (49). Unlike

BRCA2, which directly guided RAD51 to damage sites in the process of DNA repair, BRCA1 plays more diverse roles. Besides DNA repair, BRCA1 also played critical roles in checkpoint control, mitotic spindle assembly, sister chromatid decatenation, and centrosome duplication (9, 39). From there, the interaction between PARPi and the above functions of BRCA1 beyond DNA damage is less known and worth vast further investigation.

The major limitation of this meta-analysis was the considerable heterogeneity observed in the pooled results. It is highly likely that such heterogeneity was related to the design of the study itself, including different lines, conduction on different solid tumors, and different study methodology. Therefore, a randomized effect model was applied to take into account such heterogeneity. More importantly, numerous subgroup analyses based on the above factors were performed, and the results were consistent. No statistically significant differences in the efficacy of PARPi were found between BRCA1 mutation carriers and BRCA2 mutation carriers in any of these analyses. Secondly, an optimal treatment strategy is needed to maximize the benefit as well as minimize the risk of toxicities. However, in our study, the information regarding adverse events from the two subgroups separately was not available. The safety difference between BRCA1-mutated patients and BRCA2-mutated patients was unclear. Thirdly, since our study only extracted data at a trial level rather than an individual level, other variables, including BRCA1 mutation location, age, and different races, could possibly affect the response to the PARPi. Due to the lack of the necessary resources, we were only able to




FIGURE 2 | Forest plots of pooled analyses for PARP inhibitors vs. controls on progression-free survival stratified by BRCA1 mutation or BRCA2 mutation subgroups. Squares represent study-specific HRs. Horizontal lines indicate the 95% CIs. Diamonds indicate the meta-analytic pooled HRs, calculated separately in BRCA1 mutated (red) and BRCA2 mutated (blue) patients, with their corresponding 95% CIs. The p value for heterogeneity was calculated by the interaction test.




FIGURE 3 | Analyses of BRCA-specific pooled PFS hazard ratios by subgroups. Squares represent subgroup-specific pooled hazard ratios (red for BRCA1 mutation carriers, blue for BRCA2 mutation carriers). Horizontal lines indicate the 95% CIs. The p value for heterogeneity is from the interaction test.

utilize the published data. Therefore, a future individual patient data meta-analysis would be warranted. In addition, the limited number of pancreatic cancer and prostate cancer might not increase the significance of this study. Although, we performed a subgroup analysis to exclude the two small sample studies and obtained still outcome. For this reason, future studies were urgently needed to verify the similar efficacy between the two groups among other malignancies. Fourthly, we could not determine overall survival (OS) because of the lack of OS data in the RCTs, which may have provided a more convincing result. From there, an updated meta-analysis would be needed in the future.

CONCLUSION

Currently, there is no evidence that BRCA2-mutated patients would benefit more from PARPi than BRCA1-mutated patients. BRCA1-mutated patients and BRCA2-mutated patients could benefit from PARPi in clinical practice, and the efficacy is comparable. More clinical trials are currently warranted to verify this trend in various cancer such as pancreatic adenocarcinoma and prostate cancer.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**. Further inquiries can be directed to the corresponding authors.

AUTHOR CONTRIBUTIONS

LD and QG were in charge of funding acquisition, validation, and supervision. SL, LT, and HD performed data curation,

conceptualization and methodology of the study. SL, LT, HD, and XG performed data validation and formal analysis. SL, YGZ, HX, and YYZ completed the project administration, visualization and writing - original draft. SL, LT, and LD finished the manuscript writing - review & editing. All authors contributed to the article and approved the submitted version.

FUNDING

This work was supported by the Joint Project Foundation of Chongqing Health Commission and Chongqing Scientific and Technological Commission (grant number 2021MSXM103).

ACKNOWLEDGMENT

We want to show our gratitude and respect to Taylor Swift, Angelina Jolie, Hugh Jackman, Wang Nan, and other celebrities worldwide who share their own stories about the fight against cancer with positive energy fearlessly and who gives generous donations to cancer research. What they do could have a positive impact on the advocacy of cancer prevention and emphasizes that conquering cancer is never a lonely fight, but is a war for every human being.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fonc.2021.718871/full#supplementary-material>

REFERENCES

- Moynahan ME, Chiu JW, Koller BH, Jasin M. Brca1 Controls Homology-Directed DNA Repair. *Mol Cell* (1999) 4:511–8. doi: 10.1016/S1097-2765(00)80202-6
- Moynahan ME, Pierce AJ, Jasin M. BRCA2 Is Required for Homology-Directed Repair of Chromosomal Breaks. *Mol Cell* (2001) 7:263–72. doi: 10.1016/S1097-2765(01)00174-5
- Tutt A, Ashworth A. The Relationship Between the Roles of BRCA Genes in DNA Repair and Cancer Predisposition. *Trends Mol Med* (2002) 8:571–6. doi: 10.1016/S1471-4914(02)02434-6
- Venkitaraman AR. Cancer Susceptibility and the Functions of BRCA1 and BRCA2. *Cell* (2002) 108:171–82. doi: 10.1016/S0092-8674(02)00615-3
- Oh M, McBride A, Yun S, Bhattacharjee S, Slack M, Martin JR, et al. BRCA1 and BRCA2 Gene Mutations and Colorectal Cancer Risk: Systematic Review and Meta-Analysis. *J Natl Cancer Inst* (2018) 110:1178–89. doi: 10.1093/jnci/djy148
- King M-C, Marks JH, Mandell JB. Breast and Ovarian Cancer Risks Due to Inherited Mutations in BRCA1 and BRCA2. *Science* (2003) 302:643–6. doi: 10.1126/science.1088759
- Thompson D, Easton DF. Cancer Incidence in BRCA1 Mutation Carriers. *J Natl Cancer Inst* (2002) 94:1358–65. doi: 10.1093/jnci/94.18.1358
- Momozawa Y, Iwasaki Y, Hirata M, Liu X, Kamatani Y, Takahashi A, et al. Germline Pathogenic Variants in 7636 Japanese Patients With Prostate Cancer and 12 366 Controls. *J Natl Cancer Inst* (2020) 112:369–76. doi: 10.1093/jnci/djz124
- Liu G, Yang D, Sun Y, Shmulevich I, Xue F, Sood AK, et al. Differing Clinical Impact of BRCA1 and BRCA2 Mutations in Serous Ovarian Cancer. *Pharmacogenomics* (2012) 13:1523–35. doi: 10.2217/pgs.12.137
- Hyman DM, Zhou Q, Iasonos A, Grisham RN, Arnold AG, Phillips MF, et al. Improved Survival for BRCA2-Associated Serous Ovarian Cancer Compared With Both BRCA-Negative and BRCA1-Associated Serous Ovarian Cancer. *Cancer* (2012) 118:3703–9. doi: 10.1002/cncr.26655
- Yang D, Khan S, Sun Y, Hess K, Shmulevich I, Sood AK, et al. Association of BRCA1 and BRCA2 Mutations With Survival, Chemotherapy Sensitivity, and Gene Mutator Phenotype in Patients With Ovarian Cancer. *JAMA* (2011) 306:1557–65. doi: 10.1001/jama.2011.1456
- Bolton KL, Chenevix-Trench G, Goh C, Sadetzki S, Ramus SJ, Karlan BY, et al. Association Between BRCA1 and BRCA2 Mutations and Survival in Women With Invasive Epithelial Ovarian Cancer. *JAMA* (2012) 307:382–90. doi: 10.1001/jama.2012.20
- Pilié PG, Tang C, Mills GB, Yap TA. State-Of-the-Art Strategies for Targeting the DNA Damage Response in Cancer. *Nat Rev Clin Oncol* 16 (2019) 16 (2):81–104. doi: 10.1038/s41571-018-0114-z
- Moore K, Colombo N, Scambia G, Kim B-G, Oaknin A, Friedlander M, et al. Maintenance Olaparib in Patients With Newly Diagnosed Advanced Ovarian Cancer. *N Engl J Med* (2018) 379:2495–505. doi: 10.1056/NEJMoa1810858
- Litton JK, Rugo HS, Ettl J, Hurvitz SA, Gonçalves A, Lee K-H, et al. Talazoparib in Patients With Advanced Breast Cancer and a Germline BRCA Mutation. *N Engl J Med* (2018) 379:753–63. doi: 10.1056/NEJMoa1802905

16. Coleman RL, Fleming GF, Brady MF, Swisher EM, Steffensen KD, Friedlander M, et al. Veliparib With First-Line Chemotherapy and as Maintenance Therapy in Ovarian Cancer. *N Engl J Med* (2019) 381:2403–15. doi: 10.1056/NEJMoa1909707
17. Han HS, Diéras V, Robson M, Palácová M, Marcom PK, Jager A, et al. Veliparib With Temozolomide or Carboplatin/Paclitaxel Versus Placebo With Carboplatin/Paclitaxel in Patients With BRCA1/2 Locally Recurrent/Metastatic Breast Cancer: Randomized Phase II Study. *Ann Oncol* (2018) 29:154–61. doi: 10.1093/annonc/mdx505
18. Mirza MR, Monk BJ, Herrstedt J, Oza AM, Mahner S, Redondo A, et al. Niraparib Maintenance Therapy in Platinum-Sensitive, Recurrent Ovarian Cancer. *N Engl J Med* (2016) 375:2154–64. doi: 10.1056/NEJMoa1611310
19. Pujade-Lauraine E, Ledermann JA, Selle F, Gebbski V, Penson RT, Oza AM, et al. Olaparib Tablets as Maintenance Therapy in Patients With Platinum-Sensitive, Relapsed Ovarian Cancer and a BRCA1/2 Mutation (SOLO2/ENGOT-Ov21): A Double-Blind, Randomised, Placebo-Controlled, Phase 3 Trial. *Lancet Oncol* (2017) 18:1274–84. doi: 10.1016/S1470-2045(17)30469-2
20. Coleman RL, Oza AM, Lorusso D, Aghajanian C, Oaknin A, Dean A, et al. Rucaparib Maintenance Treatment for Recurrent Ovarian Carcinoma After Response to Platinum Therapy (ARIEL3): A Randomised, Double-Blind, Placebo-Controlled, Phase 3 Trial. *Lancet* (2017) 390:1949–61. doi: 10.1016/S0140-6736(17)32440-6
21. Robson M, Im S-A, Senkus E, Xu B, Domchek SM, Masuda N, et al. Olaparib for Metastatic Breast Cancer in Patients With a Germline BRCA Mutation. *N Engl J Med* (2017) 377:523–33. doi: 10.1056/NEJMoa1706450
22. Golan T, Hammel P, Reni M, Van Cutsem E, Macarulla T, Hall MJ, et al. Maintenance Olaparib for Germline -Mutated Metastatic Pancreatic Cancer. *N Engl J Med* (2019) 381:317–27. doi: 10.1056/NEJMoa1903387
23. Ray-Coquard I, Pautier P, Pignata S, Pérol D, González-Martín A, Berger R, et al. Olaparib Plus Bevacizumab as First-Line Maintenance in Ovarian Cancer. *N Engl J Med* (2019) 381:2416–28. doi: 10.1056/NEJMoa1911361
24. de Bono J, Mateo J, Fizazi K, Saad F, Shore N, Sandhu S, et al. Olaparib for Metastatic Castration-Resistant Prostate Cancer. *N Engl J Med* (2020) 382:2091–102. doi: 10.1056/NEJMoa1911440
25. Lord CJ, Ashworth A. PARP Inhibitors: Synthetic Lethality in the Clinic. *Science* (2017) 355:1152–8. doi: 10.1126/science.aam7344
26. Zhong L, Tran AT, Tomasino T, Nugent E, Smith JA. Cost-Effectiveness of Niraparib and Olaparib as Maintenance Therapy for Patients With Platinum-Sensitive Recurrent Ovarian Cancer. *J Manag Care Spec Pharm* (2018) 24:1219–28. doi: 10.18553/jmcp.2018.24.12.1219
27. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JPA, et al. The PRISMA Statement for Reporting Systematic Reviews and Meta-Analyses of Studies That Evaluate Health Care Interventions: Explanation and Elaboration. *PloS Med* (2009) 6:e1000100. doi: 10.1371/journal.pmed.1000100
28. Altman DG, Bland JM. Interaction Revisited: The Difference Between Two Estimates. *BMJ* (2003) 326:219. doi: 10.1136/bmj.326.7382.219
29. Fisher DJ, Carpenter JR, Morris TP, Freeman SC, Tierney JF. Meta-Analytical Methods to Identify Who Benefits Most From Treatments: Daft, Deluded, or Deft Approach? *BMJ* (2017) 356:j573. doi: 10.1136/bmj.j573
30. Begg CB, Mazumdar M. Operating Characteristics of a Rank Correlation Test for Publication Bias. *Biometrics* (1994) 50:1088–101. doi: 10.2307/2533446
31. Egger M, Davey Smith G, Schneider M, Minder C. Bias in Meta-Analysis Detected by a Simple, Graphical Test. *BMJ (Clinical Res Ed)* (1997) 315:629–34. doi: 10.1136/bmj.315.7109.629
32. Sterne JA, Sutton AJ, Ioannidis JP, Terrin N, Jones DR, Lau J, et al. Recommendations for Examining and Interpreting Funnel Plot Asymmetry in Meta-Analyses of Randomised Controlled Trials. *BMJ (Clinical Res Ed)* (2011) 343:d4002. doi: 10.1136/bmj.d4002
33. Lee CK, Friedlander ML, Tjokrowidjaja A, Ledermann JA, Coleman RL, Mirza MR, et al. Molecular and Clinical Predictors of Improvement in Progression-Free Survival With Maintenance PARP Inhibitor Therapy in Women With Platinum-Sensitive, Recurrent Ovarian Cancer: A Meta-Analysis. *Cancer* (2011) 127(14):2432–41. doi: 10.1002/cncr.33517
34. Ledermann J, Harter P, Gourley C, Friedlander M, Vergote I, Rustin G, et al. Olaparib Maintenance Therapy in Platinum-Sensitive Relapsed Ovarian Cancer. *N Engl J Med* (2012) 366:1382–92. doi: 10.1056/NEJMoa1105535
35. Ford L, Wolford JE, Brown SM, Randall LM. A Profile on the FoundationFocus CDxBRCA Tests. *Expert Rev Mol Diagn* (2020) 20:285–92. doi: 10.1080/14737159.2020.1701438
36. Gunderson CC, Moore KN. BRCAAnalysis CDx as a Companion Diagnostic Tool for Lynparza. *Expert Rev Mol Diagn* (2015) 15:1111–6. doi: 10.1586/14737159.2015.1078238
37. Vencken PMLH, Kriege M, Hoogwerf D, Beuglink S, van der Burg MEL, Hooning MJ, et al. Chemosensitivity and Outcome of BRCA1- and BRCA2-Associated Ovarian Cancer Patients After First-Line Chemotherapy Compared With Sporadic Ovarian Cancer Patients. *Ann Oncol* (2011) 22:1346–52. doi: 10.1093/annonc/mdq628
38. Reitsma W, de Bock GH, Oosterwijk JC, ten Hoor KA, Hollema H, Mourits MJE. Clinicopathologic Characteristics and Survival in BRCA1- and BRCA2-Related Adnexal Cancer: Are They Different? *Int J Gynecol Cancer* (2012) 22:579–85. doi: 10.1097/IGC.0b013e31823d1b5c
39. Yarden RI, Pardo-Reoyo S, Sgagias M, Cowan KH, Brody LC. BRCA1 Regulates the G2/M Checkpoint by Activating Chk1 Kinase Upon DNA Damage. *Nat Genet* (2002) 30:285–9. doi: 10.1038/ng837
40. Joukov V, Groen AC, Prokhorova T, Gerson R, White E, Rodriguez A, et al. The BRCA1/BARD1 Heterodimer Modulates Ran-Dependent Mitotic Spindle Assembly. *Cell* (2006) 127:539–52. doi: 10.1016/j.cell.2006.08.053
41. Farmer H, McCabe N, Lord CJ, Tutt ANJ, Johnson DA, Richardson TB, et al. Targeting the DNA Repair Defect in BRCA Mutant Cells as a Therapeutic Strategy. *Nature* (2005) 434:917–21. doi: 10.1038/nature03445
42. Rijnsburger AJ, Obdeijn I-M, Kaas R, Tilanus-Linthorst MMA, Boetes C, Loo CE, et al. BRCA1-Associated Breast Cancers Present Differently From BRCA2-Associated and Familial Cases: Long-Term Follow-Up of the Dutch MRISC Screening Study. *J Clin Oncol* (2010) 28:5265–73. doi: 10.1200/JCO.2009.27.2294
43. Ha SM, Chae EY, Cha JH, Kim HH, Shin HJ, Choi WJ. Association of BRCA Mutation Types, Imaging Features, and Pathologic Findings in Patients With Breast Cancer With BRCA1 and BRCA2 Mutations. *AJR Am J Roentgenol* (2017) 209:920–8. doi: 10.2214/AJR.16.16957
44. Drost R, Bouwman P, Rottenberg S, Boon U, Schut E, Klarenbeek S, et al. BRCA1 RING Function Is Essential for Tumor Suppression But Dispensable for Therapy Resistance. *Cancer Cell* (2011) 20:797–809. doi: 10.1016/j.ccr.2011.11.014
45. Huo Y, Selenica P, Mahdi AH, Pareja F, Kyker-Snowman K, Chen Y, et al. Genetic Interactions Among Brca1, Brca2, Palb2, and Trp53 in Mammary Tumor Development. *NPJ Breast Cancer* (2021) 7:45. doi: 10.1038/s41523-021-00253-5
46. Sy SMH, Huen MSY, Chen J. PALB2 Is an Integral Component of the BRCA Complex Required for Homologous Recombination Repair. *Proc Natl Acad Sci USA* (2009) 106:7155–60. doi: 10.1073/pnas.0811159106
47. Roy R, Chun J, Powell SN. BRCA1 and BRCA2: Different Roles in a Common Pathway of Genome Protection. *Nat Rev Cancer* (2011) 12:68–78. doi: 10.1038/nrc3181
48. Lin Q, Liu W, Xu S, Shang H, Li J, Guo Y, et al. PARP Inhibitors as Maintenance Therapy in Newly Diagnosed Advanced Ovarian Cancer: A Meta-Analysis. *BJOG* (2021) 128:485–93. doi: 10.1111/1471-0528.16411
49. Macciò A, Madeddu C. The Mechanism of Cancer Cell Death by PARP Inhibitors Goes Beyond DNA Damage Alone. *Int J Cancer* (2019) 145:2594–6. doi: 10.1002/ijc.32392

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Li, Tao, Dai, Gong, Zhuo, Xiang, Zhao, Gao and Deng. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.