



Case Report: Detection of Symptomatic Treatment-Related Changes in a Patient With Anaplastic Oligodendroglioma Using FET PET

OPEN ACCESS

Edited by:

Alessandro Stecco,
University of Eastern Piedmont, Italy

Reviewed by:

Francesco Cicone,
University of Catanzaro, Italy
Marco Riva,
University of Milan, Italy
Jens Gempt,
Technische Universität München,
Germany

*Correspondence:

Norbert Galldiks
n.galldiks@fz-juelich.de

Specialty section:

This article was submitted to
Cancer Imaging and
Image-directed Interventions,
a section of the journal
Frontiers in Oncology

Received: 02 July 2021

Accepted: 02 November 2021

Published: 16 November 2021

Citation:

Bauer EK, Werner J-M, Fink GR,
Langen K-J and Galldiks N (2021)
Case Report: Detection of
Symptomatic Treatment-Related
Changes in a Patient With Anaplastic
Oligodendroglioma Using FET PET.
Front. Oncol. 11:735388.
doi: 10.3389/fonc.2021.735388

Elena Katharina Bauer¹, Jan-Michael Werner¹, Gereon R. Fink^{1,2}, Karl-Josef Langen^{2,3,4} and Norbert Galldiks^{1,2,4*}

¹ Department of Neurology, Faculty of Medicine and University Hospital Cologne, University of Cologne, Cologne, Germany, ² Institute of Neuroscience and Medicine (INM-3, -4), Research Center Juelich, Juelich, Germany, ³ Department of Nuclear Medicine, University Hospital Aachen, Aachen, Germany, ⁴ Center of Integrated Oncology (CIO), Universities of Aachen, Bonn, Cologne, and Duesseldorf, Cologne, Germany

Following local and systemic treatment of gliomas, the differentiation between glioma relapse and treatment-related changes such as pseudoprogression or radiation necrosis using conventional MRI is limited. To overcome this limitation, various amino acid PET tracers such as O-[2-(¹⁸F)-fluoroethyl]-L-tyrosine (FET) are increasingly used and provide valuable additional clinical information. We here report neuroimaging findings in a clinically symptomatic 53-year-old woman with a recurrent anaplastic oligodendroglioma with MRI findings highly suspicious for tumor progression. In contrast, FET PET imaging suggested treatment-related changes considerably earlier than the regression of contrast enhancement on MRI. In patients with oligodendroglioma, the phenomenon of symptomatic treatment-related changes is not well described, making these imaging findings unique and important for clinical decision-making.

Keywords: amino acid PET, pseudoprogression, chemoradiation, oligodendroglioma, diagnostic challenge

INTRODUCTION


Treatment-related changes on conventional MRI are predominantly characterized by both an increase of contrast enhancement in the T1-weighted image and a hyperintensity in the fluid-attenuated inversion recovery (FLAIR) sequence, which either remain stable or even disappear over time without any change of anticancer treatment. Not infrequently, patients experience no clinical

deterioration. Misinterpretation of these imaging findings may result in an unnecessary discontinuation of an effective treatment with a potentially negative impact on the patient's survival. Due to the low specificity of conventional MRI, the differentiation of treatment-related changes from actual tumor progression in glioma patients following radiotherapy or chemoradiation using conventional MRI is challenging (1–4).

The additional use of amino acid PET using O-[2-(^{18}F)-fluoroethyl]-L-tyrosine (FET) in the care of glioma patients is gaining more attention. In recent years, a high diagnostic accuracy for the differentiation between tumor progression and treatment-related changes using FET PET has repeatedly been shown predominantly in patients with astrocytic gliomas (1, 2, 5, 6). In these studies, either static parameters or the

combination of static and dynamic parameters derived from FET PET seem to be of considerable clinical value for the differentiation of tumor progression from treatment-related changes (range of sensitivity, 91–99%; range of specificity, 94–100%) (1, 5, 6). Despite its clinical relevance, only a few studies described treatment-related changes in patients with oligodendrogliomas.

Oligodendroglial gliomas are characterized by the presence of an isocitrate dehydrogenase (IDH) gene mutation and a 1p/19q codeletion and are histologically corresponding to the grade II or III of the in 2016 revised World Health Organisation (WHO) classification of Tumors of the Central Nervous System (7). This rare tumor entity accounts for 5.3% of all tumors of the central nervous system.



We here report the phenomenon of symptomatic treatment-related changes in a patient with an anaplastic oligodendroglioma.

CASE PRESENTATION

A 53-year-old female patient with a recurrent anaplastic oligodendroglioma (IDH mutant, 1p/19q co-deleted) was diagnosed with the second local tumor recurrence 15 years after initial diagnosis. The first-line treatment included tumor resection and radiotherapy. Eleven years later, a first local tumor recurrence was treated with re-resection followed by adjuvant temozolomide chemotherapy. Four years later, the tumor recurred locally again. Following complete resection and re-irradiation, she underwent adjuvant nitrosourea-based chemotherapy with procarbazine and lomustine. After two cycles, the patient experienced a clinical deterioration with an expressive aphasia and a right-sided hemiparesis. According to current response assessment criteria (3), the findings of the first and second follow-up MRI were consistent with “Progressive Disease”, whereas the corresponding FET PET scans remained unchanged compared to baseline (see **Figure 1**). Without changing the treatment regimen, symptomatic therapy with dexamethasone prompted a rapid clinical improvement. During follow-up, dexamethasone could subsequently be tapered from 8 mg to 0.5 mg. The diagnosis of treatment-related changes was based on the MRI findings performed 26 weeks after baseline imaging, which showed especially an almost complete decrease of contrast enhancement, accompanied by a further improvement of the clinical condition.

DISCUSSION

Following radiotherapy with or without alkylating chemotherapy, treatment-related changes such as pseudoprogression or radiation necrosis represent a major diagnostic challenge in patients with astrocytic glioma. In this group of brain tumors, the steadily increasing number of studies highlights the value of amino acid PET for the differentiation of treatment-related changes from actual tumor progression with a high diagnostic accuracy (1, 2, 4, 9).

On the other hand, oligodendrogliomas represent a small glioma subgroup and constitute only approximately 5% of primary intracranial tumors. Importantly, it has been demonstrated that in oligodendrogliomas the rate of treatment-related changes is significantly lower than in patients with gliomas without 1p/19q codeletion (i.e., astrocytoma, glioblastoma) (10, 11). For example, Lin and co-workers reported a pseudo-progression rate of 31% for tumors without codeletions, and only 3% for tumors with codeletions (11). Moreover, amino acid uptake in 1p/19q-codeleted oligodendroglial tumors is generally significantly

increased, even higher when compared to IDH mutant astrocytic tumors (12, 13), and similar to glioblastoma (13). Thus, a considerably increased FET uptake can be expected in the case of oligodendroglioma relapse. Consequently, the absence of this uptake pattern in our case further supports the diagnosis of treatment-related changes.

Another important aspect of the present case is the extensive increase of the FLAIR signal hyperintensity, suggesting non-enhancing tumor progression. In contrast, FET PET showed in spatial correspondence no pathological increased tracer uptake. Thus, FET PET seems also to be of value to rule out this particular phenomenon. In summary, this case highlights the value of FET PET for the diagnosis of treatment-related changes in a glioma subtype characterized by a naturally increased amino acid uptake and a low rate of treatment-related changes.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/supplementary material. Further inquiries can be directed to the corresponding author.

ETHICS STATEMENT

All procedures performed in the studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards. The patients/participants provided their written informed consent to participate in this study. Written informed consent was obtained from the individual(s) for the publication of any potentially identifiable images or data included in this article.

AUTHOR CONTRIBUTIONS

Data acquisition, EB, NG, and J-MW. Writing of manuscript drafts, EB, NG, and J-MW. All authors contributed to the article and approved the submitted version.

FUNDING

Supported by the Deutsche Forschungsgemeinschaft (DFG), project number 428090865 (EB and NG), and the Cologne Clinician Scientist Program (CCSP)/Faculty of Medicine/University of Cologne, funded by the DFG, FI 773/15-1 (J-MW).

REFERENCES

- Galldiks N, Dunkl V, Stoffels G, Hutterer M, Rapp M, Sabel M, et al. Diagnosis of Pseudoprogression in Patients With Glioblastoma Using O-(2-[18F]

Fluoroethyl)-L-Tyrosine PET. *Eur J Nucl Med Mol Imaging* (2015) 42(5):685–95. doi: 10.1007/s00259-014-2959-4

- Langen KJ, Galldiks N, Hattungen E, Shah NJ. Advances in Neuro-Oncology Imaging. *Nat Rev Neurol* (2017) 13(5):279–89. doi: 10.1038/nrneuro.2017.44

3. Wen PY, Macdonald DR, Reardon DA, Cloughesy TF, Sorensen AG, Galanis E, et al. Updated Response Assessment Criteria for High-Grade Gliomas: Response Assessment in Neuro-Oncology Working Group. *J Clin Oncol* (2010) 28(11):1963–72. doi: 10.1200/JCO.2009.26.3541
4. Albert NL, Weller M, Suchorska B, Galldiks N, Soffiotti R, Kim MM, et al. Response Assessment in Neuro-Oncology Working Group and European Association for Neuro-Oncology Recommendations for the Clinical Use of PET Imaging in Gliomas. *Neuro Oncol* (2016) 18(9):1199–208. doi: 10.1093/neuonc/now058
5. Bashir A, Mathilde Jacobsen S, Molby Henriksen O, Broholm H, Urup T, Grunnet K, et al. Recurrent Glioblastoma Versus Late Posttreatment Changes: Diagnostic Accuracy of O-(2-[18F]Fluoroethyl)-L-Tyrosine Positron Emission Tomography (18F-FET PET). *Neuro Oncol* (2019) 21(12):1595–606. doi: 10.1093/neuonc/noz166
6. Galldiks N, Stoffels G, Filss C, Rapp M, Blau T, Tscherpel C, et al. The Use of Dynamic O-(2-[18F]-Fluoroethyl)-L-Tyrosine PET in the Diagnosis of Patients With Progressive and Recurrent Glioma. *Neuro Oncol* (2015) 17(9):1293–300. doi: 10.1093/neuonc/nov088
7. Louis DN, Perry A, Reifenberger G, von Deimling A, Figarella-Branger D, Cavenee WK, et al. The 2016 World Health Organization Classification of Tumors of the Central Nervous System: A Summary. *Acta Neuropathol* (2016) 131(6):803–20. doi: 10.1007/s00401-016-1545-1
8. Law I, Albert NL, Arbizu J, Boellaard R, Drzezga A, Galldiks N, et al. Joint EANM/EANO/RANO Practice Guidelines/SNMMI Procedure Standards for Imaging of Gliomas Using PET With Radiolabelled Amino Acids and [(18)F]FDG: Version 1.0. *Eur J Nucl Med Mol Imaging* (2019) 46(3):540–57. doi: 10.1007/s00259-018-4207-9
9. Werner JM, Stoffels G, Lichtenstein T, Borggrefe J, Lohmann P, Ceccon G, et al. Differentiation of Treatment-Related Changes From Tumour Progression: A Direct Comparison Between Dynamic FET PET and ADC Values Obtained From DWI MRI. *Eur J Nucl Med Mol Imaging* (2019) 46(9):1889–901. doi: 10.1007/s00259-019-04384-7
10. Lin AL, White M, Miller-Thomas MM, Fulton RS, Tsien CI, Rich KM, et al. Molecular and Histologic Characteristics of Pseudoprogression in Diffuse Gliomas. *J Neurooncol* (2016) 130(3):529–33. doi: 10.1007/s11060-016-2247-1
11. Lin AL, Liu J, Evans J, Leuthardt EC, Rich KM, Dacey RG, et al. Codeletions at 1p and 19q Predict a Lower Risk of Pseudoprogression in Oligodendrogliomas and Mixed Oligoastrocytomas. *Neuro Oncol* (2014) 16(1):123–30. doi: 10.1093/neuonc/not142
12. Takei H, Shinoda J, Ikuta S, Maruyama T, Muragaki Y, Kawasaki T, et al. Usefulness of Positron Emission Tomography for Differentiating Gliomas According to the 2016 World Health Organization Classification of Tumors of the Central Nervous System. *J Neurosurg* (2019) 1–10. doi: 10.3171/2019.5.JNS19780
13. Kim D, Chun JH, Kim SH, Moon JH, Kang SG, Chang JH, et al. Re-Evaluation of the Diagnostic Performance of (11)C-Methionine PET/CT According to the 2016 WHO Classification of Cerebral Gliomas. *Eur J Nucl Med Mol Imaging* (2019) 46(8):1678–84. doi: 10.1007/s00259-019-04337-0

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Bauer, Werner, Fink, Langen and Galldiks. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.