



# Increased Expression of microRNA-141-3p Improves Necrotizing Enterocolitis of Neonates Through Targeting MNX1

Hui Chen\*, Lichun Zeng, Wei Zheng, Xiaoli Li and Baixing Lin

Department of Neonatology, Shenzhen Hospital of Southern Medical University, Shenzhen, China

**Objective:** MicroRNA-141-3p (miR-141-3p) has been investigated in various kinds of cancers. This research delves into the functions and regulatory mechanisms of miR-141-3p in necrotizing enterocolitis (NEC) of neonates.

**Methods:** NEC tissues were obtained from neonatal mice, and subsequently, expression of miR-141-3p and motor neuron and pancreas homeobox 1 (MNX1) was assayed via RT-qPCR. Moreover, the intestinal histopathological changes and histiocytic apoptosis were observed via hematoxylin and eosin (H&E) and TUNEL staining. The correlative inflammatory factors and oxidative stress markers were evaluated to uncover the influence of miR-141-3p in NEC tissue damage. Further, the relation between MNX1 and miR-141-3p was predicated, and the functions of MNX1 in inflammatory response and cell growth of IEC-6 cells were investigated.

**Results:** Downregulated miR-141-3p and upregulated MNX1 were discovered in NEC tissues. Moreover, miR-141-3p clearly alleviated inflammation response and oxidative stress damage in NEC, which was achieved through regulating inflammatory cytokines (IL-1 $\beta$ , IL-6, and TNF- $\alpha$ ) and oxidative stress markers (MPO, MDA, and SOD) expression. MNX1 was forecasted as a target gene of miR-141-3p; meanwhile, MNX1 overexpression overturned the influence of miR-141-3p in the inflammatory response and cell growth process of IEC-6 cells.

**Conclusion:** These explorations reveal that increased expression of miR-141-3p could improve the damage to intestinal tissues in NEC through targeting MNX1. The research might exhibit a neoteric therapeutic strategy for NEC.

**Keywords:** necrotizing enterocolitis, microRNA-141-3p, motor neuron and pancreas homeobox 1, cell viability, apoptosis

## OPEN ACCESS

### Edited by:

Yogen Singh,  
Cambridge University Hospitals NHS  
Foundation Trust, United Kingdom

### Reviewed by:

Vikranth Babu Anna Venugopalan,  
Sandwell & West Birmingham  
Hospitals NHS Trust, United Kingdom  
Ömer Erdeve,  
Ankara University, Turkey

### \*Correspondence:

Hui Chen  
chenhui246000@163.com

### Specialty section:

This article was submitted to  
Neonatology,  
a section of the journal  
Frontiers in Pediatrics

Received: 24 April 2020

Accepted: 08 June 2020

Published: 28 July 2020

### Citation:

Chen H, Zeng L, Zheng W, Li X and  
Lin B (2020) Increased Expression of  
microRNA-141-3p Improves  
Necrotizing Enterocolitis of Neonates  
Through Targeting MNX1.  
Front. Pediatr. 8:385.  
doi: 10.3389/fped.2020.00385

## INTRODUCTION

Necrotizing enterocolitis (NEC) is an acquired illness, which is the most familiar neonatal gastrointestinal emergency, and it is also a dominating reason for intestinal perforation and systemic inflammatory response syndrome (SIRS) in neonates (1, 2). NEC affects digestive system function, and severe cases can cause systemic symptoms, such as shock, acidosis, and jaundice (3, 4). The representative symptoms of NEC in the clinic incorporate abdominal distension, vomiting,

diarrhea, and bloody stools (5). Adequate parenteral nutrition; maintaining the balance of water, electrolyte, acid, and base; and nutritional support are the treatment policy of NEC (6). Although there are advances in neonatal nursing, NEC is still a leading cause of morbidity and mortality in premature infants (7). Thus, to clarify the pathogenesis of NEC and explore its possible diagnostic and therapeutic targets will be beneficial to the clinical therapy of this disease.

MicroRNA (miRNA) is one kind of endogenous RNA molecule with a length of ~22 nucleotides, which joins in post-transcriptional gene expression in plants and animals (8). Recently, a correlative report disclosed the momentous roles of miRNAs in NEC (9). Among these, the importance of miR-21 has been testified to regulate the inflammatory responses in NEC (9). Additionally, the dysregulation of miR-431 was discovered in intestinal tissues of infants with NEC, which might participate in reinforcing the inflammatory response in NEC tissues and be conducive to pro-inflammatory pathophysiology (10). MiR-141-3p is an extensively researched miRNA, which has been investigated in various cancers (11, 12). Shen et al. reveals that miR-141-3p could allay chronic inflammatory pain through suppressing HMGB1 *in vitro* and *in vivo* (13). Nevertheless, relevant research about the influences of miR-141-3p in NEC remains unreported.

Motor neuron and pancreas homeobox 1 (MNX1) is also named Hb9 or Hlxb9, which can encode transcription factor (14). MNX1 was ascertained to be a newfangled reason for permanent neonatal diabetes in a kindred study (15). Moreover, one research illustrates that upregulated MNX1 has been discovered in prostate cancer cells, which might be an innovative biomarker of this cancer (16). Other research also uncovers that abnormally expressed MNX1 is linked to infantile acute myelocytic leukemia (AML) (17). But the effect and the potential regulated mechanisms of MNX1 are still indistinct in NEC of neonates.

In our research, we discovered the ectopic expression of miR-141-3p and MNX1 in NEC tissues. In addition, miR-141-3p improved the inflammation response and oxidative stress damage in NEC through targeting MNX1. Outside of these, MNX1 joined in modulating the influences of miR-141-3p in IEC-6 cell vitality and apoptosis. These explorations might offer a new therapeutic thought for NEC.

## METHODS AND METHODS

### NEC Model Construction and Groups

Experimental male C57BL/6 mice (4–7 days) were achieved from Jiesi Experimental Animal Co., Ltd. (Shanghai, China). The mice were divided into four groups: Control, NEC, NEC+miR-141 NC, and NEC+miR-141 mimics. Breast-feeding and enema with normal saline was utilized to establish a Control group; artificial formula feeding combined with hypoxic cold stimulation was used to establish the NEC group; breast-feeding and enema with 0.2 mL miRNA NC was employed to build the NEC+miR-141 NC group; artificial formula feeding combined with hypoxic cold stimulation and enema with 0.2 mL miR-141-3p was used to structure the NEC+miR-141 mimics group. The disposed

detailed procedures were performed according to previous research (18). After anesthetizing the mice with pentobarbital sodium, the intestinal tissues were collected and washed by normal saline. The obtained samples were stored in liquid nitrogen and placed in a  $-80^{\circ}\text{C}$  refrigerator.

### Histopathologic Examination

Two centimeters of the terminal ileum were taken, which were then immediately fixed in 10% neutral formaldehyde as well as embedded in paraffin and prepared in  $3\ \mu\text{m}$  sections. Subsequently, the hematoxylin and eosin (H&E) staining was carried out to disclose the functions of miR-141-3p in NEC. Additionally, the intestinal histopathological score was judged on the basis of a report from (19). The highest score observed in the sample was taken to determine the degree of intestinal damage, and the histological score  $\geq 2$  points was determined as NEC.

### TUNEL Staining Procedure

Cell apoptosis from the NEC tissues was evaluated through utilizing the DeadEnd™ fluorometric TUNEL system kit (Promega, Madison, WI, USA). The detailed procedures refer to the previous research (20, 21).

### Cell Transfection

The experimental vectors of miR-141 mimics/mimics NC and pcDNA-MNX1/pcDNA-NC were established by GenePharma (Shanghai, China). On the basis of the lipofectamine 3000 reagent (Invitrogen), a cell transfection assay was carried out for 48 h.

### Evaluation of Inflammatory Cytokine Levels

The paired ELISA kits of IL-6, IL-1 $\beta$ , and TNF- $\alpha$  (all from Shanghai Biotechnology Co., Ltd., Shanghai, China) were applied to evaluate the concentrations of IL-6, IL-1 $\beta$ , and TNF- $\alpha$  in intestinal tissues and IEC-6 cells after the corresponding vector transfection or LPS treatment. The optical density (OD) values of each well were measured by a microplate reader (Thermo Scientific Corporation, Massachusetts, USA) at 450 nm.

### RT-qPCR

Total RNA from intestinal tissues and IEC-6 cells was isolated with the aid of TRIzol reagent (Invitrogen). The cDNA template was compounded on the PCR-Cycler. The RT-qPCR procedure was implemented via applying ABI7500 quantitative PCR instrument (ABI, Foster, CA, USA). The CT values achieved from the experiment were analyzed through executing the  $2^{-\Delta\Delta\text{Ct}}$  method (22). The primer sequences of IL-6, IL-1 $\beta$ , TNF- $\alpha$ , and GAPDH were synthesized by the Sangon Biotech (Shanghai, China) and are listed in **Table 1**.

### Detection of MPO, MDA, and SOD Levels

Distalileum tissue samples were achieved to prepare the tissue homogenate (10% w/v). Conforming to the description by Hillegass et al. (23), myeloperoxidase (MPO) activity was examined. Additionally, the levels of malondialdehyde (MDA) and superoxide dismutase (SOD) were measured according to the corresponding kit specifications (Jiancheng Bioengineering Institute, Nanjing, China).

**TABLE 1** | The correlative primer sequences for RT-qPCR.

Name	Primer sequences (5'-3')
IL-6-F	GATACCACTCCCAACAGAC
IL-6-R	CTTTTCTCATTCCACGAT
IL-1 $\beta$ -F	GTGGTGTGTGACGTTCCCATTA
IL-1 $\beta$ -R	CCGACAGCACGAGGCTTT
TNF- $\alpha$ -F	TGCAGCAGGACATCAAGTTC
TNF- $\alpha$ -R	TACGCCTCAGCAGTCTCCTT
GAPDH-F	GCTTCGGCAGCACATATACTAAAT
GAPDH -R	CGCTTCACGAATTTGCGTGTGCAT

## Luciferase Reporter Assay

The miR-141-3p fragment or the 3'-UTR of MNX1 was amplified by PCR, and subsequently, the MNX1-WT was structured by using the pmirGLO luciferase vector (Promega, Madison, WI, USA). MNX1-MUT was established through applying a site-directed mutagenesis kit (Stratagene, La Jolla, CA, USA). These above-involved vectors were co-transfected into IEC-6 cells with miR-141-3p mimics and mimics NC. The luciferase activity was evaluated on the basis of the dual luciferase reporter assay system (Promega) after transfection for 48 h.

## Western Blot Assay

After mimics NC and miR-141 mimics transfection, the proteins from IEC-6 cells were extracted in line with the protein extraction kit (Beyotime, Shanghai, China). The protein concentration was assessed by a BCA kit (Beyotime). The experimental steps of SDS-PAGE and PVDF membrane transfer were carried out according to the foregone research. After this, the membranes were blocked by BSA, and the primary antibody against MNX1 and the corresponding second antibody were utilized to co-culture with the PVDF membranes. The protein bands emerged via using the enhanced chemiluminescence (ECL) reagent (Amersham Biosciences Corp., Piscataway, NJ) and, meanwhile, quantified through Quantity One software (Bio-Rad, Hercules, CA, USA).

## Detection of Cell Viability

The intestinal epithelial cell line, IEC-6, was obtained from ATCC (Rockville, MD, USA), which was cultured in DMEM mixed with 10% FBS. After LPS irritation and the corresponding vector transfection, the disposed IEC-6 cells were collected and inoculated into 96-well plates. After 24 h culture, cells were reacted with 20  $\mu$ L MTT solution (5 mg/mL, Sigma, St. Louis, MO, USA) for an extra 4 h. When terminating the incubation, 150  $\mu$ L DMSO (4%, Sigma) was used to accelerate the crystal dissolution. After accomplishing these procedures, a microplate reader (BioTek, Winooski, VT, USA) was applied to test the absorbance at 570 nm.

## Determination of Cell Death

After LPS irritation and the corresponding vector transfection, an Annexin V-PI Kit (Invitrogen, Life Technologies, USA) was utilized for exploring the apoptosis ability of IEC-6 cells. The disposed IEC-6 cells were collected and, meanwhile, washed

once with PBS (Invitrogen) as well as resuspended in 1  $\times$  binding buffer. Afterward, 5  $\mu$ L Annexin V-FITC and 5  $\mu$ L PI staining solutions were added into 400  $\mu$ L cell suspension; in the meantime, cells were incubated for 30 min at 4°C under a dark environment. Immediately, the apoptotic proportion was monitored via executing a flow cytometer (BD Biosciences, Heidelberg, Germany) and analyzed by the FlowJo 7.6 software (Tree Star Inc., San Carlos, CA, USA). Furthermore, the level of lactate dehydrogenase (LDH) was examined by the paired kits (Beyotime, Shanghai, China).

## Statistical Analysis

Relying on GraphPad Prism 7.0 software (San Diego, CA, USA), the data were managed and statistical analyses conducted, and these data were presented as mean + SD. The one-way ANOVA accompanied by Tukey's multiple comparisons test was utilized to compare the differences in multi-groups.  $P < 0.05$  manifested that the consequences possessed significant differences.

This study was conducted after obtaining approval of Shenzhen Hospital of Southern Medical University's ethical committee, and all experiments were in accordance with the guide for the care and use of laboratory animals established by the U.S. National Institutes of Health (Bethesda, MD, USA). All the animal experiments were conducted in the Science Building of Shenzhen Hospital of Southern Medical University.



## RESULTS

### Abnormal Expression of miR-141-3p and MNX1 in NEC Tissues

The NEC tissues obtained from eight male C57BL/6 neonatal mice were used in the experiment; thereafter, the RT-qPCR assay was carried out to examine miR-141-3p and MNX1 expression in NEC tissues. We noted that miR-141-3p was significantly down-expressed in NEC tissues in comparison with that in the control group ( $P < 0.001$ , **Figure 1A**). On the contrary, up-expressed MNX1 was observed in NEC tissues ( $P < 0.001$ , **Figure 1B**). These consequences revealed the aberrant expression of miR-141-3p and MNX1 in NEC tissues.

### miR-141-3p Mitigated the Damage to Intestinal Tissues in NEC


To certify the protection of miR-141-3p for NEC, the weight of neonatal mice was monitored in different groups (Control, NEC, NEC+miR-141 NC, and NEC+miR-141 mimics). We discovered that the weight of neonatal mice was markedly decreased in the NEC group compared with that in the Control group ( $P < 0.01$ , **Figure 2A**). After transfection with miR-141-3p mimics, the weight of neonatal mice was clearly recovered when constructing the NEC model for 72 h ( $P < 0.001$ , **Figure 2A**). HE staining results displayed that the structure of the intestinal tissues were normal and no inflammatory cell infiltration in the Control group. However, the observed lesions from the NEC group presented the severely necrotic intestinal tissue with the increased numbers of infiltrating inflammatory cells. Surprisingly, overexpression of miR-141-3p significantly improved the damage of intestinal tissue (**Figure 2B**).



Additionally, the intestinal histopathological score in the NEC group was increased as contrasted with the Control group ( $P < 0.001$ ). Inversely, overexpressed miR-141-3p decreased the NEC score ( $P < 0.001$ , **Figure 2C**). Furthermore, TUNEL staining revealed apoptosis of intestinal epithelial cells. Microscopically, we noted that a large number of cell nuclei were blue in the NEC group, which presented strong positive. But partial cell nuclei were blue in the NEC+miR-141 mimics group, which exhibited as positive (**Figure 2D**). These observations hinted that miR-141-3p distinctly inhibited cell apoptosis. In a word, the above results showcased that miR-141-3p could mitigate the damage to intestinal tissues of NEC.

### miR-141-3p Alleviated Inflammation Response and Oxidative Stress Damage in NEC

In the next experiment, we determined the accumulation of inflammatory cytokines (IL-1 $\beta$ , IL-6, and TNF- $\alpha$ ) in the Control, NEC, NEC+miR-141 NC, and NEC+miR-141 mimics groups. We observed that the concentrations and mRNA expression of IL-1 $\beta$ , IL-6, and TNF- $\alpha$  were significantly increased in the NEC group ( $P < 0.001$ , **Figures 3A–F**). Nevertheless, in the NEC+miR-141 mimics group, the above phenomenon was obviously overturned ( $P < 0.001$ , **Figures 3A–F**). Beyond these, compared with the Control group, the levels of MPO and




MDA were conspicuously augmented, but the level of SOD was clearly declined in the NEC group ( $P < 0.001$ , **Figures 3G–I**). Compared with the NEC+miR-141 NC group, the NEC+miR-141 mimics group prohibited the levels of MPO and MDA and, meanwhile, increased SOD content ( $P < 0.01$ , **Figures 3G–I**). The above-involved consequences evinced that miR-141-3p could allay inflammation response and oxidative stress damage in NEC.

### MNX1 Was a Direct Target of miR-141-3p

The relevance between MNX1 and miR-141-3p was forecasted through applying Starbase software and the dual luciferase

reporter assay. In **Figure 4A**, the supposed binding site between MNX1 and miR-141-3p sequences was exhibited. Additionally, the dual luciferase reporter assay revealed that co-transfection with miR-141-3p mimics and MNX1-WT significantly restrained the luciferase activity ( $P < 0.01$ , **Figure 4B**). The RT-qPCR assay disclosed the up-expressed miR-141-3p in miR-141-3p mimics transfected cells ( $P < 0.01$ , **Figure 4C**). Further, the Western blot assay showcased that the protein level of MNX1 was visibly declined in miR-141-3p mimics transfected cells ( $P < 0.01$ , **Figure 4D**). In short, the results explained that MNX1 was a new-predicted target gene of miR-141-3p and negatively regulated by miR-141-3p.



## MNX1 Eliminated the Functions of miR-141-3p in Inflammatory Response of IEC-6 Cells

To explore whether MNX1 joins in regulating the protective function of miR-141-3p in NEC, the vector of pcDNA-MNX1 was transfected into IEC-6 cells to overexpress MNX1 expression. After LPS irritation, the concentrations and mRNA expression of IL-1 $\beta$ , IL-6, and TNF- $\alpha$  were all significantly upgraded in IEC-6 cells ( $P < 0.01$ , Figures 5A–F). But the above results were reversed by overexpression of miR-141-3p ( $P < 0.01$ , Figures 5A–F). When pcDNA-MNX1 was transfected into IEC-6 cells, we surprisingly discovered that the decreased productions of IL-1 $\beta$ , IL-6, and TNF- $\alpha$  triggered by overexpressed miR-141-3p were overturned in LPS-treated IEC-6 cells ( $P < 0.05$ ,  $P < 0.01$ , Figures 5A–F). These findings revealed that MNX1 could eliminate the protective functions of miR-141-3p in LPS-damaged IEC-6 cells.

## MNX1 Regulated the Functions of miR-141-3p in IEC-6 Cell Growth

Further exploration uncovered the regulatory functions of MNX1 in IEC-6 cell viability and apoptosis. The MTT assay explained that LPS irritation clearly restrained cell viability, promoted apoptosis, and increased LDH activity in IEC-6 cells ( $P < 0.01$ , Figures 6A–C). Nonetheless, overexpressed miR-141-3p expedited cell viability, prohibited apoptosis, and declined LDH activity in LPS-irritated IEC-6 cells ( $P < 0.01$ , Figures 6A–C). More interestingly, MNX1 overexpression crippled the functions


of miR-141-3p in LPS-affected IEC-6 cell viability and apoptosis ( $P < 0.05$ ,  $P < 0.01$ , Figures 6A–C). All aforesaid consequences imparted that MNX1 subdued the functions of miR-141-3p in IEC-6 cell growth.

## DISCUSSION

NEC is a common acute intestinal inflammatory necrotizing ailment in neonates (24). Recently, studies on the pathogenesis of NEC have primarily focused on the intestinal mucosal barrier and the interaction of intestinal innate immunity and the microbial host (25). Its detailed pathogenesis has not been fully defined. In the existing research, the significant roles of miRNAs in NEC have been reported; however, functions of miR-141-3p in this disease are still ambiguous. Here, we found the aberrant expression of miR-141-3p and MNX1 in NEC tissues. Moreover, the protective functions of miR-141-3p in inflammation response and oxidative stress damage in NEC were certified. Additionally, we found that MNX1 was a target gene of miR-141-3p, and overexpressed MNX1 overturned the influence of miR-141-3p in LPS-damaged IEC-6 cells.

Various studies have discovered that miRNA can play important roles in inflammatory response; innate immunity; tumor formation, proliferation, and differentiation; and apoptosis of intestinal epithelial cells (26, 27). Meanwhile, miRNAs are linked to various intestinal physiological and pathological processes through modulating the intestinal immune or inflammatory response (28). In newborns, miR-146a is reported to downregulate IRAK1 expression to prevent





cell proliferation via interaction with NONO protein (44). In the course of inflammation, the capacity of cell viability and apoptosis tend to change naturally (45). LDH activity has been confirmed to be a vital indicator to detect cell apoptosis and cell necrosis (46). Based on this foregoing research, we discovered that overexpressed miR-141-3p clearly accelerates cell viability while it restrained cell apoptosis and LDH activity in LPS-damaged IEC-6 cells. What's more, MNX1 overexpression reversed the functions of miR-141-3p in cell viability and apoptosis. These explorations hinted that MNX1 might regulate the functions of miR-141-3p in LPS-damaged IEC-6 cells.

Taken together, our research reveals that miR-141-3p could improve inflammation response and oxidative stress damage in NEC via targeting MNX1. This conclusion might provide a therapeutic strategy for curing NEC of neonates. Nonetheless, further research is still necessary for expounding the detail mechanisms of miR-141-3p-MNX1 axis in NEC.

## REFERENCES

1. Tayman C, Tonbul A, Kahveci H, Uysal S, Koseoglu B, Tatli MM, et al. C5a, a complement activation product, is a useful marker in predicting the severity of necrotizing enterocolitis. *Tohoku J Exp Med.* (2011) 224:143–50. doi: 10.1620/tjem.224.143

## DATA AVAILABILITY STATEMENT

All datasets generated for this study are included in the article.

## ETHICS STATEMENT

This study was conducted after obtaining approval of Shenzhen Hospital of Southern Medical University's ethical committee and all experiments were in accordance with the guide for the care and use of laboratory animals established by United States National Institutes of Health (Bethesda, MD, USA).

## AUTHOR CONTRIBUTIONS

HC: conception and design and analysis of data. HC, LZ, and WZ: drafting the article. XL and BL: revising the article critically for important intellectual content. All authors read and approved the final manuscript.

2. McBride W, Roy S, Brudnicki A, Stringel G. Correlation of complex ascites with intestinal gangrene and perforation in neonates with necrotizing enterocolitis. *J Pediatr Surg.* (2010) 45:887–9. doi: 10.1016/j.jpedsurg.2010.02.011
3. Blackwood BP, Yuan CY, Wood DR, Nicolas JD, Grothaus JS, Hunter CJ. Probiotic lactobacillus species strengthen intestinal barrier

- function and tight junction integrity in experimental necrotizing enterocolitis. *J Probiotics Health*. (2017) 5:159. doi: 10.4172/2329-8901.100159
4. Ding G. Neonatal infection and neonatal necrotizing enterocolitis. *Chin Pediatr Emerg Med*. (2017) 24:330–3.
  5. Terrin G, Stronati L, Cucchiara S, De Curtis M. Serum markers of necrotizing enterocolitis: a systematic review. *J Pediatr Gastroenterol Nutr*. (2017) 65:e120–32. doi: 10.1097/MPG.0000000000001588
  6. Perks P, Abad-Jorge A. Nutritional management of the infant with necrotizing enterocolitis. *Pract Gastroenterol*. (2008) 32:133–42. doi: 10.1002/ncp.10115
  7. Hunter CJ, Upperman JS, Ford HR, Camerini V. Understanding the susceptibility of the premature infant to necrotizing enterocolitis (NEC). *Pediatr Res*. (2016). 63:117–23. doi: 10.1203/PDR.0b013e31815ed64c
  8. Wang C, Zhao WM, Mou YL. Research progress of microRNA on animal phenotypic regulation. *J Agr Sci Tech*. (2013) 15:108–12.
  9. Requena P, Beaupré J, Goldstein AM, Vanderburg CR, Nanthakumar NN. The role of MicroRNA in the pathogenesis associated with necrotizing enterocolitis. *Gastroenterology*. (2010) 138:S–301. doi: 10.1016/S0016-5085(10)61384-4
  10. Wu YZ, Chan KYY, Leung KT, Lam HS, Tam YH, Lee KH, et al. Dysregulation of miR-431 and target gene FOXA1 in intestinal tissues of infants with necrotizing enterocolitis. *FASEB J*. (2019) 33:5143–52. doi: 10.1096/fj.201801470R
  11. Zheng L, Xu M, Xu J, Wu K, Fang Q, Liang Y, et al. ELF3 promotes epithelial–mesenchymal transition by protecting ZEB1 from miR-141-3p-mediated silencing in hepatocellular carcinoma. *Cell Death Dis*. (2018) 9:387. doi: 10.1038/s41419-018-0399-y
  12. Fang M, Huang W, Wu X, Gao Y, Ou J, Zhang X, et al. MiR-141-3p suppresses tumor growth and metastasis in papillary thyroid cancer via targeting Yin Yang 1. *Anat Rec*. (2018) 302:258–68. doi: 10.1002/ar.23940
  13. Shen WS, Xu XQ, Zhai NN, Zhou ZS, Shao J, Yu YH. Potential mechanisms of microRNA-141-3p to alleviate chronic inflammatory pain by down-regulation of downstream target gene HMGB1: *in vitro* and *in vivo* studies. *Gene Therapy*. (2017) 24:353–60. doi: 10.1038/gt.2017.28
  14. Leotta CG, Federico C, Brundo MV, Tosi S, Saccone S. HLXB9 gene expression, and nuclear location during *in vitro* neuronal differentiation in the SK-N-BE neuroblastoma cell line. *PLoS ONE*. (2014) 9:e105481. doi: 10.1371/journal.pone.0105481
  15. Bonnefond A, Vaillant E, Philippe J, Skrobek B, Vaxillaire M. Transcription factor gene MNX1 is a novel cause of permanent neonatal diabetes in a consanguineous family. *Diabetes Metab*. (2013) 39:276–80. doi: 10.1016/j.diabet.2013.02.007
  16. Zhang L, Wang J, Wang Y, Zhang Y, Ittmann M. MNX1 is oncogenically upregulated in African-American prostate cancer. *Cancer Res*. (2016) 76:6290–8. doi: 10.1158/0008-5472.CAN-16-0087
  17. von Bergh AR, van Drunen E, van Wering ER, van Zutven LJ, Hainmann I, Lönnerholm G, et al. High incidence of t (7; 12)(q36; p13) in infant AML but not in infant ALL, with a dismal outcome and ectopic expression of HLXB9. *Genes Chromosomes Cancer*. (2006) 45:731–9. doi: 10.1002/gcc.20335
  18. Li P, Fu D, Sheng Q, Yu S, Bao X, Lv Z. TUDCA attenuates intestinal injury and inhibits endoplasmic reticulum stress-mediated intestinal cell apoptosis in necrotizing enterocolitis. *Int Immunopharmacol*. (2019) 74:105665. doi: 10.1016/j.intimp.2019.05.050
  19. Sodhi CP, Shi XH, Richardson WM, Grant ZS, Shapiro RA, Prindle T Jr, et al. Toll-like receptor-4 inhibits enterocyte proliferation via impaired  $\beta$ -catenin signaling in necrotizing enterocolitis. *Gastroenterology*. (2010) 138:185–96. doi: 10.1053/j.gastro.2009.09.045
  20. Chen J, Yang C, Xu X, Yang Y, Xu B. The effect of focal cerebral ischemia-reperfusion injury on TLR4 and NF- $\kappa$ B signaling pathway. *Exp Ther Med*. (2018) 15:897–903. doi: 10.3892/etm.2017.5463
  21. Zhang Q, Lai S, Hou X, Cao W, Zhang Y, Zhang Z. Protective effects of PI3K/Akt signal pathway induced cell autophagy in rat knee joint cartilage injury. *Am J Transl Res*. (2018) 10:762–70.
  22. Dhas DB, Ashmi A, Bhat B, Parija SC, Banupriya N. Modified low cost SNP genotyping technique using cycle threshold (Ct) & melting temperature (Tm) values in allele specific real-time PCR. *Indian J Med Res*. (2015) 142:555. doi: 10.4103/0971-5916.171282
  23. Hilleagass LM, Griswold DE, Brickson B, Albrightsonwinslow C. Assessment of myeloperoxidase activity in whole rat kidney. *J Pharmacol Methods*. (1990). 24:285–95. doi: 10.1016/0160-5402(90)90013-B
  24. Halpern MD, Holubec H, Dominguez JA, Meza YG, Williams CS, Ruth MC, et al. Hepatic inflammatory mediators contribute to intestinal damage in necrotizing enterocolitis. *Am J Gastrointest Liver Physiol*. (2003) 284:G695–702. doi: 10.1152/ajpgi.00353.2002
  25. Sharma R, Hudak ML. A clinical perspective of necrotizing enterocolitis: past, present, and future. *Clin Perinatol*. (2013) 40:27–51. doi: 10.1016/j.clp.2012.12.012
  26. Zhang L, Cheng J, Fan X. MicroRNAs: new therapeutic targets for intestinal barrier dysfunction. *World J Gastroenterol*. (2014) 20:5818–25. doi: 10.3748/wjg.v20.i19.5818
  27. Chivukula R, Shi G, Acharya A, Mills E, Zeitels L, Anandam J, et al. An essential mesenchymal function for miR-143/145 in intestinal epithelial regeneration. *Cell*. (2014) 157:1104–16. doi: 10.1016/j.cell.2014.03.055
  28. Chen W-X, Ren L-H, Shi R-H. Implication of miRNAs for inflammatory bowel disease treatment: systematic review. *World J Gastrointest Pathophysiol*. (2014) 5:63–70. doi: 10.4291/wjgp.v5.i2.63
  29. Chassin C, Kocur M, Pott J, Duerr C, Gütle D, Lotz M, et al. miR-146a mediates protective innate immune tolerance in the neonate intestine. *Cell Host Microbe*. (2010) 8:358–68. doi: 10.1016/j.chom.2010.09.005
  30. Ayyadurai S, Charania M, Xiao B, Viennois E, Zhang Y, Merlin D. Colonic miRNA expression/secretion, regulated by intestinal epithelial PePT1, plays an important role in cell-to-cell communication during colitis. *PLoS ONE*. (2014) 9:e87614. doi: 10.1371/journal.pone.0087614
  31. Bian Z, Li L, Cui J, Zhang H, Liu Y, Zhang C-Y, et al. Role of miR-150-targeting c-Myb in colonic epithelial disruption during dextran sulphate sodium-induced murine experimental colitis and human ulcerative colitis. *J Pathol*. (2011) 225:544–53. doi: 10.1002/path.2907
  32. Qiu W, Kassem M. miR-141-3p inhibits human stromal (mesenchymal) stem cell proliferation and differentiation. *Biochim Biophys Acta*. (2014) 1843:2114–21. doi: 10.1016/j.bbamcr.2014.06.004
  33. Feng J, Xue S, Pang Q, Rang Z, Cui F. miR-141-3p inhibits fibroblast proliferation and migration by targeting GAB1 in keloids. *Biochem Biophys Res Commun*. (2017) 490:302–8. doi: 10.1016/j.bbrc.2017.06.040
  34. Ranjha D, Aggarwal S, Bopanna S, Ahuja V, Paul J. Site-specific MicroRNA expression may lead to different subtypes in ulcerative colitis. *PLoS ONE*. (2015) 10:e0142869. doi: 10.1371/journal.pone.0142869
  35. Pan A, Tan Y, Wang Z, Xu G. STAT4 silencing underlies a novel inhibitory role of microRNA-141-3p in inflammation response of mice with experimental autoimmune myocarditis. *Am J Physiol Heart Circul Physiol*. (2019) 317:H531–40. doi: 10.1152/ajpheart.00048.2019
  36. Cheng L-B, Li K-R, Yi N, Li X-M, Wang F, Xue B, et al. miRNA-141 attenuates UV-induced oxidative stress via activating Keap1-Nrf2 signaling in human retinal pigment epithelium cells and retinal ganglion cells. *Oncotarget*. (2017) 8:13186–94. doi: 10.18632/oncotarget.14489
  37. Yang X, Pan Q, Lu Y, Jiang X, Zhang S, Wu J. MNX1 promotes cell proliferation and activates Wnt/ $\beta$ -catenin signaling in colorectal cancer. *Cell Biol Int*. (2019) 43:402–8. doi: 10.1002/cbin.11096
  38. Chen M, Wu R, Li G, Liu C, Tan L, Xiao K, et al. Motor neuron and pancreas homeobox 1/HLXB9 promotes sustained proliferation in bladder cancer by upregulating CCNE1/2. *J Exp Clin Cancer Res*. (2018) 37:154. doi: 10.1186/s13046-018-0829-9
  39. Xiao L, Hong L, Zheng W. Motor neuron and pancreas homeobox 1 (MNX1) is involved in promoting squamous cervical cancer proliferation via regulating cyclin E. *Med Sci Monit*. (2019) 25:6304. doi: 10.12659/MSM.914233
  40. Quan B, Zhang H, Xue R. miR-141 alleviates LPS-induced inflammation injury in WI-38 fibroblasts by up-regulation of NOX2. *Life Sci*. (2019) 216:271–8. doi: 10.1016/j.lfs.2018.11.056
  41. Nagel S, Scherr M, Quentmeier H, Kaufmann M, Zaborski M, Drexler H, et al. HLXB9 activates IL6 in Hodgkin lymphoma cell lines and is regulated by PI3K signalling involving E2F3. *Leukemia*. (2005) 19:841–6. doi: 10.1038/sj.leu.2403716
  42. Lin L, Liang H, Wang Y, Yin X-M, Hu Y, Huang J, et al. MicroRNA-141 inhibits cell proliferation and invasion and promotes apoptosis by targeting

- hepatocyte nuclear factor-3 $\beta$  in hepatocellular carcinoma cells. *BMC Cancer*. (2014) 14:879. doi: 10.1186/1471-2407-14-879
43. Zhou X, Wu W, Zeng A, Nie E, Jin X, Yu T, et al. MicroRNA-141-3p promotes glioma cell growth and temozolomide resistance by directly targeting p53. *Oncotarget*. (2017) 8:71080–94. doi: 10.18632/oncotarget.20528
44. Zhang X, Wu C, Xiong W, Chen C, Li R, Zhou G. Knockdown of p54nrb inhibits migration, invasion and TNF- $\alpha$  release of human acute monocytic leukemia THP1 cells. *Oncol Rep*. (2016) 35:3742–8. doi: 10.3892/or.2016.4756
45. Dong Q, Jie Y, Ma J, Li C, Yang D. Renal tubular cell death and inflammation response are regulated by the MAPK-ERK-CREB signaling pathway under hypoxia-reoxygenation injury. *J Recept Sig Transd*. (2019) 39:1–9. doi: 10.1080/10799893.2019.1698050
46. Kabakov AE, Gabai VL. Cell death and survival assays. *Methods Mol Biol*. (2018) 1709:107–27. doi: 10.1007/978-1-4939-7477-1\_9

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Chen, Zeng, Zheng, Li and Lin. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.