



Prevalence Mapping of *Schistosoma mansoni* Among Pre-school Age Children in Rwanda

Nadine Rujeni^{1*}, Jean Bosco Bayingana^{1,2}, Elias Nyandwi³, Amans Ntakarutimana¹, Joseph Kagabo¹, Reverien Rutayisire¹, Eliah Shema¹, Philbert Kanimba¹, Jean Bosco Mbonigaba⁴ and Eugene Ruberanziza⁴

¹ School of Health Sciences, College of Medicine and Health Sciences, University of Rwanda, Kigali, Rwanda, ² Single Project Implementation Unit, University of Rwanda, Kigali, Rwanda, ³ Centre for Geographic Information Systems and Remote Sensing, College of Science and Technology, University of Rwanda, Kigali, Rwanda, ⁴ Neglected Tropical Diseases and Other Parasitic Diseases Unit, Rwanda Biomedical Center, Ministry of Health, Kigali, Rwanda

OPEN ACCESS

Edited by:

Edward Oliveira,
René Rachou Institute, Oswaldo Cruz
Foundation (Fiocruz), Brazil

Reviewed by:

Tadesse Hailu,
Bahir Dar University, Ethiopia
Arthur Vengesai,
Midlands State University, Zimbabwe
Lim Boon Huat,
Universiti Sains Malaysia (USM),
Malaysia
Takafira Mduluzi,
University of Zimbabwe, Zimbabwe

*Correspondence:

Nadine Rujeni
nrujeni@gmail.com

Specialty section:

This article was submitted to
Pediatric Infectious Diseases,
a section of the journal
Frontiers in Pediatrics

Received: 28 March 2022

Accepted: 20 May 2022

Published: 24 June 2022

Citation:

Rujeni N, Bayingana JB, Nyandwi E, Ntakarutimana A, Kagabo J, Rutayisire R, Shema E, Kanimba P, Mbonigaba JB and Ruberanziza E (2022) Prevalence Mapping of *Schistosoma mansoni* Among Pre-school Age Children in Rwanda. *Front. Pediatr.* 10:906177. doi: 10.3389/fped.2022.906177

Schistosoma mansoni is endemic in Rwanda, and control programs have been implemented with a special focus on school-age children (SAC), ignoring pre-school age children (pre-SAC) for which the actual prevalence of the disease is not well established. This study consisted of a cross-sectional quantitative mapping of the distribution of *Schistosoma mansoni* and identification of associated risk factors among pre-SAC throughout the country. The study covered all the 17 districts of Rwanda endemic for *Schistosoma mansoni*, with a total sample of 4,675 children enrolled from 80 purposively selected villages. The parasitological assessment of children's urine and stool samples was conducted using CCA and Kato Katz methods, respectively, for infection detection. A standard questionnaire was used to collect data on the risk factors, and geospatial assessment was performed using tablets and GPS to record geographic coordinates for plotting locations on maps using ArcGIS software. The overall prevalence of *S. mansoni* infection across the surveyed areas was 24 and 0.8% by CCA and Kato-Katz, respectively. Infection was significantly associated with bathing children in open water bodies. Furthermore, pre-SAC looked after by siblings (sisters) were two times as much likely to be infected compared to those looked after by mothers. Schistosomiasis control interventions are needed for pre-SAC to limit their exposure to open water bodies with expectations of adapted chemotherapy to be availed. Community-based deworming campaigns may be the best way to ensure good treatment coverage of pre-SAC in Rwanda.

Keywords: schistosomiasis, pre-school aged children, risk factors, *Schistosoma mansoni*, Rwanda

INTRODUCTION

Schistosomiasis is a chronic parasitic infection that is caused by trematodes of the genus *Schistosoma*, transmitted through fresh water snails. It can cause vascular damage in individuals and is associated with growth retardation and impaired metabolism and cognition (1, 2). Schistosomiasis remains a public health concern and highly prevalent in low and middle-income countries, including Rwanda, mostly affecting poor communities with no access to potable water (3). Globally, more than 250 million people are affected by schistosomiasis, and sub-Saharan

Africa accounts for 90% of all these infection burdens (4, 5). Control interventions mainly rely on regular targeted treatments (using Praziquantel) of populations in endemic areas, coupled with health education. Although the disease is targeted (as per the WHO road map) for elimination as a public health problem (defined as prevalence of <1% of heavy-intensity schistosomiasis infections) by 2030 (6), control interventions do not currently reach all populations in need in most endemic areas.

For many years, pre-school-aged children (pre-SAC) have been left out in most schistosomiasis treatment programs. This was partly due to the assumption that they are not at great risk of contracting schistosomiasis. However, considerable advances have been made in this regard, and a number of studies have shown that most of the pre-SAC in endemic areas are significantly exposed to schistosomiasis in many countries (7–14). This has prompted the WHO to formulate a new guideline, including recommendations to include pre-SAC aged 2 years+ in preventive chemotherapy, while those below 2 years should be treated on an individual basis (15).

With the realization that pre-SAC are in need of treatment came a number of challenges, including the absence of an appropriate child-friendly formulation of Praziquantel and the lack of information on the real burden of schistosomiasis in this young age group in some countries. Indeed, inclusion of pre-SAC in schistosomiasis control programs should be justified by quantification of the burden of infection in that age group in each country for optimal intervention. Furthermore, an assessment of their accessibility and local acceptability should guide implementation strategies.

Rwanda is among the countries where *S. mansoni* is endemic. It has been proved by three nationwide schistosomiasis mapping and routine health data that, in Rwanda, only *S. mansoni* is prevalent and no *Schistosoma haematobium* has ever been detected. However, the burden of schistosomiasis among pre-SAC is not well documented. Our own study on Nkombo Island in the western part of the country reported an overall prevalence of 9.5% (9), but a countrywide assessment of the disease in this age group is necessary to better inform the control program.

The Rwandan Ministry of Health has recently introduced a community-based deworming strategy whereby community health workers (CHWs) and community leaders coordinate MDAs with minimal supervision from the central level. Such a deworming strategy should be suitable for treating schistosomiasis among pre-SAC. However, this should be guided by a detailed map of the disease distribution in the country.

In light of the above, this study was conducted to estimate the prevalence, identify risk factors, and produce an illustrative, countrywide map of *S. mansoni* distribution in all moderate- to high-risk areas of the country.

MATERIALS AND METHODS

Study Design, Population, and Sample Size

This was a cross-sectional quantitative study that involved parasitological assessment of schistosomiasis among children

aged 7–68 months. Structured interviews were also held with the participants' parents/caregivers to assess risk factors of infection in this young population. The study covered all the 17 districts of Rwanda at moderate to high risk of schistosomiasis (16, 17) and targeted 40 sectors reported with prevalence of 10% and above among school children (based on incidence and mapping data) (16, 18). From each targeted sector, 2 villages were selected based on their proximity to existing water bodies and/or wetlands. Because of the focal distribution of schistosomiasis, such purposive samplings have been recommended in mapping studies to better guide interventions toward schistosomiasis elimination (19, 20). In our study, the villages layer was intersected with layers of lakes, multipurpose water dams, fish ponds, and/or important wetlands (wetland of ≥ 0.7 ha hosting a socioeconomic activity, mostly irrigated agriculture). From eligible villages (high-risk villages), final selection was purposively decided, considering the spatial distribution pattern analyzed visually by researchers and complemented with field verification, making 80 study villages with proximity to open water sources and wetlands. A total of 4,675 children were randomly selected and enrolled from the purposively selected villages.

Parasitological Assessment

Urine samples were collected and tested for schistosomiasis-circulating antigen using the Point-Of-Care Circulating Cathodic Antigen (POC-CCA or CCA), following the manufacturer's instructions. CCA test results were recorded as negative (– or trace) or positive (+; ++ or +++ according to the intensity of the test line in comparison to a test control). Stool samples were also collected and tested using the Kato Katz method for the detection of patient infection, following published protocols (21). A single specimen was collected from which 2 slides were prepared and read by 2 laboratory technicians independently. Ten per cent (10%) of all the slides were retested by the National Reference Laboratory senior technicians for quality control. The POC-CCA estimate was considered as the overall prevalence, given the high sensitivity of the assay (16). However, KK results were considered for infection intensity, where raw fecal worm egg count (FWEC) was multiplied by 24 to estimate the eggs per gram (epg) (21, 22). The level of infection intensity was determined based on WHO guidelines (23).

Risk Factors and Spatial Assessment

A standard questionnaire was designed in the Kobo toolbox and used to collect data from parents/guardians by a well-trained data collector using a tablet. Geographic coordinates were recorded using tablets, and ArcGIS software version 10.4 was used to produce maps.

Statistical Analysis

Data collected were analyzed by Stata 13 (STATA Corp., College Station, TX, United States). Descriptive analysis was conducted. Categorical variables were presented as frequencies and their respective percentages. Numerical variables were presented as mean and standard deviations. For comparison purposes, cross-tabulation analyses were conducted, and contingency tables

were produced, while chi-square tests were used to compare proportions. To assess the association between variables, logistic regression was performed. For binary logistic regression, each risk factor was assessed with the outcome (infection status) separately, while, for multivariable logistic regression, a backward stepwise logistic regression was run, and all risk factors with a *p*-value less or equal to 0.05 were considered in the model. Odds ratios and the corresponding 95% confidence interval (CI) were reported. A two-tailed significance level of 0.05 was considered in all analyses, and only those variables that were significant in bivariate analysis were included in multivariate analysis.

RESULTS

Demographic Characteristics of the Study Participants

The total sample was 4,675 pre-SAC, with a mean age of 37.5 months (± 13.0), and a sex ratio M:F of 1.01: 1 (Table 1). Majority of the study participants (corresponding to 96.2%) were born in their respective villages (permanent residents).

Water Contact Activities (Exposure) for Children in Study Areas

More than 90% parents reported farming as their occupation, and 83.3% (*n* = 3,895) of the families use open water sources (lakes, dams, or swamps) for daily domestic activities. This results in more than 60% of enrolled children being regularly in contact with open water sources (Table 2).

Prevalence and Infection Intensity of Schistosomiasis Among Pre-school-Aged Children

Prevalence of schistosomiasis among pre-school-aged children was assessed using both KK and POC-CCA. The POC-CCA

results yielded prevalence of 24% among children tested (trace results considered negative), while the KK only yielded 0.8% prevalence. Based on WHO guidelines, 16.2% (*n* = 6) of the 37 children who tested positive on Kato Katz had moderate infection, while the rest had light infection (Table 3).

Prevalence of Schistosomiasis by Age, Gender, and Location Among Pre-school Children

The results on the prevalence of schistosomiasis infection by gender revealed no statistically significant difference in infection between males and females. However, a statistically significant difference in infection was observed across age groups, the youngest having the lowest infection burden (Table 4).

TABLE 1 | Characteristics of pre-school aged children enrolled in the study.

	<i>n</i>	%
Age in months (Mean, SD) (37.5, 13.0)		
Age categories (in months)		
12 months and less	73	1.6
13–24 months	842	18
25–36 months	1230	26.3
37–48 months	1359	29.1
49 months and above	1171	25
Total	4675	100
Sex		
Female	2317	49.6
Male	2358	50.4
Total	4675	100
Child born in village		
No	177	3.8
Yes	4498	96.2
Total	4675	100

TABLE 2 | Water contact activities for parents and their children in the study areas.

	<i>n</i>	%
Parent's occupation		
Casual worker	143	3.18
Farmer	4,060	90.26
Government employee	52	1.16
Self employed	157	3.49
Unemployed	86	1.91
Total	4,498	100
Fetching water in the open water source		
No	780	16.7
Yes	3895	83.3
Total	4675	100
Child taken to open water source		
No	1444	30.9
Yes	3222	69.1
Total	4666	100
Frequency child taken to lake		
Once a day	2055	63.4
Once a week	824	25.4
Once a month	185	5.7
Once a year	53	1.6
Once in his life	123	3.8
Total	3240	100
Child bathed in the lake		
No	1669	36.2
Yes	2937	63.8
Total	4606	100
Frequency child bathed IN the lake		
Once a day	1943	65.2
Once a week	761	25.5
Once a month	155	5.2
Once a year	38	1.3
Once in his life	83	2.8
Total	2980	100
Child ever bathed using FETCHED lake water		
No	947	20.3
Yes	3728	79.7
Total	4675	100

TABLE 3 | Prevalence and infection intensity of the study population.

	<i>n</i>	%
POC-CCA test		
Negative	3554	76
Positive	1121	24
Total	4675	100
KK test results		
Negative	4638	99.2
Positive	37	0.8
Total	4675	100
Intensity of infection		
Light	31	83.8
Moderate	6	16.2
Heavy	0	0
Total	37	100

Bold number is the percentage of infected children (prevalence).

TABLE 4 | Infection prevalence per age category and gender.

Age	CCA – traces considered negative		KK		
	Negative	Positive	Negative	Positive	
≤12 months	<i>n</i>	52	21	73	0
	%	1.5	1.9	1.6	0.0
13–24 months	<i>n</i>	617	225	841	1
	%	17.4	20.1	18.1	2.7
25–36 months	<i>n</i>	915	315	1222	8
	%	25.7	28.1	26.3	21.6
37–48 months	<i>n</i>	1068	291	1346	13
	%	30.1	26.0	29.0	35.1
≥49 months	<i>n</i>	902	269	1156	15
	%	25.4	24.0	24.9	40.5
		χ^2 (<i>P</i> -value) = 11.7493 (0.019)		χ^2 (<i>P</i> -value) = 9.7921 (0.044)	
Gender					
Female	<i>n</i>	1782	535	2298	19
	%	50.1	47.7	49.5	51.4
Male	<i>n</i>	1772	586	2340	18
	%	49.9	52.3	50.5	48.6
		χ^2 (<i>P</i> -value) = 1.9890 (0.158)		χ^2 (<i>P</i> -value) = 0.0478 (0.827)	

Bold numbers are the percentages of infected children in each age group. Significant p-values (for X2 tests comparing age groups).

There were significant differences in infection prevalence across study sites, Gisagara district having the highest prevalence, with 60.4%, followed by Nyagatare and Gasabo districts, respectively, with 46.8 and 40%, while Kamonyi district had the lowest prevalence (7.1%). Infection distribution across study sites is illustrated on **Figure 1**.

Factors Associated With Schistosomiasis Infection

The logistic regression performed highlighted the age as a contributing factor to infection status as expected. In addition, two behavioral factors were significantly associated with the

occurrence of infection (**Table 5**): children taken care of by a sister, and those who are bathed in the lake/pond were approximately 2 times more at risk of being infected (OR = 1.9, $p < 0.0001$ and OR = 1.8, $p < 0.0001$, respectively) compared to their counterparts.

DISCUSSION

The present study aimed to estimate the burden of schistosomiasis among pre-SAC in Rwanda and risk factors for the infection. Overall, the prevalence of *Schistosoma mansoni* infection was 24% using the CCA diagnostic method and 0.8% using the KK method. The high positivity of the CCA compared to KK has well been documented in other settings (24–26), and this may be related to the fact that pre-patent and single-sex infections are detected by CCA as opposed to KK, which only detects eggs in stool (26, 27). This is more relevant in pre-SAC, who are most likely carrying pre-patent and/or light infections (27). However, although there is a lack of a more sensitive and more specific field diagnostic tool for Schistosomiasis mapping, one of the limitations of the CCA is the possibility of false positives in low endemic settings (28). Nevertheless, our surveyed areas were known to have a moderate to high prevalence of *S. mansoni*, which would limit the number of false positives.

The CCA prevalence detected was comparable to findings among SAC in the same areas (16), highlighting an urgent need to include these children in treatment campaigns. As expected, the distribution of infection was skewed, few sectors harboring most of the infection burden as highlighted by the significant differences between sites (i.e., 60.5%, maximum; and 7.1%, minimum). In agreement with the previous mapping among SAC (16, 17), sectors of Nyagatare and Gisagara districts showed the highest infection prevalence (in 4 sectors, the prevalence exceeded 50%), but, also, the districts of Ngoma, Bugesera, Gatsibo, Kayonza, Nyamasheke, Gicumbi, and Karongi were heavily burdened. With regard to infection intensity, and based on WHO criteria (23), most of the pre-SAC showed light-to-moderate infection intensities.

In this study, 83.3% of the parents/guardians fetched their water for domestic use from open water sources, and 63.4% confirmed taking their children to these water sites every day. Most importantly, 63.8% of the parents/guardians interviewed bath their children in open water bodies, and our logistic regression analysis indicates that their children were two times as much likely to be infected compared to those who were not bathed in these waters. Previous studies elsewhere have also reported this passive nature of exposure to schistosomiasis among pre-SAC (10, 11, 29). Interestingly, our logistic regression highlights that children who are looked after by a sibling (a sister in this case) – most likely while their parents conduct domestic/professional chores – are two times more likely to be infected compared to those who are, constantly, with their mothers. Although we were not able to conduct a thorough observational investigation, these findings suggest that older sisters play and/or bathe their younger siblings at water




FIGURE 1 | A map illustrating the distribution of *Schistosoma mansoni* infection among pre-SAC in Rwanda. Infection prevalence was based on CCA findings (trace as negative); geographical coordinates were collected for each site (village) and were extrapolated to the lowest administrative boundary (sector).

contact points that are far enough from where mothers wash dishes/clothes and, therefore, far from detergents that may destroy cercaria (30) or snails (31). These detergents may – to

some extent - protect those children who are, constantly, with their mothers. Another explanation could be the fact that older siblings spend more time with pre-SAC than mothers who are busy with domestic chores or income-generating activities away from water sites. It is also possible that water contact sites for older siblings are more stagnant (for playing purposes) than contact sites for mothers who need more “flowing” (and therefore “cleaner”) water for domestic use.

TABLE 5 | Association between behavioral factors and schistosomiasis.

CCA Results	Odds ratio	Std. Err.	P	95% Conf. Interval
Age group				
≤12 months	REF			
13–24 months	0.80	0.21	0.39	0.48 1.34
25–36 months	0.72	0.18	0.20	0.43 1.19
37–48 months	0.58	0.15	0.03	0.35 0.96
≥49 months	0.60	0.16	0.05	0.36 1.01
Child carer				
Mum	REF			
Dad	0.81	0.13	0.19	0.58 1.11
Sister	1.91	0.32	<0.0001	1.37 2.66
Brother	0.37	0.20	0.07	0.13 1.07
Other	0.50	0.06	<0.0001	0.40 0.63
Child bathed in Lake				
No	REF			
Yes	1.82	0.15	<0.0001	1.56 2.14

Bold numbers indicate significant p-values.

This study was limited in terms of depth of health education assessment as the main focus was on screening infection in the child and interviewing the caregivers to assess exposure risk factors. Nevertheless, given the findings on infection prevalence, schistosomiasis control interventions are specifically needed that focus on health education for household members’ behavior change and practices to (a) limit the children’s exposure to open water bodies and (b) improve children’s environmental sanitary conditions (32). Additional interventions, such as provision of potable water supplies for these vulnerable communities, as well as public latrines, may provide longer term solutions. Indeed, more than 90% of the interviewed parents/guardians are farmers, and most of our study sites were in close proximity to wetlands that are exploited for agriculture. Farmers generally spend several hours in the fields and, in the absence of latrines, are likely to defecate in these fields and contaminate the water bodies that are later used for bathing children. Investigations into *S. mansoni*

intermediate hosts are ongoing, but preliminary results indicate that they are present in most water bodies included in our study (unpublished data).

Taken together, the present study findings stress that pre-SAC in Rwanda are part of the population at risk of schistosomiasis, and control interventions of the disease in Rwanda must include them. In particular, we provide a detailed map of pre-SAC infection distribution that should guide intervention strategies countrywide and prepare the ground for Mass Drug Administration (MDA) among pre-SAC once the Praziquantel (PZQ) pediatric formulation becomes available. Furthermore, the Rwandan Ministry of Health has recently started implementing a community-based deworming that takes place during the mother-child health (MCH) week, which should be the best way to reach pre-SAC.

Further assessment of the parents/CHW knowledge, attitudes, and practices toward schistosomiasis and WASH-recommended practices is recommended. We also recommend an integrated control of schistosomiasis in the country, including MDA, WASH interventions, and vector control.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by the Institutional Review Board (IRB) of the

University of Rwanda, College of Medicine and Health Sciences (Approval Notice No. 148/CMHS/IRB/2019). Written informed consent to participate in this study was provided by the participants' legal guardian/next of kin.

AUTHOR CONTRIBUTIONS

NR and ER designed and coordinated the study. EN coordinated the geographical mapping of study sites and produced the map. JB, AN, JM, PK, JK, and ES conducted the field work and parasitological analyses. JB and RR conducted the statistical analyses. JB drafted the manuscript. NR secured funding and critically reviewed the manuscript. All authors had reviewed the manuscript before submission.

FUNDING

This research was commissioned by the National Institute for Health Research (NIHR) Global Health Research Program (16/136/33) using United Kingdom aid from the United Kingdom Government.

ACKNOWLEDGMENTS

We would like to extend thanks to the study participants: Ministry of Health, District Hospitals and Health Centers for all facilitations to this research activity.

REFERENCES

- Gasparotto J, Senger MR, Telles de Sá Moreira E, Brum PO, Carazza Kessler FG, Peixoto DO, et al. Neurological impairment caused by *Schistosoma mansoni* systemic infection exhibits early features of idiopathic neurodegenerative disease. *J Biol Chem.* (2021) 297:100979. doi: 10.1016/j.jbc.2021.100979
- Osakunor DNM, Mduluzi T, Osei-Hyiaman D, Burgess K, Woolhouse MEJ, Mutapi F. *Schistosoma haematobium* infection is associated with alterations in energy and purine-related metabolism in preschool-aged children. *PLoS Negl Trop Dis.* (2020) 14:e0008866. doi: 10.1371/journal.pntd.0008866
- Rujeni N, Morona D, Ruberanziza E, Mazigo HD. *Schistosomiasis* and soil-transmitted helminthiasis in Rwanda: an update on their epidemiology and control. *Infect Dis Poverty.* (2017) 6:8. doi: 10.1186/s40249-016-0212-z
- Engels D, Chitsulo L, Montresor A, Savioli L. The global epidemiological situation of *Schistosomiasis* and new approaches to control and research. *Acta Tropica.* (2002) 82:139–46. doi: 10.1016/s0001-706x(02)00045-1
- Sokolow SH, Wood CL, Jones IJ, Swartz SJ, Lopez M, Hsieh MH, et al. Global assessment of *Schistosomiasis* control over the past century shows targeting the snail intermediate host works best. *PLoS Negl Trop Dis.* (2016) 10:e0004794. doi: 10.1371/journal.pntd.0004794
- WHO. *Ending-the-Neglect-to-Attain-the-SDGs-NTD-Roadmap.* Geneva: WHO (2021).
- Mutapi F, Rujeni N, Bourke C, Mitchell K, Appleby L, Nausch N, et al. *Schistosoma haematobium* treatment in 1-5 year old children: safety and efficacy of the antihelminthic drug praziquantel. *PLoS Negl Trop Dis.* (2011) 5:e1143. doi: 10.1371/journal.pntd.0001143
- Imai N, Rujeni N, Nausch N, Bourke CD, Appleby LJ, Cowan G, et al. Exposure, infection, systemic cytokine levels and antibody responses in young children concurrently exposed to *Schistosomiasis* and malaria. *Parasitology.* (2011) 138:1519–33. doi: 10.1017/S0031182011001181
- Rujeni N, Mazimpaka A, Tumusiime M, Nyandwi E, Rutayisire G, Kayiranga P, et al. Pre-school aged children are exposed to schistosoma through lake Kivu in Rwanda. *AAS Open Res.* (2019) 2:1–13. doi: 10.12688/aasopenres.12930.1
- Stothard JR, Gabrielli AF. *Schistosomiasis* in African infants and preschool children: to treat or not to treat? *Trends Parasitol.* (2007) 23:83–6.
- Poole H, Terlouw DJ, Naunje A, Mzembe K, Stanton M, Betson M, et al. *Schistosomiasis* in pre-school-age children and their mothers in Chikhwawa district, Malawi with notes on characterization of schistosomes and snails. *Parasit Vectors.* (2014) 7:153. doi: 10.1186/1756-3305-7-153
- Sousa-figueiredo JC, Pleasant J, Day M, Betson M, Rollinson D, Montresor A, et al. Treatment of intestinal *Schistosomiasis* in Ugandan preschool children: best diagnosis, treatment efficacy and side-effects, and an extended praziquantel dosing pole. *Int Health.* (2010) 2:103–13. doi: 10.1016/j.inhe.2010.02.003
- Ekpo UF, Oluwole AS, Abe EM, Etta HE, Olamiju F, Mafiana CF. *Schistosomiasis* in infants and pre-school-aged children in sub-Saharan Africa: implication for control. *Parasitology.* (2012) 139:835–41. doi: 10.1017/S0031182012000029
- Ekpo UF, Laja-deile A, Oluwole AS, Sam-wobo SO, Mafiana CF. Urinary *Schistosomiasis* among preschool children in a rural community near Abeokuta Nigeria. *Parasit Vectors.* (2010) 3:58. doi: 10.1186/1756-3305-3-58
- World Health Organization [WHO]. *WHO Guideline on control and Elimination of Human Schistosomiasis. Evidence-Based Recommendations.* Geneva: World Health Organization (2022).
- Ruberanziza E, Wittmann U, Mbituyumuremyi A, Mutabazi A, Campbell CH, Colley DG, et al. Nationwide remapping of *Schistosoma mansoni* infection in Rwanda using circulating cathodic antigen rapid test: taking steps toward

- elimination. *Am J Trop Med Hyg.* (2020) 103:315–24. doi: 10.4269/ajtmh.19-0866
17. Clark NJ, Umulisa I, Ruberanziza E, Owada K, Colley DG, Ortu G, et al. Mapping *Schistosoma mansoni* endemicity in Rwanda: a critical assessment of geographical disparities arising from circulating cathodic antigen versus Kato-Katz diagnostics. *PLoS Negl Trop Dis.* (2019) 13:e0007723. doi: 10.1371/journal.pntd.0007723
 18. Nyandwi E, Osei FB, Veldkamp T, Amer S. Modeling *Schistosomiasis* spatial risk dynamics over time in Rwanda using zero-inflated poisson regression. *Sci Rep.* (2020) 10:19276. doi: 10.1038/s41598-020-76288-8
 19. Stothard JR, Rollinson D, Tchuente LT. Precision mapping : an innovative tool and way forward to shrink the map, better target interventions, and accelerate toward the elimination of *Schistosomiasis*. *PLoS Negl Trop Dis.* (2018) 12:e0006563. doi: 10.1371/journal.pntd.0006563
 20. Tchuente LA, Rollinson D, Stothard JR, Molyneux D. Moving from control to elimination of *Schistosomiasis* in sub-Saharan Africa: time to change and adapt strategies. *Infect Dis Poverty.* (2017) 6:1–14. doi: 10.1186/s40249-017-0256-8
 21. Barbosa CS, Gomes ECS, Marcelino JMR, Cavalcante KRLJ, Nascimento WRC. Quality control of the slides by Kato-Katz method for the parasitological diagnosis of *Schistosomiasis* infection by *Schistosoma mansoni*. *J Bras Patol Med Lab.* (2017) 53:110–4.
 22. Leuenberger A, Nassoro T, Said K, Fenner L, Sikalengo G, Letang E, et al. Assessing stool quantities generated by three specific Kato-Katz thick smear templates employed in different settings. *Infect Dis Poverty.* (2016) 5:58. doi: 10.1186/s40249-016-0150-9
 23. Sturrock RF. Guidelines for the evaluation of soil-transmitted helminthiasis and *Schistosomiasis* at community level: a guide for managers of control programmes. A. Montresor, DWT Crompton, DAP Bundy, A Hall & L. Savioli. Geneva: world health organization, 1998. *Trans R Soc Trop Med Hyg.* (1998) 92:470–1.
 24. Ferreira FT, Fidelis TA, Pereira TA, Otoni A, Queiroz LC, Amâncio FF. Sensitivity and specificity of the circulating cathodic antigen rapid urine test in the diagnosis of *Schistosomiasis mansoni* infection and evaluation of morbidity in a low- endemic area in Brazil. *Rev Soc Bras Med Trop.* (2017) 50:358–64. doi: 10.1590/0037-8682-0423-2016
 25. Chieng B, Okoyo C, Simiyu E, Gichuki P, Mwatele C, Kepha S, et al. Comparison of quantitative polymerase chain reaction, Kato-Katz and circulating cathodic antigen rapid test for the diagnosis of *Schistosoma mansoni* infection : a cross-sectional study in Kirinyaga County, Kenya. *Curr Res Parasitol Vector Borne Dis.* (2021) 1:100029. doi: 10.1016/j.crvbd.2021.100029
 26. Oliveira WJ, Magalhães FDC, Elias AMS, de Castro VN, Favero V, Lindholz CG, et al. Evaluation of diagnostic methods for the detection of intestinal *Schistosomiasis* in endemic areas with low parasite loads : saline gradient, Helminx, Kato-Katz and rapid urine test. *PLoS Negl Trop Dis.* (2018) 12:e0006232. doi: 10.1371/journal.pntd.0006232
 27. Stothard JR, Sousa-figuereido JC, Betson M, Adriko M, Arinaitwe M, Rowell C, et al. *Schistosoma mansoni* infections in young children : when are Schistosome antigens in Urine, eggs in stool and antibodies to eggs first detectable? *PLoS Negl Trop Dis.* (2011) 5:e938. doi: 10.1371/journal.pntd.0000938
 28. De Sousa SRM, Dias IHL, Fonseca ALS, Contente BR, Nogueira JFC, Da Costa Oliveira TN, et al. Concordance of the point-of-care circulating cathodic antigen test for the diagnosis of intestinal *Schistosomiasis* in a low endemicity area. *Infect Dis Poverty.* (2019) 8:37. doi: 10.1186/s40249-019-0551-7
 29. Osakunor DNM, Woolhouse MEJ, Mutapi F. Paediatric *Schistosomiasis*: what we know and what we need to know. *PLoS Negl Trop Dis.* (2018) 12:e0006144. doi: 10.1371/journal.pntd.0006144
 30. Okwuosa VN, Osuala FO. Toxicity of washing soaps to *Schistosoma mansoni* cercariae and effects of sublethal concentrations on infectivity in mice. *Appl Parasitol.* (1993) 34:69–75.
 31. Abebe F, Erko B, Gemetchu T, Gundersen SG. Control of *Biomphalaria pfeifferi* population and *Schistosomiasis* transmission in Ethiopia using soap berry endod (*Phytolacca dodecandra*), with special emphasis on application methods. *Trans R Soc Trop Med Hyg.* (2005) 99:787–94. doi: 10.1016/j.trstmh.2005.04.013
 32. World Health Organization [WHO]. *Children's Environmental Health Indicators (CEHI): Presenting Regional Successes Learning for the Future: Summary.* Geneva: World Health Organization (2009).
- Author Disclaimer:** The views expressed in this publication are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.
- Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.
- Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.
- Copyright © 2022 Rujeni, Bayingana, Nyandwi, Ntakarutimana, Kagabo, Rutayisire, Shema, Kanimba, Mbonigaba and Ruberanziza. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.