



Methylene Blue Inhibits the SARS-CoV-2 Spike–ACE2 Protein-Protein Interaction—a Mechanism that can Contribute to its Antiviral Activity Against COVID-19

Damir Bojadzic¹, Oscar Alcazar¹ and Peter Buchwald^{1,2*}

¹Diabetes Research Institute, University of Miami, Miami, FL, United States, ²Department of Molecular and Cellular Pharmacology, Miller School of Medicine, University of Miami, Miami, FL, United States

OPEN ACCESS

Edited by:

Rafael Maldonado,
Pompeu Fabra University, Spain

Reviewed by:

Enzo Nisoli,
University of Milan, Italy
Gianluigi Lauro,
University of Salerno, Italy

*Correspondence:

Peter Buchwald
pbuchwald@med.miami.edu

Specialty section:

This article was submitted to
Experimental Pharmacology
and Drug Discovery,
a section of the journal
Frontiers in Pharmacology

Received: 29 August 2020

Accepted: 03 December 2020

Published: 13 January 2021

Citation:

Bojadzic D, Alcazar O and Buchwald P
(2021) Methylene Blue Inhibits the
SARS-CoV-2 Spike–ACE2 Protein-
Protein Interaction—a Mechanism that
can Contribute to its Antiviral Activity
Against COVID-19.

Front. Pharmacol. 11:600372.
doi: 10.3389/fphar.2020.600372

Due to our interest in the chemical space of organic dyes to identify potential small-molecule inhibitors (SMIs) for protein-protein interactions (PPIs), we initiated a screen of such compounds to assess their inhibitory activity against the interaction between SARS-CoV-2 spike protein and its cognate receptor ACE2, which is the first critical step initiating the viral attachment and entry of this coronavirus responsible for the ongoing COVID-19 pandemic. As part of this, we found that methylene blue, a tricyclic phenothiazine compound approved by the FDA for the treatment of methemoglobinemia and used for other medical applications (including the inactivation of viruses in blood products prior to transfusion when activated by light), inhibits this interaction. We confirmed that it does so in a concentration-dependent manner with a low micromolar half-maximal inhibitory concentration ($IC_{50} = 3 \mu M$) in our protein-based ELISA-type setup, while chloroquine, siramesine, and suramin showed no inhibitory activity in this assay. Erythrosine B, which we have shown before to be a promiscuous SMI of PPIs, also inhibited this interaction. Methylene blue inhibited the entry of a SARS-CoV-2 spike bearing pseudovirus into ACE2-expressing cells with similar IC_{50} ($3.5 \mu M$). Hence, this PPI inhibitory activity could contribute to its antiviral activity against SARS-CoV-2 even in the absence of light by blocking its attachment to ACE2-expressing cells and making this inexpensive and widely available drug potentially useful in the prevention and treatment of COVID-19 as an oral or inhaled medication.

Keywords: ACE2, antiviral, chloroquine, COVID-19, methylene blue, protein-protein interaction, SARS-CoV-2, spike protein

Abbreviations: ACE2, angiotensin converting enzyme 2; CoV, coronavirus; MeBlu, methylene blue; PPI, protein-protein interaction; SARS, severe acute respiratory syndrome; SMI, small-molecule inhibitor.

INTRODUCTION

Severe acute respiratory syndrome-coronavirus 2 (SARS-CoV-2), a novel betacoronavirus and the most recent one of the seven coronaviruses (CoVs) known to infect humans, is responsible for COVID-19, which has been declared a pandemic by the World Health Organization in March 2020 and continues to spread worldwide (Liu et al., 2020; Matheson and Lehner, 2020; Moore and June, 2020). While four CoVs (HCoV 229E, OC43, NL63, and HKU1) are responsible for about one third of the common cold cases in humans, three have caused recent epidemics associated with considerable mortality: SARS-CoV-1 (2002–2003, causing ~10% mortality), MERS-CoV (Middle East respiratory syndrome coronavirus; 2012, causing ~35% mortality), and now SARS-CoV-2 (2019–2020), which seems to be less lethal but more transmissible (Guy et al., 2020). SARS-CoV-2 is the most infectious agent in a century (Tiwari et al., 2020) and has already caused infections in the order of tens of millions and deaths that are likely to be in the order of millions worldwide. According to current early estimates, about 3% of infected individuals need hospitalization and 0.5% die, a range that is strongly age-dependent increasing from 0.001% in <20 years old to 8.3% in those >80 years old (Salje et al., 2020). Accordingly, there is considerable interest in possible preventive or therapeutic treatments. There are several possible targets in the coronavirus life cycle for therapeutic interventions including attachment and entry, uncoating, gRNA replication, translation in endoplasmic reticulum (ER) and Golgi, assembly, and virion release (Guy et al., 2020). Viral attachment and entry are particularly promising as they are the first steps in the replication cycle and take place at a relatively accessible extracellular site; hence, they have been explored for intervention purposes for several viruses (Melby and Westby, 2009). CoVs use their glycosylated spike (S) protein to bind to their cognate cell surface receptors and initiate membrane fusion and virus entry. For both SARS-CoV and SARS-CoV-2, the S protein mediates entry into cells by binding to angiotensin converting enzyme 2 (ACE2) via its receptor-binding domain (RBD) followed by proteolytic activation by human proteases (Lan et al., 2020; Matheson and Lehner, 2020; Shang et al., 2020; Sivaraman et al., 2020). Blockade of this RBD–ACE2 protein-protein interaction (PPI) can disrupt infection efficiency; for example, SARS-CoV-2 RBD protein was shown to block S protein mediated SARS-CoV-2 pseudovirus entry into ACE2 receptor-expressing target cells (Tai et al., 2020). Antibodies can be quite effective PPI inhibitors, and they are highly target-specific and relatively stable *in vivo*. However, they cannot reach intracellular targets and, as all other protein therapies, are hindered by problems such as low solubility, propensity for immunogenicity, long elimination half-lives, lack of oral bioavailability, product heterogeneity, and possible manufacturing and storage stability issues. Since they are foreign proteins, they elicit strong immune response in certain patients (Suntharalingam et al., 2006; Wadman, 2006; Leader et al., 2008), and even if approved for clinical use, they

tend to have more post-market safety issues than small-molecule drugs (Downing et al., 2017). Small-molecule inhibitors (SMIs) are more challenging to identify for PPIs, but it is now well established that they can be effective against certain PPIs and can offer useful alternatives. There are now >40 PPIs targeted by SMIs that are in preclinical development, and two such SMIs are approved for clinical use (venetoclax and lifitegrast) (Arkin and Wells, 2004; Milroy et al., 2014; Scott et al., 2016; Bojadzic and Buchwald, 2018).

Due to our interest in the chemical space of organic dyes to identify potential SMIs for PPIs (Margolles-Clark et al., 2009a; Margolles-Clark et al., 2009b; Ganesan et al., 2011; Song et al., 2014; Chen et al., 2017; Bojadzic and Buchwald, 2018; Bojadzic et al., 2018), we initiated a screen of such compounds for their ability to inhibit the interaction between SARS-CoV-2 spike protein and its cognate receptor ACE2, which is the first critical step initiating the viral attachment and entry of this CoV. As part of this, we found that methylene blue, a tricyclic phenothiazine compound approved for the treatment of acquired methemoglobinemia and some other uses (Clifton and Leikin, 2003; Schirmer et al., 2011; Bistas and Sanghavi, 2020), inhibits this interaction, and we have confirmed that it does so in a concentration-dependent manner. This can contribute to the antiviral activity of this inexpensive and widely available dye-based drug against SARS-CoV-2 making it potentially useful in the prevention and treatment of COVID-19, especially in non-industrialized nations.

MATERIALS AND METHODS

Binding Assays

Methylene blue and other test compounds used here were obtained from Sigma-Aldrich (St. Louis, MO, United States) and used as such. Purities (and catalog numbers) were as follows: methylene blue >95% (M4159), chloroquine >98.5% (C6628), erythrosine B 90% (198269), siramesine >98% (SML0976), sunset yellow FCF 90% (465224), and trypan blue 60% (302643). Suramin (>99%; cat. no. 1472) was from Tocris Bioscience (Biotechnique, Minneapolis, MN, United States). ACE2-Fc and SARS-CoV-2 S1 or RBD with His tag proteins used in the binding assays were obtained from Sino Biological (Wayne, PA, United States); catalog no. 10108-H05H, 40591-V08H, and 40592-V08H). Binding inhibition assays were performed in a 96-well cell-free format similar to the one described before (Margolles-Clark et al., 2009b; Ganesan et al., 2011; Song et al., 2014; Chen et al., 2017). Briefly, microtiter plates (Nunc F Maxisorp, 96-well; Thermo Fisher Scientific, Waltham, MA, United States) were coated overnight at 4°C with 100 μ L/well of Fc-conjugated ACE2 receptor diluted in PBS pH 7.2. This was followed by blocking with 200 μ L/well of SuperBlock (PBS) (Thermo Fisher Scientific) for 1 h at room temperature. Then, plates were washed twice using washing solution (PBS pH 7.4, 0.05% Tween-20) and tapped dry before the addition of the tagged ligand (SARS-CoV-2 S1 or RBD) and test compounds diluted in binding buffer (100 mM HEPES, pH 7.2) to give a total


volume of 100 μL /well. After 1 h incubation, three washes were conducted, and a further 1 h incubation with anti-His HRP conjugate (BioLegend; San Diego, CA, United States; catalog no. 652504) diluted (1:2,500) in SuperBlock (PBS) was used to detect the bound His-tagged ligand. Plates were washed four times before the addition of 100 μL /well of HRP substrate TMB (3,3',5,5'-tetramethylbenzidine) and kept in the dark for up to 15 min. The reaction was stopped using 20 μL of 1 M H_2SO_4 , and the absorbance value was read at 450 nm. The plated concentrations of ACE2 receptor were 1.0 $\mu\text{g}/\text{ml}$ for SARS-CoV-2 RBD and 2.0 $\mu\text{g}/\text{ml}$ for SARS-CoV-2 S1. The concentrations of the ligand used in the inhibitory assays were 0.5 $\mu\text{g}/\text{ml}$ for RBD and 1.0 $\mu\text{g}/\text{ml}$ for S1. These values were selected following preliminary testing to optimize response (i.e., to produce a high-enough signal at conditions close to half-maximal response, EC_{50}). Binding assessments for CD40-CD40L and TNF-R1-TNF- α were performed as previously described (Bojadzic et al., 2018). Stock solutions of compounds at 10 mM in DMSO were used.

SARS-CoV-2 Pseudovirus Assay

Assay from Montana Molecular (Bozeman, MT, United States; catalog no. C1100R and C1100G) was used per the instructions of the manufacturer with minor modifications. Briefly, HEK293T cells (ATCC, Manassas, VA, United States; catalog no. CRL-1573) were seeded onto 96-well plates at a density of 5×10^4 cells per well in 100 μL complete medium (DMEM supplemented with 10% fetal bovine serum). A transduction mixture containing ACE2 BacMam Red-Reporter virus (1.8×10^8 VG/ml) and 2 mM sodium butyrate prepared in complete medium was added (50 μL per well) and incubated for 24 h at 37°C and 5% CO_2 . Medium was removed, washed once with PBS, and replaced with 100 μL fresh medium containing methylene blue at selected concentrations, pre-incubating for 30 min at 37°C and 5% CO_2 . A transduction mixture containing Pseudo SARS-CoV-2 Green-Reporter pseudovirus (3.3×10^8 VG/ml) and 2 mM sodium butyrate prepared in complete medium was added (50 μL per well) and incubated for 48 h at 37°C and 5% CO_2 . Medium was removed, washed once with PBS, replaced with 150 μL fresh medium, and cells incubated for additional 48 h at 37°C and 5% CO_2 . Cell fluorescence was detected using an EVOS FL microscope (Life Technologies, Carlsbad, CA, United States) and was quantified in ImageJ (United States National Institutes of Health, Bethesda, MD, United States) (Schneider et al., 2012) using the Analyze Particles tool after thresholding for the corresponding colors.

Statistics and Data Fitting


All binding inhibition and cell assays were tested in at least duplicate per plates, and assays were performed as at least two independent experiments. As before (Ganesan et al., 2011; Song et al., 2014; Chen et al., 2017), binding data were converted to percent inhibition and fitted with standard log inhibitor vs. normalized response models (Buchwald, 2020) using nonlinear regression in GraphPad Prism (GraphPad, La Jolla, CA, United States) to establish half-maximal (median) effective or inhibitory concentrations (EC_{50} , IC_{50}).



RESULTS


As part of our work to identify SMIs for co-signaling PPIs that are essential for the activation and control of immune cells, we discovered that the chemical space of organic dyes, which is particularly rich in strong protein binders, can offer a useful starting point. Accordingly, it seemed logical to explore it for possible inhibitors of the SARS-CoV-2 S protein-ACE2 PPI that is an essential first step for the viral entry of this novel, highly infectious coronavirus. As a first step, we explored the feasibility of setting up a screening assays using a cell-free ELISA-type 96-well format similar to those used in our previous works with Fc-conjugated receptors coated on the plate and FLAG- or His-tagged ligands in the solution (Margolles-Clark et al., 2009b; Ganesan et al., 2011; Song et al., 2014; Chen et al., 2017). To establish assay conditions, we first performed concentration-response assessments using such a format with ACE2-Fc and SARS-CoV-2 S1 or RBD with His tag, and they indicated that both bindings follow classic sigmoid patterns with a slightly stronger binding for RBD than S1 (Figure 1). Fitting of data gave median effective concentrations (EC_{50} s) and hence binding affinity constant (K_d) estimates of 5 and 13 nM, respectively (127 and 1,008 ng/ml)—in good agreement with the specifications of the manufacturer and published values that are also in the low nanomolar range (4–90 nM), typically based on surface plasmon resonance (SPR) studies (Sivaraman et al., 2020).

Accordingly, we can use this format for inhibitory screening, and we decided to use hACE2 with SARS-CoV-2 RBD-His, as it showed stronger binding. In fact, this assay setup is very similar to one recently shown to work as a specific and sensitive SARS-CoV-2 surrogate virus neutralization test based on antibody-mediated blockage of this same PPI (CoV-S-ACE2) (Tan et al., 2020). With



this setup in our hands, we performed a preliminary screening of representative organic dyes from our in-house library plus a few compounds that are or have been considered of possible interest in inhibiting SAR-CoV-2 by different mechanisms of action, e.g., chloroquine, siramesine, and suramin (Colson et al., 2020; Gordon et al., 2020; McKee et al., 2020; Salgado-Benvindo et al., 2020; Xiu et al., 2020). Screening at

5 μ M indicated that most have no activity and, hence, are unlikely to interfere with the S-protein-ACE2 binding needed for viral attachment. Nevertheless, some showed activity; those of selected compounds of interest are shown in **Figure 2** together with corresponding chemical structures. Erythrosine B (ErB, FD and C red no.3), an FDA approved food colorant, was included as a possible positive control since




we have shown it previously to be a promiscuous PPI inhibitor together with other xanthene dyes (Ganesan et al., 2011), and it indeed showed strong inhibition here. Of particular interest, methylene blue (MeBlu), which has a long history of diverse medical applications (Clifton and Leikin, 2003; Schirmer et al., 2011; Bistas and Sanghavi, 2020), also showed promising inhibitory activity.

Therefore, to confirm its activity, we performed detailed concentration-response assessments as recommended per experimental guidelines in pharmacology and experimental biology (Curtis et al., 2018; Michel et al., 2020). As shown in **Figure 3**, this confirmed that MeBlu indeed inhibited in concentration-dependent manner with an estimated IC₅₀ of 3.0 μM (95% CI: 2.5–3.6 μM), whereas chloroquine and suramin showed no inhibitory activity in this assay. Chloroquine, an anti-parasitic and immunosuppressive drug primarily used to prevent and treat malaria, was included as it has potential antiviral activity against SARS-CoV-2 (subject to controversies) (Colson et al., 2020). Suramin, an antiparasitic drug approved for the prophylactic treatment of African sleeping sickness (trypanosomiasis) and river blindness (onchocerciasis), was incorporated because it was claimed to inhibit SARS-CoV-2 infection in cell culture most likely by preventing binding or entry of the virus (Salgado-Benvindo et al., 2020) (as well as because we found it earlier to inhibit the CD40–CD40L PPI (Margolles-Clark et al., 2009a)). ErB also inhibited with an IC₅₀ of 0.4 μM, which is consistent with our previous observation of promiscuous PPI inhibition by this compound with a possibly slightly higher activity than found for other PPIs tested before (1–20 μM)

(Ganesan et al., 2011). Sunset yellow FCF (FD and C yellow no. 6), an azo dye and an FDA approved food colorant included as a possible negative control, indeed showed no inhibitory activity.

Next, we were also able to show that MeBlu inhibits the entry of a pseudovirus bearing the SARS-CoV-2 S spike protein into ACE2-expressing HEK293T cells. This BacMam-based pseudovirus assay allows the quantification of viral entry as the pseudovirus expresses bright green fluorescent protein that is targeted to the nucleus of ACE2 and red fluorescence reporter expressing host cells, while it can be handled using biosafety level 1 containment because they do not replicate in human cells. Pseudovirus entry is indicated by expression of green fluorescence in the nucleus; if the entry is blocked, the cell nucleus remains dark. MeBlu showed clear concentration-dependent inhibition with an estimated IC₅₀ of 3.5 μM (95% CI: 1.6–7.4 μM) (**Figure 4**). In this assay involving 48 h exposure, MeBlu showed signs of cytotoxicity at higher concentrations (45 μM) affecting viability even if not affecting overall ACE2-expression (red bars, **Figure 4**). Since the IC₅₀ obtained for MeBlu here (3 μM) is within the range of its circulating levels following normal clinical dosage (e.g., peak blood concentration of 19 μM after 500 mg p.o. with an elimination half-life of ~14 h (Walter-Sack et al., 2009) or trough levels of 6–7 μM in healthy human volunteers following oral doses of 207 mg/day (69 mg, t.i.d.) (Baddeley et al., 2015), this inhibitory effect on viral attachment can contribute to the possible antiviral activity of MeBlu against SARS-CoV-2 and possibly other ACE2-binding CoVs such as SARS-CoV and the α-coronavirus HCoV NL63.



DISCUSSION

Results here confirm again the usefulness of our strategy to rely on the chemical space of organic dyes, known to contain strong protein binders, as a starting platform to identify SMI scaffolds for PPI inhibition. Using this strategy, we have achieved considerable progress in targeting co-signaling interactions as we have identified the first SMIs for CD40–CD40L (Margolles-Clark et al., 2009b) and OX40–OX40L PPIs (Song et al., 2014) as well as the first promiscuous SMIs of PPIs (Ganesan et al., 2011). Organic dyes contain privileged structures for protein binding (Che et al., 2006; Fletcher and Hamilton, 2006; Hershberger et al., 2007), and, contrary to usual drug-like libraries, whose chemical space does not correspond well with that of promising PPI inhibitors (Neugebauer et al., 2007; Reynès et al., 2010; Sperandio et al., 2010), they are a good starting point to identify SMIs of PPIs. Most dyes, however, are unsuitable for therapeutic development because of their strong color and, in the case of azo dyes, their quick metabolic degradation (Levine, 1991; Feng et al., 2012); hence further medicinal chemistry is needed to optimize their clinical potential (Chen et al., 2017).

More importantly, our results indicate that MeBlu, an organic dye in clinical use for some therapeutic applications in the developed world (Clifton and Leikin, 2003; Schirmer et al., 2011; Bistas and Sanghavi, 2020) and with additional potential for certain developing world applications such as malaria (Dicko et al., 2018), can inhibit the viral attachment and entry of SARS-CoV-2 by blocking the PPI of its spike protein with ACE2 on the host cell. MeBlu is a tricyclic phenothiazine dye approved by the FDA for clinical use for the treatment of methemoglobinemia, and it is also used for other applications such as prevention of urinary tract infections in elderly patients; ifosfamid-induced neurotoxicity in cancer patients; vasoplegic syndrome, a type of distributive shock that occurs during coronary procedures; and intraoperative visualization of nerves, nerve tissues, and

endocrine glands (Schirmer et al., 2011; Bistas and Sanghavi, 2020). MeBlu is included in the WHO List of Essential Medicines and was, in fact, the very first fully synthetic drug used in medicine, as it was used to treat malaria since 1891 (Schirmer et al., 2011). This utilization spanned through WW2 until it was replaced by chloroquine; although, due to the blue urine it could cause, MeBlu was not well liked among the soldiers (“Even at the loo we see, we pee, navy blue”) (Schirmer et al., 2011). It also served as the lead compound for the development of chlorpromazine and tricyclic antidepressants (Schirmer et al., 2011). Moreover, there is resurgent interest in its antimalarial application (Dicko et al., 2018), and it has potential for the treatment of neurodegenerative disorders such as Alzheimer’s disease (AD), due to its putative inhibitory action on the aggregation of tau protein (Schirmer et al., 2011). Notably, MeBlu was also part of the first method developed for pathogen inactivation in plasma, where it has been used since 1991 to inactivate viruses in combination with light (Lozano et al., 2013). MeBlu intercalates within nucleic acid strands, and application of light causes its excitation generating highly reactive singlet oxygen that oxidizes guanosine and breaks nucleic strands (Lozano et al., 2013). Hence, in the presence of light, MeBlu has broad-spectrum virucidal activity and is used to inactivate viruses in blood products prior to transfusions.

Notably, there is also recent evidence of possible *in vitro* antiviral activity for MeBlu even in the absence of UV-induced activation. For example, one group found that MeBlu showed virucidal activity at low micromolar concentrations when incubated with Vero E6 cells and SARS-CoV-2 for 20 h in the dark (Cagno et al., 2020). Another group also found non-photoactivated MeBlu to inhibit SARS-CoV-2 replication in Vero E6 *in vitro* with an IC_{50} of $0.3 \pm 0.03 \mu$ M at multiplicity of infection (MOI) of 0.25 (Gendrot et al., 2020). The ability of MeBlu to inhibit the SARS-CoV-2-S-ACE2 PPI could be a mechanism of action contributing to such activity especially as

we also showed MeBlu to inhibit the entry of SARS-CoV-2 pseudovirus into ACE2-expressing cells with low micromolar IC_{50} (**Figure 4**). If this PPI inhibitory activity of MeBlu is retained at similar levels *in vivo* as found here ($IC_{50} \approx 3 \mu\text{M}$), it is within a range that can be obtained in blood following typical doses (e.g., 200 mg/day) as indicated by pharmacokinetic studies in humans. For example, in one study, peak blood concentration of MeBlu was $19 \mu\text{M}$ after 500 mg p.o., and the elimination half-life was also more than adequate being around 14 h (Walter-Sack et al., 2009). In another study, trough levels of $6\text{--}7 \mu\text{M}$ were obtained following total daily oral doses of 207 mg/day (administered as 69 mg, p.o., t.i.d.) (Baddeley et al., 2015). Hence, oral administration could provide adequate concentrations (e.g., $>7 \mu\text{M}$) and inhaled applications, which have been explored in less developed countries for some respiratory treatments (Golwalkar, 2020), could be even more advantageous. MeBlu is generally safe, but it shows dose-dependent toxicity with nausea, vomiting, hemolysis, and other undesired side effects starting to occur at doses $>7 \text{ mg/kg}$ (i.e., $>500 \text{ mg}$) (Clifton and Leikin, 2003; Bistas and Sanghavi, 2020). It also is contraindicated in certain populations, e.g., in those taking serotonin reuptake inhibitors and in persons with hereditary glucose-6-phosphate dehydrogenase deficiency (G6PD deficiency) (Schirmer et al., 2011; Bistas and Sanghavi, 2020).

It has to be noted, however, that MeBlu also inhibited the CD40–CD40L and TNF-R1–TNF α PPIs in our assays with low-to mid-micromolar potency (data not shown); hence, it is possible that MeBlu is a somewhat promiscuous PPI inhibitor limiting its usefulness. Its three-ring phenothiazine framework resembles somewhat the three-ring xanthene framework of erythrosine B (**Figure 2**), which we have shown before to act as promiscuous PPI inhibitor together with some other structural analog xanthene dyes such as rose Bengal and phloxine (Ganesan et al., 2011). MeBlu certainly shows polypharmacology and acts on a multitude of targets (Schirmer et al., 2011); many of these however can have further beneficial effects in COVID-19 patients (Scigliano and Scigliano, 2020). Its main mechanism of action is reducing the oxidized ferric form of hemoglobin (Fe^{3+}) when in a state of methemoglobinemia, which binds oxygen irreversibly, to the ferrous (Fe^{2+}) form (Bistas and Sanghavi, 2020). This increases the oxygen-binding capacity of hemoglobin and, thus, oxygen delivery to tissues—an important benefit for COVID-19 patients. COVID-19 patients often exhibiting low oxygen levels, typically incompatible with life without dyspnea—a phenomenon termed silent hypoxemia (or happy hypoxia in public media) (Tobin et al., 2020). Possibly relevant to this, MeBlu was found to improve hypoxemia and hyperdynamic circulation in patients with liver cirrhosis and severe hepatopulmonary syndrome (Schenk et al., 2000). MeBlu is being used for the treatment of pneumonia and other respiratory ailments in less developed countries with some success (Golwalkar, 2020).

Further, MeBlu was recently shown to block the PD-1–SHP2 PPI, which is downstream from the PD-1–PD-L1 co-signaling PPI, with low micromolar potency and effectively enough to

counteract the suppressive activity of PD-1 on cytotoxic T lymphocytes and restore their cytotoxicity, activation, proliferation, and cytokine-secreting activity (Fan et al., 2020). This mechanism of action targeting this co-signaling pathway (PD-1) could contribute to restoring T cell homeostasis and function from exhausted state (Barber et al., 2006; Vardhana and Wolchok, 2020), which is of interest to improve viral clearance and rein-in the inflammatory immune response and the associated cytokine storm during anti-viral responses such as those causing the high mortality of COVID-19 patients (Di Cosimo et al., 2020; Liu et al., 2020; Ye et al., 2020).

As far as clinical applications, one promising indication comes from a report of a cohort of 2,500 French patients treated with MeBlu as part of their cancer care none of whom developed influenza like illness during the COVID-19 epidemics (Henry et al., 2020). MeBlu has also been explored in one Phase one clinical trial (NCT04370288) for treatment of critically ill COVID-19 patients in Iran as part of a three-drug last therapeutic option add-on cocktail (MeBlu 1 mg/kg, vitamin C 1500 mg/kg, and N-acetyl cysteine 2000 mg/kg) based on the hypothesis that this combination could rebalance NO, methemoglobin, and oxidative stress. Four of the five patients responded well to treatment (Alamdari et al., 2020).

In conclusion, screening of our organic dye-based library identified MeBlu as a low-micromolar inhibitor of the interaction between SARS-CoV-2 spike protein and its cognate receptor ACE2, a PPI that is the first critical step initiating the viral entry of this coronavirus. While MeBlu shows strong polypharmacology and might be a somewhat promiscuous PPI inhibitor, its ability to inhibit this PPI could contribute to the antiviral activity of MeBlu against SARS-CoV-2 even in the absence of light making this inexpensive and widely available drug potentially useful in the prevention and treatment of COVID-19 as an oral or inhaled medication.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

AUTHOR CONTRIBUTIONS

DB performed most of the experiments, OA performed the pseudovirus assay. PB originated and designed the project, provided study guidance, and wrote the draft manuscript. All authors contributed to writing and read the final manuscript.

ACKNOWLEDGMENTS

Financial support by the Diabetes Research Institute Foundation (www.diabetesresearch.org) is gratefully acknowledged.

REFERENCES

- Alamdari, D. H., Moghaddam, A. B., Amini, S., Keramati, M. R., Zarmehri, A. M., Alamdari, A. H., et al. (2020). Application of methylene blue -vitamin C -N-acetyl cysteine for treatment of critically ill COVID-19 patients, report of a phase-I clinical trial. *Eur. J. Pharmacol.* 885, 173494. doi:10.1016/j.ejphar.2020.173494
- Arkin, M. R., and Wells, J. A. (2004). Small-molecule inhibitors of protein-protein interactions: progressing towards the dream. *Nat. Rev. Drug Discov.* 3 (4), 301–317. doi:10.1038/nrd1343
- Baddeley, T. C., McCaffrey, J., Storey, J. M., Cheung, J. K., Melis, V., Horsley, D., et al. (2015). Complex disposition of methylthioninium redox forms determines efficacy in tau aggregation inhibitor therapy for Alzheimer's disease. *J. Pharmacol. Exp. Therapeut.* 352 (1), 110–118. doi:10.1124/jpet.114.219352
- Barber, D. L., Wherry, E. J., Masopust, D., Zhu, B., Allison, J. P., Sharpe, A. H., et al. (2006). Restoring function in exhausted CD8 T cells during chronic viral infection. *Nature*. 439 (7077), 682–687. doi:10.1038/nature04444
- Bistas, E., and Sanghavi, D. (2020). Methylene blue. (Treasure Island, FL: StatPearls).
- Bojadzic, D., and Buchwald, P. (2018). Toward small-molecule inhibition of protein-protein interactions: general aspects and recent progress in targeting costimulatory and coinhibitory (immune checkpoint) interactions. *Curr. Top. Med. Chem.* 18 (8), 674–699. doi:10.2174/1568026618666180531092503
- Bojadzic, D., Chen, J., Alcazar, O., and Buchwald, P. (2018). Design, synthesis, and evaluation of novel immunomodulatory small molecules targeting the CD40-CD154 costimulatory protein-protein interaction. *Molecules*. 23 (5), 1153. doi:10.3390/molecules23051153
- Buchwald, P. (2020). A single unified model for fitting simple to complex receptor response data. *Sci. Rep.* 10 (1), 13386. doi:10.1038/s41598-020-70220-w
- Cagno, V., Medaglia, C., Cerny, A., Cerny, T., Tapparel, C., and Cerny, E. (2020). Methylene blue has a potent antiviral activity against SARS-CoV-2 in the absence of UV-activation in vitro. Available at: <https://www.biorxiv.org/content/10.1101/2020.08.14.251090v1>.
- Che, Y., Brooks, B. R., and Marshall, G. R. (2006). Development of small molecules designed to modulate protein-protein interactions. *J. Comput. Aided Mol. Des.* 20 (2), 109–130. doi:10.1007/s10822-006-9040-8
- Chen, J., Song, Y., Bojadzic, D., Tamayo-Garcia, A., Landin, A. M., Blomberg, B. B., et al. (2017). Small-molecule inhibitors of the CD40-CD40L costimulatory protein-protein interaction. *J. Med. Chem.* 60 (21), 8906–8922. doi:10.1021/acs.jmedchem.7b01154
- Clifton, J. I., and Leikin, J. B. (2003). Methylene blue. *Am. J. Therapeut.* 10 (4), 289–291. doi:10.1097/00045391-200307000-00009
- Colson, P., Rolain, J. M., and Raoult, D. (2020). Chloroquine for the 2019 novel coronavirus SARS-CoV-2. *Int. J. Antimicrob. Agents.* 55 (3), 105923. doi:10.1016/j.ijantimicag.2020.105923
- Curtis, M. J., Alexander, S., Cirino, G., Docherty, J. R., George, C. H., Gienbycz, M. A., et al. (2018). Experimental design and analysis and their reporting II: updated and simplified guidance for authors and peer reviewers. *Br. J. Pharmacol.* 175 (7), 987–993. doi:10.1111/bph.14153
- Di Cosimo, S., Malfettone, A., Perez-Garcia, J. M., Lombart-Cussac, A., Miceli, R., Curigliano, G., et al. (2020). Immune checkpoint inhibitors: a physiology-driven approach to the treatment of coronavirus disease 2019. *Eur. J. Canc.* 135, 62–65. doi:10.1016/j.ejca.2020.05.026
- Dicko, A., Roh, M. E., Diawara, H., Mahamar, A., Soumare, H. M., Lanke, K., et al. (2018). Efficacy and safety of primaquine and methylene blue for prevention of Plasmodium falciparum transmission in Mali: a phase 2, single-blind, randomised controlled trial. *Lancet Infect. Dis.* 18 (6), 627–639. doi:10.1016/s1473-3099(18)30044-6
- Downing, N. S., Shah, N. D., Aminawung, J. A., Pease, A. M., Zeitoun, J. D., Krumholz, H. M., et al. (2017). Postmarket safety events among novel therapeutics approved by the United States food and drug administration between 2001 and 2010. *J. Am. Med. Assoc.* 317 (18), 1854–1863. doi:10.1001/jama.2017.5150
- Fan, Z., Tian, Y., Chen, Z., Liu, L., Zhou, Q., He, J., et al. (2020). Blocking interaction between SHP2 and PD-1 denotes a novel opportunity for developing PD-1 inhibitors. *EMBO Mol. Med.* 12, e11571. doi:10.15252/emmm.201911571
- Feng, J., Cerniglia, C. E., and Chen, H. (2012). Toxicological significance of azo dye metabolism by human intestinal microbiota. *Front. Biosci.* 4, 568–586. doi:10.2741/400
- Fletcher, S., and Hamilton, A. D. (2006). Targeting protein-protein interactions by rational design: mimicry of protein surfaces. *J. R. Soc. Interface.* 3 (7), 215–233. doi:10.1098/rsif.2006.0115
- Ganesan, L., Margolles-Clark, E., Song, Y., and Buchwald, P. (2011). The food colorant erythrosine is a promiscuous protein-protein interaction inhibitor. *Biochem. Pharmacol.* 81 (6), 810–818. doi:10.1016/j.bcp.2010.12.020
- Gendrot, M., Andreani, J., Dufloy, I., Boxberger, M., Bideau, M. L., Mosnier, J., et al. (2020). Methylene blue inhibits replication of SARS-CoV-2 in vitro. *Int. J. Antimicrob. Agents.* 56, 106202. doi:10.1016/j.ijantimicag.2020.106202
- Golwalkar, D. (2020). Treatment for COVID-19 using methylene blue. Medium. Available at: <https://medium.com/@dr.deepak.golwalkar/treatment-for-covid-19-using-methylene-blue-d23fc5a31a4d> (Accessed April 12, 2020).
- Gordon, D. E., Jang, G. M., Bouhaddou, M., Xu, J., Obernier, K., White, K. M., et al. (2020). A SARS-CoV-2 protein interaction map reveals targets for drug repurposing. *Nature*. 583 (7816), 459–468. doi:10.1038/s41586-020-2286-9
- Guy, R. K., DiPaola, R. S., Romanelli, F., and Dutch, R. E. (2020). Rapid repurposing of drugs for COVID-19. *Science*. 368 (6493), 829–830. doi:10.1126/science.abb9332
- Henry, M., Summa, M., Patrick, L., and Schwartz, L. (2020). A cohort of cancer patients with no reported cases of SARS-CoV-2 infection: the possible preventive role of methylene blue. *Substantia*. 4 (1), 888. doi:10.13128/Substantia-888
- Hershberger, S. J., Lee, S. G., and Chmielewski, J. (2007). Scaffolds for blocking protein-protein interactions. *Curr. Top. Med. Chem.* 7 (10), 928–942. doi:10.2174/156802607780906726
- Lan, J., Ge, J., Yu, J., Shan, S., Zhou, H., Fan, S., et al. (2020). Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2 receptor. *Nature*. 581 (7807), 215–220. doi:10.1038/s41586-020-2180-5
- Leader, B., Baca, Q. J., and Golan, D. E. (2008). Protein therapeutics: a summary and pharmacological classification. *Nat. Rev. Drug Discov.* 7 (1), 21–39. doi:10.1038/nrd2399
- Levine, W. G. (1991). Metabolism of azo dyes: implication for detoxication and activation. *Drug Metab. Rev.* 23 (3-4), 253–309. doi:10.3109/03602539109029761
- Liu, P. P., Blet, A., Smyth, D., and Li, H. (2020). The science underlying COVID-19: implications for the cardiovascular system. *Circulation*. 142, 68–78. doi:10.1161/CIRCULATIONAHA.120.047549
- Lozano, M., Cid, J., and Muller, T. H. (2013). Plasma treated with methylene blue and light: clinical efficacy and safety profile. *Transfus. Med. Rev.* 27 (4), 235–240. doi:10.1016/j.tmr.2013.08.001
- Margolles-Clark, E., Jacques-Silva, M. C., Ganesan, L., Umland, O., Kenyon, N. S., Ricordi, C., et al. (2009a). Suramin inhibits the CD40-CD154 costimulatory interaction: a possible mechanism for immunosuppressive effects. *Biochem. Pharmacol.* 77 (7), 1236–1245. doi:10.1016/j.bcp.2009.01.001
- Margolles-Clark, E., Umland, O., Kenyon, N. S., Ricordi, C., and Buchwald, P. (2009b). Small-molecule costimulatory blockade: organic dye inhibitors of the CD40-CD154 interaction. *J. Mol. Med.* 87 (11), 1133–1143. doi:10.1007/s00109-009-0519-3
- Matheson, N. J., and Lehner, P. J. (2020). How does SARS-CoV-2 cause COVID-19? *Science*. 369 (6503), 510–511. doi:10.1126/science.abc6156
- McKee, D. L., Sternberg, A., Stange, U., Laufer, S., and Naujokat, C. (2020). Candidate drugs against SARS-CoV-2 and COVID-19. *Pharmacol. Res.* 157, 104859. doi:10.1016/j.phrs.2020.104859
- Melby, T., and Westby, M. (2009). Inhibitors of viral entry. *Handb. Exp. Pharmacol.* 189, 177–202. doi:10.1007/978-3-540-79086-0_7
- Michel, M. C., Murphy, T. J., and Motulsky, H. J. (2020). New author guidelines for displaying data and reporting data analysis and statistical methods in experimental biology. *J. Pharmacol. Exp. Therapeut.* 372 (1), 136–147. doi:10.1124/jpet.119.264143
- Milroy, L. G., Grossmann, T. N., Hennig, S., Brunsfeld, L., and Ottmann, C. (2014). Modulators of protein-protein interactions. *Chem. Rev.* 114 (9), 4695–4748. doi:10.1021/cr400698c
- Moore, J. B., and June, C. H. (2020). Cytokine release syndrome in severe COVID-19. *Science*. 368 (6490), 473–474. doi:10.1126/science.abb8925

- Neugebauer, A., Hartmann, R. W., and Klein, C. D. (2007). Prediction of protein-protein interaction inhibitors by chemoinformatics and machine learning methods. *J. Med. Chem.* 50 (19), 4665–4668. doi:10.1021/jm070533j
- Reynès, C., Host, H., Camproux, A. C., Laconde, G., Leroux, F., Mazars, A., et al. (2010). Designing focused chemical libraries enriched in protein-protein interaction inhibitors using machine-learning methods. *PLoS Comput. Biol.* 6 (3), e1000695. doi:10.1371/journal.pcbi.1000695
- Salgado-Benvindo, C., Thaler, M., Tas, A., Ogando, N. S., Bredenbeek, P. J., Ninaber, D. K., et al. (2020). Suramin inhibits SARS-CoV-2 infection in cell culture by interfering with early steps of the replication cycle. *Antimicrob. Agents Chemother.* 64 (8), e00900–e00920. doi:10.1128/AAC.00900-20
- Salje, H., Tran Kiem, C., Lefrancq, N., Courtejoie, N., Bosetti, P., Paireau, J., et al. (2020). Estimating the burden of SARS-CoV-2 in France. *Science.* 369 (6500), 208–211. doi:10.1126/science.abc3517
- Schenk, P., Madl, C., Rezaie-Majid, S., Lehr, S., and Müller, C. (2000). Methylene blue improves the hepatopulmonary syndrome. *Ann. Intern. Med.* 133 (9), 701–706. doi:10.7326/0003-4819-133-9-200011070-00012
- Schirmer, R. H., Adler, H., Pickhardt, M., and Mandelkow, E. (2011). Lest we forget you--methylene blue. *Neurobiol. Aging.* 32 (12), 2325–2416. doi:10.1016/j.neurobiolaging.2010.12.012
- Schneider, C. A., Rasband, W. S., and Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods.* 9 (7), 671–675. doi:10.1038/nmeth.2089
- Scigliano, G., and Scigliano, G. A. (2020). Acute respiratory distress syndrome from Covid-19: a perfect storm from free radicals? Proposal for a new treatment. *Med. Hypotheses.* 144, 110120. doi:10.1016/j.mehy.2020.110120
- Scott, D. E., Bayly, A. R., Abell, C., and Skidmore, J. (2016). Small molecules, big targets: drug discovery faces the protein-protein interaction challenge. *Nat. Rev. Drug Discov.* 15 (8), 533–550. doi:10.1038/nrd.2016.29
- Shang, J., Wan, Y., Luo, C., Ye, G., Geng, Q., Auerbach, A., et al. (2020). Cell entry mechanisms of SARS-CoV-2. *Proc. Natl. Acad. Sci. United States.* 117 (21), 11727–11734. doi:10.1073/pnas.2003138117
- Sivaraman, H., Er, S. Y., Choong, Y. K., Gavor, E., and Sivaraman, J. (2020). Structural basis of the SARS-CoV-2/SARS-CoV receptor binding and small-molecule blockers as potential therapeutics. *Annu. Rev. Pharmacol. Toxicol.* 61, 1. doi:10.1146/annurev-pharmtox-061220-093932
- Song, Y., Margolles-Clark, E., Bayer, A., and Buchwald, P. (2014). Small-molecule modulators of the OX40-OX40 ligand co-stimulatory protein-protein interaction. *Br. J. Pharmacol.* 171 (21), 4955–4969. doi:10.1111/bph.12819
- Sperandio, O., Reynès, C. H., Camproux, A. C., and Villoutreix, B. O. (2010). Rationalizing the chemical space of protein-protein interaction inhibitors. *Drug Discov. Today.* 15 (5-6), 220–229. doi:10.1016/j.drudis.2009.11.007
- Suntharalingam, G., Perry, M. R., Ward, S., Brett, S. J., Castello-Cortes, A., Brunner, M. D., et al. (2006). Cytokine storm in a phase 1 trial of the anti-CD28 monoclonal antibody TGN1412. *N. Engl. J. Med.* 355 (10), 1018–1028. doi:10.1056/nejmoa063842
- Tai, W., He, L., Zhang, X., Pu, J., Voronin, D., Jiang, S., et al. (2020). Characterization of the receptor-binding domain (RBD) of 2019 novel coronavirus: implication for development of RBD protein as a viral attachment inhibitor and vaccine. *Cell. Mol. Immunol.* 17 (6), 613–620. doi:10.1038/s41423-020-0400-4
- Tan, C. W., Chia, W. N., Qin, X., Liu, P., Chen, M. I., Tiu, C., et al. (2020). A SARS-CoV-2 surrogate virus neutralization test based on antibody-mediated blockage of ACE2-spike protein-protein interaction. *Nat. Biotechnol.* 38, 1073–1078. doi:10.1038/s41587-020-0631-z
- Tiwari, V., Beer, J. C., Sankaranarayanan, N. V., Swanson-Mungerson, M., and Desai, U. R. (2020). Discovering small-molecule therapeutics against SARS-CoV-2. *Drug Discov. Today.* 25 (8), 1535–1544. doi:10.1016/j.drudis.2020.06.017
- Tobin, M. J., Laghi, F., and Jubran, A. (2020). Why COVID-19 silent hypoxemia is baffling to physicians. *Am. J. Respir. Crit. Care Med.* 202 (3), 356–360. doi:10.1164/rccm.202006-2157CP
- Vardhana, S. A., and Wolchok, J. D. (2020). The many faces of the anti-COVID immune response. *J. Exp. Med.* 217 (6), e20200678. doi:10.1084/jem.20200678
- Wadman, M. (2006). London's disastrous drug trial has serious side effects for research. *Nature.* 440 (7083), 388–389. doi:10.1038/440388a
- Walter-Sack, I., Rengelshausen, J., Oberwittler, H., Burhenne, J., Mueller, O., Meissner, P., et al. (2009). High absolute bioavailability of methylene blue given as an aqueous oral formulation. *Eur. J. Clin. Pharmacol.* 65 (2), 179–189. doi:10.1007/s00228-008-0563-x
- Xiu, S., Dick, A., Ju, H., Mirzaie, S., Abdi, F., Cocklin, S., et al. (2020). Inhibitors of SARS-CoV-2 entry: current and future opportunities. *J. Med. Chem.* 63 (21), 12256–12274. doi:10.1021/acs.jmedchem.0c00502
- Ye, Q., Wang, B., and Mao, J. (2020). The pathogenesis and treatment of the 'cytokine storm' in COVID-19. *J. Infect.* 80 (6), 607–613. doi:10.1016/j.jinf.2020.03.037

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Bojadzic, Alcazar and Buchwald. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.