



OPEN ACCESS

Edited by:

Timothy E Albertson,
UC Davis Medical Center,
United States

Reviewed by:

Arnab K Chatterjee,
Calibr at Scripps Research,
United States
Darrell O. Rieke,
Massachusetts Institute of
Technology, United States

***Correspondence:**

Md. Moklesur Rahman Sarker
moklesur2002@yahoo.com
dr.moklesur2014@gmail.com
prof.moklesur@sub.edu.bd
Isa Naina Mohamed
isanaina@ppukm.ukm.edu.my

†ORCID ID

Safaet Alam
orcid.org/0000-0002-1831-2278
Taslima Binte Kamal
orcid.org/0000-0003-3687-1388
Md. Moklesur Rahman Sarker
orcid.org/0000-0001-9795-0608
Jin-Rong Zhou
orcid.org/0000-0002-6745-9495
S. M. Abdur Rahman
orcid.org/0000-0002-9963-8885
Isa Naina Mohamed
orcid.org/0000-0001-8891-2423

†These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Respiratory Pharmacology,
a section of the journal
Frontiers in Pharmacology

Received: 27 January 2021

Accepted: 28 April 2021

Published: 14 June 2021

Citation:

Alam S, Kamal TB, Sarker MMR,
Zhou J-R, Rahman SMA and
Mohamed IN (2021) Therapeutic
Effectiveness and Safety of
Repurposing Drugs for the Treatment
of COVID-19: Position Standing
in 2021.
Front. Pharmacol. 12:659577.
doi: 10.3389/fphar.2021.659577

Therapeutic Effectiveness and Safety of Repurposing Drugs for the Treatment of COVID-19: Position Standing in 2021

Safaet Alam^{1†}, Taslima Binte Kamal^{1†}, Md. Moklesur Rahman Sarker^{1,2*†}, Jin-Rong Zhou^{3†}, S. M. Abdur Rahman^{4†} and Isa Naina Mohamed^{5*†}

¹Department of Pharmacy, State University of Bangladesh, Dhaka, Bangladesh, ²Pharmacology and Toxicology Research Division, Health Med Science Research Limited, Dhaka, Bangladesh, ³Nutrition/Metabolism Laboratory, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, United States, ⁴Department of Clinical Pharmacy and Pharmacology, Faculty of Pharmacy, University of Dhaka, Dhaka, Bangladesh, ⁵Pharmacology Department, Medical Faculty, Universiti Kebangsaan Malaysia (The National University of Malaysia), Kuala Lumpur, Malaysia

COVID-19, transmitted by SARS-CoV-2, is one of the most serious pandemic situations in the history of mankind, and has already infected a huge population across the globe. This horrendously contagious viral outbreak was first identified in China and within a very short time it affected the world's health, transport, economic, and academic sectors. Despite the recent approval of a few anti-COVID-19 vaccines, their unavailability and insufficiency along with the lack of other potential therapeutic options are continuing to worsen the situation, with valuable lives continuing to be lost. In this situation, researchers across the globe are focusing on repurposing prospective drugs and prophylaxis such as favipiravir, remdesivir, chloroquine, hydroxychloroquine, ivermectin, lopinavir-ritonavir, azithromycin, doxycycline, ACEIs/ARBs, rivaroxaban, and protease inhibitors, which were preliminarily based on *in vitro* and *in vivo* pharmacological and toxicological study reports followed by clinical applications. Based on available preliminary data derived from limited clinical trials, the US National Institute of Health (NIH) and USFDA also recommended a few drugs to be repurposed i.e., hydroxychloroquine, remdesivir, and favipiravir. However, World Health Organization later recommended against the use of chloroquine, hydroxychloroquine, remdesivir, and lopinavir/ritonavir in the treatment of COVID-19 infections. Combining basic knowledge of viral pathogenesis and pharmacodynamics of drug molecules as well as *in silico* approaches, many drug candidates have been investigated in clinical trials, some of which have been proven to be partially effective against COVID-19, and many of the other drugs are currently under extensive screening. The repurposing of prospective drug candidates from different stages of evaluation can be a handy wellspring in COVID-19 management and treatment along with approved anti-COVID-19 vaccines. This review article combined the information from completed clinical trials, case series, cohort studies, meta-analyses, and retrospective studies to focus on the current status of repurposing drugs in 2021.

Keywords: COVID-19, repurposing drugs, favipiravir, remdesivir, doxycycline, hydroxychloroquine, ivermectin, rivaroxaban

INTRODUCTION

Coronaviruses, genus of the Coronaviridae family, have been the most tragic pathogen of the 21st century, causing three major epidemics since 2002. The recent coronavirus outbreak of 2019 is the most dreadful of all the previous ones, halting every aspect of human life worldwide. The World Health Organization (WHO) recognized coronavirus disease-2019 (COVID-19) as a pandemic on March 11, 2020 (Singh et al., 2020). According to WHO, a novel coronavirus (severe acute respiratory syndrome coronavirus 2, SARS-CoV-2) case, found in China, was first reported to the WHO Country Office on December 31, 2019. Up to January 25, 2021, 98,794,942 confirmed cases and 2,124,193 deaths had been reported globally due to COVID-19 (WHO Coronavirus Disease (COVID-19) Dashboard, 2020). SARS-CoV-2 contains a receptor binding domain on its surface's crown shaped spikes protein which helps it to bind to human angiotensin-converting enzyme 2 (ACE2), promoting its endocytosis into human lung cells. After entry, this virus continues its replication in lung cells by synthesizing and assembling its own protein using host cells' protein synthesis machinery (Rothan and Byrareddy, 2020).

Although a few vaccines against SARS-CoV-2 infections have recently been approved by the regulatory authorities and been used, there are no other proven and/or potential therapeutic options in hand to treat this pandemic. To worsen the issue, allocation and production in sufficient quantities and the affordability and deployment of these vaccines are also notable challenges faced by the policy makers and developers (Folegatti et al., 2020; Polack et al., 2020; Logunov et al., 2021; Wouters et al., 2021). Thus, as the search for new effective therapeutic options are still being conducted and will need more time and resources, scientists are exploring the possibility of the usage of existing drugs that may be therapeutically effective for the treatment of COVID-19 along with the discovery and availability of vaccines. Repurposing of a drug indicates the use of currently existing drugs for new therapeutic purposes rather than its primary and main indications (Khan et al., 2020). There have been several investigations on drug repurposing for COVID-19 recently and clinical trials are still being conducted on several drugs. The current review aims to present a clear idea on the mechanism of actions and outcomes from completed clinical trials along with the data from case series, retrospective studies, cohort studies, and meta-analyses on the efficacy and safety of prospective repurposing drugs in the treatment of COVID-19 infections.

Article Search Strategy

To summarize and compile the desired findings related to repurposing drugs in COVID-19 treatment, a literature search was conducted using Web of Science, PubMed, ScienceDirect, Scopus, Wiley Online Library, and Google Scholar databases. In addition, clinical trial reports and other relevant information were also checked critically from authentic sources including WHO, USFDA, and ClinicalTrials.gov websites. The keywords used during the searches included 'COVID-19', 'COVID', 'Repurposing drugs', 'Anti-COVID-19 drugs', 'Coronavirus' and 'Clinical trial', 'Case study', 'Retrospective study', 'Cohort study', and 'Meta-analysis'. Only the completed studies on clinical


trials, case series, cohort studies, meta-analyses and retrospective data (up to 9th April, 2021) from aforementioned authentic databases were included in this review, while incomplete studies of any of the above categories were excluded. Of the 412 identified papers and reports, after discarding titles and abstracts mismatched with the inclusion criteria, 176 unique sources were identified to be finally included in the review.

Remdesivir

Remdesivir, developed by Gilead Sciences, was considered as a potential treatment option for RNA viruses like EBOV (Ebola Virus), MERS (Middle East respiratory syndrome), and SARS (Severe acute respiratory syndrome), all of which could cause a global pandemic. In 2014, during the ebola virus outbreak, clinical evaluation for remdesivir started. But being inferior to antibody-based therapeutics for the treatment of EBOV and having one serious adverse event (hypotension) along with increased plasma levels of creatinine and aspartate aminotransferase, its evaluation study against Ebola virus was terminated (Eastman et al., 2020). A randomized controlled trial of four investigational therapies (triple monoclonal antibody ZMapp, remdesivir, the single monoclonal antibody MAb114, or the triple monoclonal antibody REGN-EB3) was conducted for ebola virus disease (EVD) after the EVD outbreak of 2018 in the Democratic Republic of Congo. Data from this trial indicated that remdesivir was inferior to the other therapies and the highest mortality was reported in the remdesivir group than the other groups. Random assignment of patients to the remdesivir group was recommended to be stopped depending on efficacy and mortality analysis of patients. The median time of the first negative result for EBOV nucleoprotein was also higher in the remdesivir group (28 days) compared to the other groups (Mulangu et al., 2019). With the COVID-19 outbreak, several *in-vitro* and *in-vivo* animal studies were conducted to assess the effectiveness of remdesivir against SARS-CoV-2 and most of the studies showed positive results. A recent *in vitro* study in infected Vero E6 involving qRT-PCR quantification of viral copy number demonstrated an IC50 of 770 nM (Wang et al., 2020).

Supportive *in-vitro* data for the use of remdesivir against SARS-CoV-2 and a lack of alternative treatment options led to the development of clinical trials with remdesivir. While clinical trials with remdesivir were being developed in China, the first case of a COVID-19 positive patient was reported in the USA on January 20, 2020. The patient was given IV remdesivir infusion and showed improved clinical condition with no identified adverse event. However, the patient also received Paracetamol, Ibuprofen, Guaifenesin, Vancomycin, and Cefepime along with remdesivir (Holshue et al., 2020). A benefit risk assessment study for remdesivir showed that one clinical trial reported non-significant benefit of recovery time (21 vs. 23 days) while another study reported non-significant reduction of mortality risk (8 vs. 12%) compared to placebo (Davies et al., 2020).

Remdesivir is an investigational nucleotide analogue and as its triphosphate form resembles Adenosine triphosphate it can compete with it to act as a substrate for various viral RNA polymerase inhibiting RNA synthesis. Remdesivir triphosphate form adds to the growing RNA chain more efficiently than ATP, causing chain termination at 3 nucleotides downstream. This




delayed chain termination mechanism can contribute to the effectiveness of remdesivir against all strain of coronaviruses including SARS-CoV and SARS-CoV-2 (Saha et al., 2020). The anti-COVID-19 action of remdesivir is displayed in **Figure 1**.

Remdesivir, being an investigational drug, received the first FDA authorization for emergency use in treating SARS-CoV-2 only in hospital or healthcare settings capable of providing acute care. Further research is required to establish its role in managing COVID-19 patients (Ridwan, 2020).

A retrospective cohort study in Korea involving 86 severe COVID-19 patients hospitalized between June 1 and July 31, 2020, suggested better clinical and virologic effectiveness of remdesivir compared to supportive care in the case of mechanical ventilator requirement (22.9 vs. 44.7%, $p = 0.032$), mechanical ventilator duration (average, 1.97 vs. 5.37 days; $p = 0.017$), and viral load reduction (increase of cycle threshold (Ct) values: average, 10.19 vs. 5.36; $p = 0.007$ and the slope of Ct value increase: average, 5.10 vs. 2.68; $p = 0.007$) (Joo et al., 2021). Another retrospective analysis demonstrated a well-tolerated safety profile of remdesivir and 84% improvement rate of the drug in COVID-19 patients (Gupte et al., 2021). On the contrary, a large observational study in Japan concluded that remdesivir may not have any benefit on hospitalized COVID-19 patients' clinical outcomes and reductions in need of invasive/non-invasive respiratory support. Results from the study showed that the remdesivir case group and non-remdesivir control group had no significant difference in fatality risk (12.2 vs. 13.3%; $p = 1.0$), risk of invasive mechanical ventilation or extracorporeal membrane oxygenation (IMV/ECMO 5.4 vs. 4.6%; $p = 0.757$), length of intensive care unit stay (6 vs. 6 days; $p = 0.473$), or length of IMV (13 vs. 5 days; $p = 0.509$), while longer length of stay (LoS) in healthcare facilities was

observed in the case group than in the control group (14 vs. 11 days; $p < 0.001$) (Tsuzuki et al., 2021). Again, a retrospective, cohort study involving 142 COVID-19 patients admitted to a large tertiary center in Israel showed no nasopharyngeal viral load changes with remdesivir (Goldberg et al., 2021).

The Adaptive COVID-19 Treatment Trial (ACTT-1) involving 1,062 patients and Solidarity involving 6,838 patients are the two largest trials among all the published randomized controlled trials (RCTs) aiming to evaluate remdesivir as a COVID-19 treatment option. ACTT-1 found reduced recovery time (shortened from 15 to 10 days) but no change in mortality (hazard ratio 0.73, 95% confidence interval [CI] 0.52–1.03) with remdesivir and Solidarity also showed no mortality benefit for the drug (rate ratio 0.95, 95% CI 0.81–1.11). Both the trials demonstrated a lack of effectiveness of remdesivir in critically ill patients (Wu and Morris, 2021). A meta-analysis of four randomized controlled trials (RCTs) of remdesivir in moderate or severe COVID-19 patients showed increased recovery rate and hospital discharge with no effect on clinical improvement and mortality (Al-Abdoh et al., 2021). Another meta-analysis involving five RCT's concluded that remdesivir may elevate the recovery percentage and slightly decrease ventilation requirement in hospitalized COVID-19 patients but has no effect in mortality. It also suggested the benefit of a 5-days course over a 10-days course of remdesivir (Kaka et al., 2021). Six published case-series and cohort series studies on the use of remdesivir against COVID-19 reported potential effectiveness of the drug but most of the study results are limited by their small group size, short follow-up duration, lack of potential data, scarcity of patient's information treated at baseline, and lack of a randomized control group (Saqrane et al., 2021).



Moreover, a few more studies are going on to determine the safety, antiviral efficacy, pharmacokinetic property, and tolerability of remdesivir alone and in combination with other agents like tocilizumab, merimepodib, and convalescent plasma therapy in cases of severe and moderate COVID-19 patients in the United States, Bangladesh, Nepal, China, and Canada (Anonymous, 2020a). However, with a lack of adequate evidence supporting remdesivir use in COVID-19, WHO recently deprecated its use regardless of hospitalized patients' severity of illness (WHO, 2020).

Doxycycline

Doxycycline (1967) is a semisynthetic second-generation class of tetracyclines with a broad spectrum of antimicrobial activity. It was developed by chemical modification of basic chlortetracycline structure and identified and named by mycologist B. M. Duggar in 1944 (Joshi and Miller, 1997). Having antiviral, immunomodulatory, and anti-inflammatory properties, doxycycline and other tetracycline antibiotics could be potential options for treating COVID-19 patients. An *in vitro* antiviral activity study of doxycycline involving VERO E6 cell line showed positive results against SARS-CoV-2 with median effective concentration (EC₅₀) of $5.6 \pm 0.4 \mu\text{M}$. For prophylaxis, doxycycline could be used in combination with hydroxychloroquine or chloroquine (Gendrot et al., 2020).

Several mechanisms have been proposed for the actions of doxycycline against SARS-CoV-2. Effectiveness of tetracyclines (e.g., tetracycline, doxycycline, and minocycline) against

COVID-19 infection can be attributed to their higher lipophilicity with higher lung tissue penetration ability facilitating their role in viral replication inhibition in the lungs. Again, tetracyclines can form chelate with zinc compounds on matrix metalloproteinases (MMPs) of the host while coronaviruses rely heavily on MMPs of the host for cell adhesion, infiltration, survival, and replication, thus inhibiting the virus' ability to replicate within the host. They might also act by inhibiting positive-sense single-stranded RNA replication of COVID-19. Again, due to their anti-inflammatory mechanism, tetracyclines are used to treat different inflammatory conditions. They can decrease the levels of inflammatory cytokines, such as tumor necrosis factor- α , interleukin (IL)-1 β , and IL-6, which are elevated when lung tissue is infected with severe acute respiratory syndrome-associated coronavirus (SARS-CoV) deteriorating the infection itself (Sodhi and Etminan, 2020). The anti-COVID-19 action of doxycycline is displayed in **Figure 2**.

Doxycycline (a tetracycline antibiotic), being a non-toxic mitochondrial biogenesis and cellular respiration inhibitor and having various pleiotropic features, can attenuate the replication of most viruses which need the energy of aerobic glycolysis to replicate and use mitochondria as cellular powerhouses. It is suggested that doxycycline in combination with Vitamin C shows more effectiveness against COVID-19 (Szolnoky, 2020). Again, Sargiacomo et al. suggested that senolytic drugs such as doxycycline can be used for SARS-CoV-2 infection to prevent fibrotic transformation. SARS-CoV-2 can cause stormy inflammation and subsequent fibrosis after binding and

replicating preferably with senescent and chronologically aged lung cells (Sargiacomo et al., 2020).

In New York, 89 high-risk COVID-19 patients (diagnosed from March 18 to May 13, 2020) received early treatment with doxycycline in long-term care facilities and demonstrated early clinical recovery, reduced hospitalization, and reduced mortality. Among 89 patients, 76 (85%) were clinically recovered, three patients (3%) showed deterioration of clinical condition requiring transfer to hospital, and 10 patients (11%) died (Alam et al., 2020). A prospective case-series study involving 21 COVID-19 patients from Iran from September 14, 2020 to September 28, 2020 suggested the potential effectiveness of doxycycline in COVID-19 treatment (Meybody et al., 2021). Recovery of four high risk COVID-19 patients with comorbid pulmonary diseases who were treated with doxycycline was reported (Narendrakumar et al., 2021).

Doxycycline shows more effectiveness against SARS-CoV-2 when used in combination with other drugs like hydroxychloroquine, ivermectin, and dapson. More studies are required to establish its effectiveness in proposed combination therapies for COVID-19 (Farouk and Salman, 2020; Malek et al., 2020). At present, various clinical trials are being conducted with doxycycline alone or in combination with other drugs for COVID-19 treatment. An experimental trial with combination of hydroxychloroquine and doxycycline involving 54 high risk COVID-19 patients demonstrated higher clinical recovery, decreased hospital stay, and reduced mortality (Narendrakumar et al., 2021). A recent double-blind and randomized interventional trial of doxycycline in combination with ivermectin was conducted with 400 participants. Early clinical improvement was reported in 60.7% of patients (111 out of 183 patients) with doxycycline and ivermectin combination compared to 44.4% (80 out of 180) with placebo (NCT04523831, 2020). More studies are still in progress in Iraq, Egypt, and Bangladesh to assess the efficacy and safety of this combination for COVID-19 treatment (Anonymous, 2020b).


Azithromycin

Azithromycin, a macrolide derivative and the first of the fifteen-membered ring azalide class of antimicrobials, shows a broad spectrum of activity against various gram-positive and gram-negative bacteria and other atypical pathogens (Ballou and Amsden, 1992). They act by inhibiting bacterial growth through interference with their protein synthesis. Azithromycin showed *in-vitro* activity against Ebola and Zika viruses and prevented severe viral infections of the respiratory tract. Previously they have been used as an adjunctive therapy to treat viral respiratory tract infections due to their anti-inflammatory and immunomodulatory activities in addition to extended antibacterial coverage (Wu et al., 2020). With no available definite treatment option for COVID-19, azithromycin has undergone several studies recently to find its effectiveness against SARS-CoV-2. An *in-vitro* screening with VeroE6 (ATCC CRL-1586) and Caco-2 (ATCC HTB-37) cells reveals Azithromycin to be a potential inhibitor of SARS-CoV-2 replication with EC_{50} (50% effective concentration) = 2.12 μ M and EC_{90} (90% effective concentration) = 8.65 μ M. (Touret et al., 2020).

There is no known specific mechanism; instead, several mechanisms have been proposed for the probable antiviral activity of Azithromycin. Any of these mechanisms could be responsible for its activity against SARS-CoV-2. For example, being a weak base, azithromycin can inhibit the uncoating of coronaviruses as the uncoating of enveloped viruses requires an acidic environment. Again, preferential accumulation of this drug in lysosomes and endosomal vesicles can elevate pH and limit replication of the virus by blocking endocytosis. Another possible mechanism can be the amplification of host's interferon (IFN) pathway-mediated antiviral responses, resulting in a reduction of viral replication. A quantum mechanical model proposes that azithromycin's interference in the binding interaction of SARS-CoV-2 spike protein and host receptor ACE2 (angiotensin converting enzyme-2) protein can be responsible for limited viral entry (Damle et al., 2020). Azithromycin has also been suggested to have senolytic activity used to inhibit viral replication and IL-6 production (Sargiacomo et al., 2020). The anti-COVID-19 action of Azithromycin is displayed in **Figure 3**.

Currently, various clinical trials are being conducted to evaluate the efficacy of Azithromycin in combination with hydroxychloroquine for COVID-19 treatment. According to a study, azithromycin and hydroxychloroquine combination therapy was encouraged in the treatment of early symptomatic and high-risk outpatients (Esper et al., 2020). Several other preliminary trials also reported positive results like the one performed by Pfizer in France and another reported by Gautret et al. (2021) which previously suggested the use of combination treatment of azithromycin and hydroxychloroquine for COVID-19 (Gautret et al., 2021; Wu et al., 2020). According to another retrospective observational study, azithromycin (introduced after 6–8 h from being diagnosed) could notably improve the hospital stay duration and requirement of respiratory support in hospital days (Ishaqui et al., 2020). Despite having no clear idea about the optimal dose of azithromycin against SARS-CoV-2, in accordance with IDSA guidelines and RECOVERY trial, 500 mg azithromycin once daily (OD) for 5 days is recommended in the case of severe patients. Again, based on previous studies, azithromycin was only suggested to early phase patients with COVID-19 infection as in later phases it failed to deliver any beneficial impacts on clinical status (Echeverría-Esnal et al., 2021).

On the contrary, one of the biggest randomized controlled clinical trials conducted on hospitalized patients with mild to moderate COVID-19 infection involved the administration of hydroxychloroquine monotherapy azithromycin and hydroxychloroquine cotherapy, which did not alleviate patient status at 15 days compared to standard care (Arshad et al., 2020; Echeverría-Esnal et al., 2021). Another experiment also failed to demonstrate any evidence of rapid antiviral activity in severe COVID-19 patients with hydroxychloroquine and azithromycin combination therapy (Molina et al., 2020). Moreover, this combination is reported to prolong QT and lacks safety data for patients with arrhythmia or renal or hepatic impairment. So, robust clinical data are required to support the efficacy and safety of this combined therapy (Machiels et al., 2020). Recently an



open-label and randomized phase-3 clinical trial of hydroxychloroquine and azithromycin for COVID-19 treatment in pregnant women has been withdrawn. In another UK-based, open label, multi arm, and randomized clinical trial, the routine use of azithromycin failed to display promising activities to shorten the recovery time or risk of hospitalization of COVID-19 infected individuals. The trial also reported inappropriate and excessive use of azithromycin during the COVID-19 pandemic which ultimately leads to antimicrobial resistance (Butler et al., 2021). A few other clinical studies also suggested the same inefficacy of azithromycin or combination of azithromycin with hydroxychloroquine in terms of virologic clearance and other poor clinical outcomes (Molina et al., 2020; Echeverría-Esnal et al., 2021). Besides, the administration of hydroxychloroquine and azithromycin in hospitalized patients has been linked to an increased risk of cardiac adverse events. More specifically, such a combination has been associated with an increased risk of QTc prolongation, ventricular arrhythmia, TdP (0.4%), atrial fibrillation, atrioventricular block, and cardiac failure (Echeverría-Esnal et al., 2021).


Interestingly, some very recent studies revealed that in combination therapy, hydroxychloroquine might cause cardiac toxicity not azithromycin (Echeverría-Esnal et al., 2021). In addition, in the only randomized-controlled clinical trial conducted to date in hospitalized patients, azithromycin

monotherapy did not confer a higher risk of adverse effects like hydroxychloroquine or combination therapy with hydroxychloroquine (Arshad et al., 2020; Echeverría-Esnal et al., 2021). Thus, though a cautious risk-benefit ratio and follow up of ADRs should be monitored, from a safety perspective azithromycin could be being studied for its use against SARS-CoV-2 infection (Echeverría-Esnal et al., 2021).

Lopinavir-Ritonavir Combination

Lopinavir-Ritonavir combination (Kaletra), approved by the FDA in 2000, was developed by Abbott Laboratories, USA to treat HIV patients. Both drugs are retroviral protease inhibitors but are given in combination to enhance Lopinavir exposure within the body. Lopinavir has a short half-life and so was co-formulated with ritonavir which can minimize extensive biotransformation of Lopinavir inhibiting the metabolizing enzyme cytochrome P450 (Chandwani and Shuter, 2008). With positive results from an *in-vitro* SARS-CoV inhibitory study, lopinavir/Ritonavir was first used in a clinical study with 152 patients during the 2003 SARS outbreak. This study reported weak antiviral activity of lopinavir-ritonavir and suggested further studies (Chu et al., 2004).

Plausible effectiveness of lopinavir/ritonavir against SARS-Cov-2 might be due to the inhibition of viral cysteine protease. Several studies involving the use of lopinavir-ritonavir in SARS-Cov-2 patients have been conducted. These



studies demonstrated positive results for old patients and pediatric patients and reported reduced hospital stay time with quick recovery. But in a recent study this combination showed adverse effects with no significant improvement in clinical recovery time and mortality rate (Singh et al., 2020). Anti-COVID-19 action of lopinavir-ritonavir combination is displayed in Figure 4.

In Korea, lopinavir-ritonavir therapy has been proposed for high-risk elderly COVID-19 patients at an early stage depending on a positive report from lopinavir/ritonavir application for treating COVID-19 pneumonia in a Korean patient (Lim et al., 2020). A retrospective case-series study involving 19 severe COVID-19 patients demonstrated 94.74% clinical improvement rate (Luo et al., 2021). A randomized controlled and open-label trial conducted by Cao et al. (2020) to investigate the specific efficacy of lopinavir/ritonavir administration showed no significant improvement of clinical recovery time compared to standard care (Cao et al., 2020). Though there is a lack of adequate clinical data to confirm the effectiveness of this combination, Chinese guidelines suggests the combination for critically ill elderly COVID-19 patients. Moreover, several case studies reported enhanced antiviral activity when lopinavir-ritonavir is administered with other drugs like ribavirin and interferon alfa (Khan et al., 2020). Triple combination of lopinavir/ritonavir, ribavirin, and interferon- β was compared with lopinavir/ritonavir in a recent multicenter, prospective, open-label, and randomized phase-2 trial. The study

demonstrated significantly better clinical response and reduced median hospitalization (9 days against 14.5 days, $p = 0.016$) in the treatment group in comparison to the control group. In the study, 86 hospitalized patients who were early assigned (within a median of 5 days) to a triple combination of lopinavir/ritonavir (400/100 mg every 12 h), ribavirin (400 mg every 12 h), and three doses of 8 million international units of interferon- β 1b on alternate days for 14 days, had a significantly reduced median time of hospital stay [(7 days against 12 days, $p = 0.0010$) than the control group of 41 patients treated only with lopinavir/ritonavir for 14 days (Meini et al., 2020). Several clinical trials are ongoing with lopinavir-ritonavir alone and in combination with other drugs like Favipiravir to reach a conclusion regarding their effectiveness and safety for COVID-19 treatment. This combination is also being compared with other agents like hydroxychloroquine sulfate and darunavir/cobicistat in recent trials (Anonymous, 2020c). However, a retrospective case-series study raised concerns about the safety of lopinavir-ritonavir use in COVID-19 patients based on the possibility of causing serious adverse events even before reaching the presumed therapeutic concentration for COVID-19 treatment. Moreover, WHO reported the recommendation of Solidarity trial regarding immediate discontinuation of this combination in COVID-19 patients due to ineffectiveness and safety concerns on July 4, 2020 (Lepage et al., 2021). Again, a large retrospective, multicenter study involving 8,553 COVID-19 patients found no overall advantage of lopinavir-ritonavir administration in early

COVID-19 patients (Lora-Tamayo et al., 2021). Moreover, based on data extracted from 17 different studies, the use of lopinavir-ritonavir was strongly discouraged by WHO due to their dubiousness on the rate of reducing mortality and mechanical ventilation along with the probability of increasing diarrhea (Lamontagne et al., 2020). Besides, the time of clinical improvement is also uncertain when lopinavir-ritonavir was used against COVID-19 infection (Lamontagne et al., 2020).

Chloroquine

Chloroquine was first extracted from the cinchona plant by Bayer Laboratories in 1934 (Savarino et al., 2003; Singh et al., 2020). It has been used clinically for more than 50 years all over the world. Initially it was synthesized and applied as an anti-plasmodial drug, but it also can act as an extended spectrum antiviral agent (Savarino et al., 2003; Paton et al., 2011). Chloroquine and hydroxychloroquine are established drug moieties with great similarities in their structures (Rynes, 1997) though compared to chloroquine, hydroxychloroquine exhibits less toxicity due to the presence of a hydroxyl group (Wu et al., 2020). Chloroquine can restrain SARS-CoV-2 from entering into the host cell. It can also effectively inhibit the virus and host cell fusion by altering glycosylation of ACE2 receptor and its binding capacity with spike protein. There is also evidence of chloroquine's efficacy during early phase of COVID-19 attack by down regulating expression of ACE2 and its activity (Savarino et al., 2006; Liu et al., 2020; Wang et al., 2020). The anti-COVID-19 action of chloroquine is displayed in **Figure 4**.

On the contrary, according to some research, chloroquine, evaluated against several viral infections, including human immunodeficiency virus (HIV), MERS-CoV, SARS-CoV, chikungunya, and dengue (Savarino et al., 2003; Keyaerts et al., 2009; Colson et al., 2020), failed to show any benevolent actions in animal models or clinical trials (Touret and de Lamballerie, 2020). According to research, during a randomized double blind placebo controlled trial, chloroquine could not deliver any promising activity against viral infections such as influenza virus (Paton et al., 2011).

Researchers in China recommended that during COVID-19 infections 500 mg chloroquine should be provided twice a day for ten days continuously (Jie et al., 2020; Smith et al., 2020). The China National Center for Biotechnology Development also documented chloroquine as one of the top three anti-COVID-19 drug candidates (Khan et al., 2020). Primary evidence from the Chinese authorities also stated that almost 100 COVID-19 affected patients experienced quicker recovery from fever, alleviated lung computed tomography (CT) images, and no pronounced side effects and recommended chloroquine in the COVID-19 treatment protocols (Gao J. et al., 2020; Jie et al., 2020). This made chloroquine one of the first line treatments throughout China and different countries, though some prevalent side effects, including cardiomyopathy and macular retinopathy, cannot be nullified (Cubero et al., 1993). Almost 10 clinical studies have been conducted to determine the efficacy and safety of chloroquine to ascertain improved therapeutic decision (Harrison, 2020; Smith et al., 2020). But some reports also stated that chloroquine in high doses (600 mg twice daily for

10 days or total dose of 12 g) can trigger cardiac arrest and thus should not be prescribed for COVID-19 infection (Borba et al., 2020). Despite being a safe choice, chloroquine, a drug with a narrow therapeutic index, can trigger retinopathy, methemoglobinemia, gastrointestinal anomalies, tinnitus, and cardiac conditions including arrhythmias and cardiomyopathy (Frisk-Holmberg et al., 1983; Waldrop et al., 2020). As the safety and efficacy profiles of chloroquine and hydroxychloroquine is still under question, health practitioners should make a critical decision considering risk vs. benefit ratio (Juurlink, 2020).

Several clinical trials were conducted to evaluate the efficacy of chloroquine against COVID-19 infection. Up to April 2020, more or less 80 clinical studies had been conducted on chloroquine and/or hydroxychloroquine and even sometimes with other drugs, to assess the efficacy of these agents against the novel corona virus strain. A few trials revealed positive outcomes (Femere and Aronson, 2020). Moreover, several trials are underway to compare chloroquine with other drugs (like hydroxychloroquine or telemedicine) or to compare chloroquine combination drugs (chloroquine + losartan) with chloroquine alone (Anonymous, 2020d). Based on a trial conducted in 2020, the candidacy of chloroquine was proposed by alleviating disease symptoms and supporting virus negative conversion and shortening of the disease period (Gao et al., 2020). But after preliminary hype, the safety and efficacy profile of chloroquine or hydroxychloroquine has imposed a severe question mark on their repurposing use. Use of chloroquine/hydroxychloroquine can induce cardiac arrest and malignant arrhythmias. Another meta-analysis conducted on COVID-19 patients also revealed risk of QTc prolongation followed by discontinuation of chloroquine/hydroxychloroquine (Kashour et al., 2021). The increasing number of ADR incidences resulted from chloroquine/hydroxychloroquine revealed its failure in COVID-19 infection. The number of ADRs (from FAERS database) due to chloroquine and hydroxychloroquine is more than double in previous years (in 2020 $n = 11,493$ and 89,607 ADRs, in 2019 $n = 5,131$ and 37,559 ADRs and in 2018 $n = 4,681$ and 25,035 ADRs) which is possibly attributable to excessive use of these drugs against SARS-CoV-2 infection. Considering these incidences and major clinical trials, the USFDA suspended emergency use authorization for these drugs on June 15, 2020 (Bull-Otterson et al., 2020). Thus, with time, the once promising drug candidates chloroquine and hydroxychloroquine failed to exert their safety and efficacy profiles against COVID-19 infections and dropped out from the race of being repurposed.

Hydroxychloroquine

Being an analog of chloroquine, hydroxychloroquine also shows immunomodulatory activities and is used to treat malaria, rheumatoid arthritis, and systemic lupus erythematosus (SLE) (Savarino et al., 2003; Wu et al., 2020). As a safer analogue of chloroquine, an FDA approved drug, hydroxychloroquine has been prescribed for the last 60 years and is also the most abundantly prescribed antimalarial drug (Shippey et al., 2018).

Despite not having an evidently elucidated mechanism of action of hydroxychloroquine against coronavirus, several mechanisms to irradiate its antiviral activity are already

projected. Reports claim that hydroxychloroquine can effectively inhibit SARS-CoV-2 infection during *in vitro* evaluation (Colson et al., 2020; Liu et al., 2020; Zhou P. et al., 2020). Being a weakly basic drug, hydroxychloroquine increases the pH within cells and at the cellular membrane and consequently arrests the capability of the virus to fuse onto the cell membrane and invade the host cell. Other anticipated mechanisms of hydroxychloroquine include cessation of DNA and RNA synthesis along with immunomodulating and anti-inflammatory properties (Reilly, 2020). Hydroxychloroquine can also arrest glycosylation of viral proteins, virus assembly, new virus particle transport, virus release, and other processes to attain its antiviral activity (Vincent et al., 2005). The anti-COVID-19 action of hydroxychloroquine is displayed in **Figure 4**.

Hydroxychloroquine has been given priority in COVID-19 management in various countries due to its abundant availability and *in vitro* activity against COVID-19 (Khan et al., 2020). Clinical trials in mild, moderate, and severe cases ensure that hydroxychloroquine can eliminate viral nasopharyngeal carriage of COVID-19 within a week, though at present no report from Randomized Controlled Trials (RCTs) is available (Gautret et al., 2020; Khan et al., 2020). Despite not establishing a clear dosage regimen, a few USA experts recommend variable doses including 400 mg BID on day one and then continuation for 5 days, 400 mg BID on day one and then continuation of 200 mg BID for 4 days, and 600 mg BID on day one and then continuation of 400 mg daily on days 2–5 (Khan et al., 2020). The China International Exchange and Promotive Association for Medical and Health Care (CPAM) has given a guideline of hydroxychloroquine use (400 mg once daily) when chloroquine cannot be availed (Smith et al., 2020).

A recent study confirmed 8 out of 10 patients' positive status for SARS-CoV-2 PCR on days 5–6 after treating with hydroxychloroquine and azithromycin (Molina et al., 2020). Another study which has not yet been peer reviewed reported clinical improvement in 80.6% cases (25 of 31 patients) when treated with hydroxychloroquine whereas there was a 54.8% (17 of 31 patients) improvement rate in control group who were treated with antivirals other than hydroxychloroquine (Chen Z. et al., 2020), though use of both chloroquine and hydroxychloroquine should be monitored strictly including checking of patients' QT measurement. Thus, off-label use should be restricted (Giudicessi et al., 2020). A few patients should be tested for glucose-6-phosphate dehydrogenase deficiency as use of these drugs in compromised cases can result in hemolytic anemia (Vijayvargiya et al., 2020).

In contrast, based on clinical data from 76 different studies, WHO strongly recommended against the use of hydroxychloroquine and chloroquine in COVID-19 patients independent of their severity. In case of special considerations, if patients with a history of tachyarrhythmias, ventricular tachycardia, prolonged QT interval, etc., need hydroxychloroquine or chloroquine, more precautions should be taken (Lamontagne et al., 2020). Data from 29 studies with 10,859 patients suggested the improbability of decreasing mortality and data from five studies indicated the unlikelihood of reduction of mechanical ventilation by hydroxychloroquine.


Other studies also recommended the uncertainty of hydroxychloroquine's effect on viral clearance, admission to hospital, and clinical improvement time (Lamontagne et al., 2020). A meta-analysis based on one randomized trial and 12 cohort studies that comprised of 19,573 hospitalized COVID-19 infected patients exerted no statistically significant relationship between hydroxychloroquine and mortality rate, while three cohort studies also reported no association of this drug with the abatement of mechanical ventilation and ICU administration requirements (Kashour et al., 2021). In addition, two more cohort studies also failed to display a statistically significant relationship between mechanical ventilation and ICU administration requirement in COVID-19 infection and hydroxychloroquine when co-administered with azithromycin (Kashour et al., 2021). Besides, according to Geleris et al., a study comprised of 1,446 hospitalized COVID-19 affected patients who received hydroxychloroquine showed no positive outcomes in lowering of death risk (Geleris et al., 2020). Another study conducted on 150 patients (hydroxychloroquine group = 75 and standard care group = 75) also failed to illustrate promising outcomes in terms of being COVID-19 negative (Tang W. et al., 2020). Considering all these factors, WHO decided to discontinue hydroxychloroquine as treatments arms against COVID-19 (Hossain and Rahman, 2020).

Ivermectin

In 1974, isolation of a new strain of a soil-dwelling bacterium in the laboratories of the Kitasato Institute in Japan and subsequent transferal of the sample to the laboratories of Merck & Co., Inc. in USA, gave rise to the serendipitous discovery of an active anthelmintic substance. Thus, avermectins were identified and with piecemeal research works were subsequently transmuted into antiparasitic products such as ivermectin. After much scientific research, in 1981, ivermectin was presented commercially against endoparasitic nematode and ectoparasitic arthropods in livestock, though initially the target was to apply it in veterinary science and animal husbandry (Campbell, 2012).

SARS-CoV-2, an RNA virus containing only a phospholipid envelope to preserve its genetic material with few protein inserted, does not show any protein capsid (Sigrist et al., 2020) which make them vulnerable against ivermectin (Cordova et al., 2003). Ivermectin can act upon this type of viruses by altering ionic balance between the internal and external environment by reentering water molecules, resulting in osmotic lysis (Rizzo, 2020). Thus, integrity and functionality of a viral membrane can be damaged seriously. Though the exact mechanism is not strongly elucidated, another report suggested that the probable anti-COVID-19 action of ivermectin is attributed by the inhibition of nuclear transport facilitated by the importin $\alpha/\beta 1$ heterodimer (Rizzo, 2020). This inhibitory impact can damage a notable sum of RNA viruses (Caly et al., 2012; Jans et al., 2019) including novel corona virus (Heidary and Gharebaghi, 2020). The anti-COVID-19 action of ivermectin is displayed in **Figure 5**.

According to a recent study, a single dose of ivermectin can reduce viral RNA of SARS-CoV-2 by 5000-fold in a Vero-hSLAM cell culture model at 48 h, which attracted researchers to consider it as a well-deserved drug candidate against COVID-19 attack,



though antiviral concentration of ivermectin demands a large dose (Caly et al., 2020). On the other hand, the ability of this drug to cross the blood brain barrier and affect GABAergic transmission at high doses makes its safety profile questionable (Hossen et al., 2020). A safe dose of ivermectin is considered as $\leq 200 \mu\text{g}/\text{kg}$ and overdose of this drug can cause severe adverse effects such as ataxia, psychosis, depression, seizure, and confusion (Gritti et al., 2020). Thus, dose can be a huge concern in ivermectin therapy and challenge the treatment protocol during COVID-19 infection.

Several clinical trials are being conducted with ivermectin alone or in combination to conclude about the efficacy and safety of the drug in COVID-19 patients. A recent double-blind randomized clinical trial confirmed early clinical improvement of COVID-19 patients with ivermectin and doxycycline combination compared to standard care (Hashim et al., 2020). Ivermectin was also subjected to another clinical trial among 72 hospitalized Bangladeshi SARS-CoV-2 infected adults as a randomized, placebo controlled, and double-blind testing to ensure the safety profile along with the ability of COVID-19 clearance. During this trial, patient pools were divided into three subgroups, where each group individually received either 12 mg ivermectin monotherapy once daily for 5 days or combination therapy comprised of 12 mg ivermectin and 200 mg doxycycline on day 1 followed by 100 mg of doxycycline every 12 h for next 4 days and the last subgroup was assigned as a placebo control group. Virological clearance reported from these three groups was respectively 9.7, 11.5, and 12.7 days and thus this study demonstrated efficacy of ivermectin therapy in COVID-19 cases without any severe adverse effects (Ahmed et al., 2021). In another study, ivermectin was also subjected to RT-PCR

proven COVID-19 affected patient in two different doses comprised of 6 and 12 mg in every 84 h pattern for two weeks and compared with control (lopinavir/ritonavir daily for 2 weeks) group. In this trial, mean days to negative were respectively 6.0, 4.65, and 9.15 days which demonstrated the candidacy of Ivermectin 12 mg in the clinical management of COVID-19 infection (Babalola et al., 2021). Besides, another clinical trial conducted on 600 participants showed significant abatement in RT-PCR conversion days along with mortality rate. This study recommended early use of ivermectin in COVID-19 infection as prophylaxis and to improve cytokines storm (Elgazzar et al., 2020). Moreover, in accordance with the Iranian Registry of Clinical Trials website (registration ID IRCT20200408046987N1), ivermectin is also a notable adjunct to reduce mortality rate, low O_2 duration, and hospitalization days among adult COVID-19 infected adults with a broad margin of safety and high therapeutic impact (Niaee et al., 2020).

On the contrary, one study conducted in Pakistan that comprised of 50 patients in total (equally divided into case group and control group where both groups received symptomatic treatments but ivermectin was only prescribed to case group) failed to demonstrate any statistically significant outcomes against COVID-19 infection and suggested that symptomatic treatments can be equally effective with or without ivermectin (Chachar et al., 2020). In another study, conducted among 62 mild to moderate SARS-CoV-2 infected patients, ivermectin again failed to demonstrate significant beneficial effects. In that study, total recovery time recorded for ivermectin group (ivermectin $200 \mu\text{g}/\text{kg}$ + symptomatic treatment) and control group (only symptomatic treatment)

was 10.09 ± 3.236 and 11.50 ± 5.32 respectively which discarded the standing of ivermectin therapy in mild to moderate SARS-CoV-2 infection (Podder et al., 2020). Again, WHO recommended against the use of ivermectin in COVID-19 patients except for clinical trials due to the unpredictable association with ivermectin and important clinical outcomes like mortality rate, mechanical ventilation, hospital admission, and duration of hospitalization based on data from 31 different studies (Lamontagne et al., 2020).

Favipiravir

Favipiravir was discovered many years ago during the primary *in vitro* experimentation for anti-influenza activity by chemical remodeling of a pyrazine analogue at Research Laboratories of Toyama Chemical Co., Ltd. In essence, it blocks the influenza viral RNA polymerase inhibiting viral transcription process (Furuta et al., 2017; Khambholja and Asudani, 2020). Favipiravir, one of the pronounced antiviral agents, has been tested against several SARS-CoV-2 envelope proteins. Favipiravir, a pyrazine carboxamide derivative molecule, acts as an antiviral molecule which is basically a prodrug against RNA viruses and desired antiviral property is attenuated by the competitive nature of favipiravir with purine nucleoside rather than pyrimidine nucleosides. Madin Darby Canine Kidney (MDCK) cell treated with favipiravir during *in vitro* testing provided favipiravir ribofuranosyl-5'-triphosphate (favipiravir-RTP), favipiravir ribofuranose (favipiravir-R) and favipiravir ribofuranosyl-5'-monophosphate (favipiravir-RMP) in HPLC analysis as metabolites among which chemically synthesized favipiravir-RTP demonstrated to the ability to obstruct the viral RNA polymerase activity in concentrations ranging from nanomolar to micromolar. This test reflects the projection that favipiravir, which is a prodrug, can emit an antiviral property when it is intracellularly phosphoribosylated to be an active form, favipiravir-RTP, to inhibit RNA polymerase. When a single molecule of Favipiravir-RTP is proficiently incorporated into a nascent RNA strand, a partial cessation of extension in viral RNA is detected and when double incorporation takes place, total inhibition results in (Furuta et al., 2017). The anti-COVID-19 action of favipiravir has been displayed in **Figure 4**.

Favipiravir has demonstrated prominent therapeutic impacts in COVID-19 treatment by arresting infection progression and viral removal and research has also stated that, due to mild adverse effects, the treatment duration of favipiravir can be prolonged as per requirements (Cai et al., 2020). In accordance with the official at the Chinese Ministry of Science and Technology along with the Guardian news report, favipiravir has exhibited auspicious clinical trials outcomes in Shenzhen and Wuhan in a study comprised of 340 patients (McCurry, 2020). The Turkish Ministry of Health has also focused on favipiravir in intensive care cases to decrease infection periods ranging from 11–12–4 days (Al-Monitor News, 2020). A recent nonrandomized, controlled, open-label trial reported efficacy of favipiravir in COVID-19 affected patients (1,600 mg twice in day 1 followed by 600 mg twice in day 2–14) compared to lopinavir-ritonavir combination including improved chest imaging reports in favipiravir group (91.43%) than lopinavir-

ritonavir group (62.22%) on day 14 of medication (Cai et al., 2020). On the contrary, favipiravir, a teratogenic agent, cannot be prescribed in pregnancy cases (Delang et al., 2018) and thus it demands more vigorous clinical trials from concerned regulatory bodies controlling medical care in each country to get the approval of widespread utilization (Khan et al., 2020).

At present, a few clinical trials of favipiravir are ongoing throughout several countries, including China and Japan, to reestablish the therapeutic profile of favipiravir against COVID-19 infection (Wu et al., 2020). A randomized control trial (ChiCTR200030254) has displayed that infected individuals treated with favipiravir experienced a faster recovery rate (71.43%) compared to treated individuals with umifenovir (55.86%), whereas fever duration and cough relief time are also alleviated prominently (Chen, C. et al., 2020). Up to mid-April, 2020, there were two clinical trials in Japan and eight in China under way in order to evaluate the efficacy of favipiravir against novel coronavirus. These clinical studies included non-randomized and randomized controlled trials assessing the safety and efficacy of favipiravir alone (ChiCTR2000030113, JPRN-jRCTs031190226, JPRN-jRCTs041190120) or in conjunction with interferon- α (ChiCTR2000029600), baloxavir marboxil (ChiCTR2000029544, ChiCTR2000029548), tocilizumab (ChiCTR2000030894, NCT04310228), or chloroquine phosphate (ChiCTR2000030987, NCT04319900) (Wu et al., 2020). Currently, several trials are ongoing in Italy, Saudi Arabia, Turkey, the United States, Iran, Russia, Germany, Mexico, Indonesia, Kuwait, Egypt, Bangladesh, Nepal, the United Kingdom, Hungary, and Thailand to evaluate the performance, safety, and efficacy of favipiravir alone or in combination with other agents and in comparison to other agents for mild, moderate, and severe COVID-19 treatment (Anonymous, 2020e). Five ongoing clinical trials are predicted to provide promising results for the use of favipiravir in COVID-19 treatment. This drug can be considered as a relatively safer treatment option for COVID-19 patients than other agents except in pregnancy due to teratogenicity (Maslub et al., 2021). A recently published preprint of a retrospective observational case control study (between March and September 2020 in Turkey) suggested the early use of favipiravir in COVID-19 treatment for better clinical outcomes and viral load reduction (Uçan et al., 2021). Three case-reports, one involving two elderly COVID-19 patients, another with a case of COVID-19 pneumonia, and the third with a critically ill patient, supported the use of favipiravir for beneficial effect in COVID-19 treatment. A randomized multicenter study including 96 COVID-19 patients performed between April and August 2020 reported reduced hospital stay and decreased need for mechanical ventilation with favipiravir than with Chloroquine (Dabbous et al., 2021). Another randomized, open-label, comparative, multicenter phase-3 clinical trial from May 14 to July 3, 2020 with 150 patients suggested the benefit of favipiravir in mild to moderate COVID-19 based on the median time of viral shedding cessation [5 days (95% CI: 4 days, 7 days) vs. 7 days (95% CI: 5 days, 8 days), $p = 0.129$] and clinical cure (3 days (95% CI: 3 days, 4 days) vs. 5 days (95% CI: 4 days, 6 days), $p = 0.030$) compared to control (Udwadia et al., 2021). On the other hand,

Medisetty et al. (2021) expressed concern about the conclusion made by Udawadia et al. (2021) and proposed the need for well-designed randomized, placebo controlled clinical trials for clearer evidence before using favipiravir in mild to moderate COVID-19 cases (Medisetty et al., 2021; Udawadia et al., 2021).

However, several studies demonstrated mixed results of efficacy and a complex pharmacokinetic profile of the prodrug, favipiravir, leading to dose determination challenges for COVID-19 treatment. Due to pharmacokinetic complexities and a lack of correlation between favipiravir concentration and viral load reduction, assessment of the active triphosphate form concentration in cell and tissue is required to understand its pharmacokinetic/pharmacodynamic profile. Thus, the challenges to achieve effective drug concentrations should be considered if using favipiravir to treat COVID-19 (Ison and Scheetz, 2021).

Rivaroxaban

In today's market, there are four clinically approved direct FXa inhibitors to be used as anticoagulants: rivaroxaban (approved in 2011; Xarelto) (Mueck et al., 2008), apixaban (2012; Eliquis) (Byon et al., 2019), edoxaban (2015; Savaysa) (Bounameaux and Camm, 2014), and betrixaban (2017; Bevyxxa) (Garland et al., 2018). In 2008, Bayer Healthcare was given approval for clinical use of rivaroxaban under the brand name Xarelto as an orally administered direct factor Xa inhibitor in order to prevent venous thromboembolism (VTE) after elective hip or knee replacement surgery (Perzborn et al., 2011). An early validation research on naturally occurring inhibitors for factor Xa resulting in the development of synthetic and orally selective factor Xa inhibitors in 1998 recommended rivaroxaban's potency and specificity along with good oral bioavailability. The preclinical profiles of rivaroxaban also ensure high oral bioavailability, dose-dependent pharmacokinetics, a faster onset/offset of action, negligible drug-drug or drug-food interactions, and ascertains rivaroxaban as prolific drug candidate which does not demand routine coagulation monitoring (Perzborn et al., 2011).

The spike protein of SARS-CoV-2 is comprised of a couple of vital subunits: the S1 subunit containing the host receptor binding domain and the S2 subunit, accountable for the fusion of virus and host cell membrane. The S1 subunit binds to its receptor ACE2 (angiotensin-converting enzyme 2) on the host cell while a proteolytic activation promoted by the host protease including transmembrane protease serine 2 (TMPRSS2), furin, and liposomal cathepsins results in the breakdown of the linkage between S1 and S2 subunits to ease the fusion (Belouzard et al., 2009; Shang et al., 2020). A few other host proteases like Human Factor Xa (FXa) also possess noteworthy viral entry and fusion regulation property (Belouzard et al., 2009; Shang et al., 2020). It is also reported that those FXa inhibitors can potentially block the viral entry of SARS-CoV-2 into the host by ceasing the fragmentation of spike protein into the S1 and S2 subunits. Thus, the concentration dependent inhibitors can obstruct the SARS-CoV-2 plaque formation in Vero E6 cells and thwart the entry of corona virus (Al-Horani, 2020). Additionally, a few recent studies have suggested that FXa inhibitors can potentially inhibit TMPRSS2 and act against a wide spectrum

of RNA and DNA viruses by ceasing the viral entry phase (Du et al., 2007; Kim et al., 2020). Therefore, FXa inhibitors can be a potential agent against COVID-19 attack (Al-Horani, 2020). The anti-COVID-19 action of rivaroxaban is displayed in **Figure 4**.

Direct FXa inhibitors possess significant anticoagulant, anti-inflammatory, and antiviral properties which make them potential candidates in COVID-19 management (Al-Horani, 2020). At present, more than 10 clinical studies are being carried out to investigate the role of these agents against novel corona virus where dosage regimen is kept similar to their conventional use given to thrombotic patients after considering renal function (Al-Horani, 2020). A few trials suggested usage of these agents, such as rivaroxaban for 21–30 days, though more concrete reports are still needed to ensure the desired benefits (Al-Horani, 2020). Another study conducted on 183 hospitalized patients in China documented that activation of the coagulation system can play a detrimental role in COVID-19 management (Tang N. et al., 2020). According to that research, the degree of activation of a coagulation system such as increased D-dimer concentrations during admission is much higher in non-survival cases. Overall, 21 (11%) of 183 patients died; 15 (71%) of the 21 non-survivors and only one (1%) of the 162 survivors met criteria for disseminated intravascular coagulation during their hospital stay. In a case series in China, elevated D-dimer concentrations during hospital admission ($>1 \mu\text{g/ml}$) was correlated with higher chances of death rate which is 18 times greater than normal individual (Zhou F. et al., 2020). Thus, the conducted study warned about a poor outcome in COVID-19 management if standard supportive therapy is absent (Zhou F. et al., 2020) and strongly recommended inclusion of thromboprophylaxis in corona virus treatment (Spyropoulos et al., 2020).

A few clinical trials are ongoing in the United States, Germany, Brazil, and the United Kingdom to evaluate the safety and efficacy of rivaroxaban alone or in combination with other drugs for moderate to severe COVID-19 treatment or for minimizing risk of major venous and arterial thrombotic events, hospitalization, and death in symptomatic COVID-19 infection (Anonymous, 2020f). Among them PREVENT-HD, a double-blind, placebo-controlled, pragmatic, event-driven phase 3 trial, started in August 2020 and was expected to enroll 4,000 participants to evaluate the efficacy and safety of rivaroxaban in COVID-19 outpatients with possible risk of thrombotic events, hospitalization, and death (Capell et al., 2021). A descriptive and analytical study with 50 COVID-19 patients (affected between April and May 2020) presenting pneumonia suggested the use of rivaroxaban for thromboprophylaxis (Carpio-Orantes et al., 2021).

ACEIs/ARBs

For more than 30 years, ACEIs, the first medication to target the Renin Angiotensin Aldosterone System (RAAS), have been used to treat a variety of indications associated with hypertension, cardiovascular diseases, and renal diseases (Yehualashet and Belachew, 2020). ARBs can also show efficacy in hypertensive disorders. Not only do ACEIs and ARBs reduce blood pressure, but they also demonstrate distinctive cardio protective efficacy (Maione et al., 2011).

SARS-CoV and other SARS-related coronaviruses (SARSr-CoV) can directly interact with ACE2 to penetrate to the target site. Studies found that ACE2 is strongly pronounced in the mouth and tongue, promoting viral entrance in the host. ACE2 is found on type I and II alveolar epithelial cells of lower lungs. After invasion, the penetration of SARS-CoV-2 begins with the adherence to ACE2 on the alveolar surface of the spike glycoprotein distributed on the viral envelope. Clathrin-dependent endocytosis of the SARS-CoV-2 and ACE2 complexes modulates the adhesion of SARS-CoV-2 to ACE2, prompting cell membrane fusion (Yehualashet and Belachew, 2020). SARS-CoV-2 uses the endogenous transcriptional machinery of alveolar tissues to replicate and spread throughout the entire lung once it enters the cells (Perico et al., 2020). ACE2 can aid SARS-CoV-2 S-mediated cell entry and thus establish it as an operational receptor for COVID-19. In the pathogenesis of COVID-19, angiotensin-converting enzyme inhibitors (ACEIs) and angiotensin receptor blockers (ARBs) have a distinct mechanism. ACEIs in clinical application cannot directly interfere with the activity of ACE2 (Brojakowska et al., 2020) and the intestinal messenger RNA concentration of ACE2 may be enhanced with ACEI but not reported with ARBs. SARS-CoV-2 abates the expression of ACE2, which resulted in reduction of the protective actions of ACE2 on various organs (Wysocki et al., 2020). The entrance is amplified by ACEIs more than twice compared to ARBs. This will certainly not infer the drawbacks or advantages of such treatments with patients displaying several variables such as age, hypertension, and the result of unspecified comorbidities on the outcome of the COVID-19 pandemic (Guo J. et al., 2020; Leclézio et al., 2020). By stabilizing ACE2-AT1R interaction and suppressing viral protein and ACE2 interaction and invasion, ACEIs or ARBs can arrest COVID-19 viral entrance. In the presence of stabilized ACE2-AT1R complexes, the interaction between the viral protein and ACE2 can be reduced in a significant manner (Leclézio et al., 2020). According to Yang et al. (2020), elevated cases of hypertension were found in serious COVID-19 patients though it is not concluded that the severity of this infection is associated with hypertension. Additionally, no strong data on the use of ACEIs and ARBs for COVID-19 infection has been published yet (Yang et al., 2020). According to Guo T. et al. (2020), during the evaluation of cardiovascular effects of COVID-19 infection, the use of ACEIs/ARBs failed to demonstrate any correlation with mortality case (Guo T. et al., 2020). However, previous studies on animal models have shown that ACEIs and ARBs can upsurge the action of ACE2. Thus, it is projected that ACEIs and ARBs, by improving the activity of ACE2, can increase COVID-19 infectivity. Some researchers have suggested that the use of ACEIs/ARBs may be effective in preventing COVID-19 infection (Li et al., 2017). The anti-COVID-19 action of ACEIs/ARBs is displayed in **Figure 4**.

There are several study reports on the use of ACEIs/ARB in COVID-19 which include mainly hospitalized patients. One study involving 1,128 hypertensive patients, 188 receiving ACEIs/ARB, showed in-hospital use of these drugs can reduce mortality risk (Zhang et al., 2020). In another larger study with 8,910 patients, 770 receiving ACEI and 556 receiving ARBs,

decreased in-hospital mortality rate was shown for ACEI but not for ARBs (Mehra et al., 2020). A cohort study with 8.3 million people concluded that ACEI/ARBs use is associated with decreased risk of COVID-19 for in-hospital patients. But initial reports from New York showed no evidence of a beneficial outcome for ACEI/ARB use (Hippisley-Cox et al., 2020).

However, use of ACEIs/ARB in COVID-19 is controversial. Several studies on animals have suggested these to be beneficial while some studies have shown the risk associated with their use in COVID-19 patients. More studies are required to draw a conclusion about the benefit/risk ratio (Onweni et al., 2020). Currently a randomized clinical trial in Denmark is underway to investigate the effects of continuation and discontinuation of ACEIs/ARB in COVID-19 patients and another multicenter, randomized trial with 208 participants is being conducted to test the positive outcome of stopping ACEI/ARBs in symptomatic SARS-CoV-2 patients (NCT04351581, 2020; NCT04353596, 2020). Other trials with ACEI/ARBs are being conducted in Ireland, the United States, France, Canada, Saudi Arabia, Ukraine, Brazil, Spain, and Italy (Anonymous, 2020g).

A recent preprint of a retrospective cohort study suggested higher survival rates for in-hospital high-risk COVID-19 patients with the use of ARBs than ACEIs (de Abajo et al., 2021). Another retrospective cohort study in Wuhan suggested the beneficial role of ACEI/ARBs over calcium channel blockers in COVID-19 patients with hypertension (Liu et al., 2021). A recent study involving 53 COVID-19 patients hospitalized from February 8, 2020 to February 24, 2020 concluded that discontinuation of ACEIs/ARBs in hypertensive COVID-19 positive patients increases length of hospital stay (Tian et al., 2021). A meta-analysis of the existing observational cohort studies (including more than 22,000 COVID-19 patients) on the use of ACEIs/ARBs in COVID-19 found that ACEIs/ARBs was not associated with increased mortality, disease severity, or hospital stay in COVID-19 patients, but rather showed association with reduced mortality, severity, and hospitalization (Zhang et al., 2021). On the contrary, a retrospective cohort study with 182 hypertensive COVID-19 patients found no significant association between severity of COVID-19 or mortality and the use of ACEIs/ARBs (Abdullah et al., 2021). Another preprint of a Meta-Regression Analysis (consisting of 53 cross-sectional, case-control, and cohort studies published before January 18th, 2021 involving 112,468 patients) showed no effect on mortality and severity in COVID-19 patients using ACEIs/ARBs (Singh et al., 2021).

Protease Inhibitors

The coronavirus glycoproteins allow viral entrance into target cells by attaching to receptors and accelerating viral and host cell membrane fusion. Nevertheless, to acquire a fusion-active form, glycoproteins are synthesized as inactive precursors and rely on activation by host cell proteases (Zhou et al., 2015). As a result, for acquiring broad-spectrum antiviral action, the respective enzymes can be considered as possible targets, and an effective guiding principle for the production of broad-spectrum antiviral medications will be the development of protease inhibitors (Zhou et al., 2011). Protease is also a crucial replication enzyme in the

case of SARS-CoV-2 and is beneficial for the development of wide-spectrum inhibitors. The protease is responsible for converting it into functional proteins and its activity is induced by residues binding to particular points on the protease known as active sites where inhibitors can block the aforementioned activity (Wang et al., 2016; Mothay and Ramesh, 2020). When an inhibitor binds to an active site, it prevents substrates from binding, resulting in complete inhibition of the protease's action. Thus, a groundbreaking step to battle this pandemic could be the development of an inhibitor for COVID-19's protease and with this hope several protease inhibitors are now under clinical trials including PF-07304,814, EIDD-2801, and AT- 527 (Mothay and Ramesh, 2020). Therefore, despite not having any completed clinical studies, drugs of this class are included in this review work, so that they can get much deserved priority on being repurposed in COVID-19 infection.

PF-07304,814 is a notable phosphate prodrug which is converted into PF-00835321 and a strong inhibitor of 3CLpro that shows promising antiviral efficacy against SARS-CoV-2 as a monotherapy or combined therapy with remdesivir (Boras et al., 2020). A phase 1 clinical trial sponsored by Pfizer is in recruiting status to assess the safety, tolerability, and pharmacokinetic parameters associated with single and multiple doses of PF-07304814 in Hospitalized SARS-CoV-2 infected patients. In this triple masking, placebo controlled trial, 72 participants were enrolled where both PF-07304814 and placebo were introduced intravenously. Researchers are estimating that the completion of the clinical trial and outcomes can be available in July 5, 2021 (NCT04535167, 2020) and to our best knowledge, along with this uncompleted clinical trial, there is another clinical trial on PF-07304814 (also sponsored by Pfizer) which has been completed and registered in ClinicalTrials.gov till 8th April, 2020 to evaluate the candidacy of PF-07304814 against SARS-CoV-2 (NCT04627532, 2020).

EIDD-2801, also known as MK-4482 or molnupiravir, is a protease inhibitor and prodrug of ribonucleoside analog β -D-N⁴-hydroxycytidine (NHC) which showed efficacy against SARS-CoV and MARS-CoV in mice model. Previous records have generated enough interest amongst researchers to launch clinical trials of EIDD-2801 to evaluate its efficacy to alleviate SARS-CoV-2 infection (Wahl et al., 2021). Like PF-07304814, very few completed clinical trials (NCT04392219 and NCT04405570) on EIDD-2801 have been completed while only four clinical trials are enlisted in ClinicalTrials.gov till 8th April, 2020 (NCT04392219, 2020; NCT04405570, 2020).

AT-527, a guanosine nucleotide analog, is an orally administered double prodrug that can selectively obstruct the viral RNA-dependent RNA polymerase analog (RdRp). During *in vitro* evaluation against several 76 human coronaviruses, including SARS-CoV-2, AT-511, the free base of AT-527, has showed

promising antiviral activity where the other active metabolite is AT-9010. The estimated concentration of the active metabolites in pulmonary tissues after oral dosing indicates that AT-527 can be an excellent therapeutic choice against SARS-CoV-2 infection (Good et al., 2021). There are only two clinical trials of this therapy against COVID-19 enlisted in ClinicalTrials.gov which are still in recruiting status and the estimated completion period is June, 2021 (NCT04396106, 2020; NCT04709835, 2020).

CONCLUSION

COVID-19 is one of the biggest pandemic situations in the past hundred years. In the current scenario, drug repositioning can be a feasible option along with available vaccines. Several *in silico*, *in vitro*, and *in vivo* studies along with clinical trials, case series, retrospective studies, cohort studies, and meta-analyses have been conducted to establish the safety and efficacy profile, mechanism of action, pharmacokinetics, pharmacodynamics, and future aspects of repurposing drugs against COVID-19. Although several drugs have shown their effectiveness against SARS-CoV-2 infection in some notable studies and been used in many countries, safety and efficacy profiles of those drugs are yet to be well established in large scale clinical trials. Based on current study findings, some drugs clearly failed to be effective in the treatment of COVID-19. As a consequence, WHO has recommended against the use of such types of drugs (namely chloroquine, hydroxychloroquine, remdesivir, and lopinavir/ritonavir) in the treatment of COVID-19 infection. Thus, establishing complete safety and efficacy profiles of the trial drugs is still a desperate call and collaborative efforts throughout the world are needed to achieve this aim.

AUTHOR CONTRIBUTIONS

MMRS and SA conceptualized the research work. SA and TBK searched and gathered the previous studies. SA, TBK, and MMRS wrote the manuscript. J-RZ, SMAR, and INM critically reviewed the manuscript. SA and TBK edited the reviewed manuscript. MMRS critically evaluated and revised the manuscript and supervised the whole project.

ACKNOWLEDGMENTS

Authors would like to deliver their thanks to all frontline fighters (medical doctors, pharmacists, nurses, and others healthcare workers) who are delivering services to patients, risking infection as well as psychological stress during the COVID-19 pandemic.

REFERENCES

- Abdullah, M., Khan, K., Javed, M., Iqbal, J., Waheed, K., Akhtar, M. N., et al. (2021). A Comparison of Disease Severity and Outcomes in COVID-19 Cases Taking Angiotensin-Converting Enzyme Inhibitors/Angiotensin Receptor Blockers versus Other Antihypertensive Drugs. *Cureus* 13 (1). doi:10.7759/cureus.12757
- Ahmed, S., Karim, M. M., Ross, A. G., Hossain, M. S., Clemens, J. D., Sumiya, M. K., et al. (2021). A Five-Day Course of Ivermectin for the Treatment of COVID-19 May Reduce the Duration of Illness. *Int. J. Infect. Dis.* 103, 214–216. doi:10.1016/j.ijid.2020.11.191
- Al-Abdoh, A., Bizanti, A., Barbarawi, M., Jabri, A., Kumar, A., Fashanu, O. E., et al. (2021). Remdesivir for the Treatment of COVID-19: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Contemp. Clin. Trials* 101, 106272. doi:10.1016/j.cct.2021.106272
- Al-Horani, R. A. (2020). Potential Therapeutic Roles for Direct Factor Xa Inhibitors in Coronavirus Infections. *Am. J. Cardiovasc. Drugs* 20, 525–533. doi:10.1007/s40256-020-00438-6
- Al-Monitor news (2020). Turkey Says It's Using 'special' Coronavirus Drug Sent from China. Available at: <https://www.al-monitor.com/pulse/originals/2020/03/turkey-using-special-coronavirus-drug-china.html> (Accessed January 25, 2021).
- Alam, M. M., Mahmud, S., Rahman, M. M., Simpson, J., Aggarwal, S., and Ahmed, Z. (2020). Clinical Outcomes of Early Treatment with Doxycycline for 89 High-Risk COVID-19 Patients in Long-Term Care Facilities in New York. *Cureus* 12 (8), e9658. doi:10.7759/cureus.9658
- Anonymous (2020a). Remdesivir, ClinicalTrials.Gov. Available at: <https://www.clinicaltrials.gov/ct2/results?cond=COVID19&term=remdesivir&cntry=&state=&city=&dist=> (Accessed January 25, 2021).
- Anonymous (2020b). Doxycycline, ClinicalTrials.Gov. <https://www.clinicaltrials.gov/ct2/results?cond=COVID19&term=doxycycline&cntry=&state=&city=&dist=> (Accessed January 25, 2021).
- Anonymous (2020c). Lopinavir/Ritonavir, ClinicalTrials.Gov. Available at: <https://www.clinicaltrials.gov/ct2/results?cond=COVID19&term=Lopinavir%2FRitonavir&cntry=&state=&city=&dist=> (Accessed January 25, 2021).
- Anonymous (2020d). Chloroquine, ClinicalTrials.Gov. Available at: <https://www.clinicaltrials.gov/ct2/results?cond=COVID19&term=chloroquine&cntry=&state=&city=&dist=> (Accessed January 25, 2021).
- Anonymous (2020e). Favipiravir, Clinical Trial, ClinicalTrials.Gov. Available at: <https://www.clinicaltrials.gov/ct2/results?cond=COVID19&term=favipiravir&cntry=&state=&city=&dist=> (Accessed January 25, 2021).
- Anonymous (2020f). Rivaroxaban, ClinicalTrials.Gov. Available at: <https://www.clinicaltrials.gov/ct2/results?cond=COVID19&term=Rivaroxaban+&cntry=&state=&city=&dist=#wrapper> (Accessed January 25, 2021).
- Anonymous (2020g). ACEIs/ARB, ClinicalTrials.Gov. Available at: <https://www.clinicaltrials.gov/ct2/results?cond=COVID19&term=ACEIs%2FARB&cntry=&state=&city=&dist=> (Accessed January 25, 2021).
- Arshad, S., Kilgore, P., Chaudhry, Z. S., Jacobsen, G., Wang, D. D., Huitsing, K., et al. (2020). Treatment with Hydroxychloroquine, Azithromycin, and Combination in Patients Hospitalized with COVID-19. *Int. J. Infect. Dis.* 97, 396–403. doi:10.1016/j.ijid.2020.06.099
- Babalola, O. E., Bode, C. O., Ajayi, A. A., Alakaloko, F. M., Akase, I. E., Otofawo, E., et al. (2021). Ivermectin Shows Clinical Benefits in Mild to Moderate Covid19 Disease: A Randomised Controlled Double Blind Dose Response Study in Lagos. *medRxiv*. doi:10.1101/2021.01.05.21249131
- Ballow, C. H., and Amsden, G. W. (1992). Azithromycin: the First Azalide Antibiotic. *Ann. Pharmacother.* 26 (10), 1253–1261. doi:10.1177/106002809202601014
- Belouzard, S., Chu, V. C., and Whittaker, G. R. (2009). Activation of the SARS Coronavirus Spike Protein via Sequential Proteolytic Cleavage at Two Distinct Sites. *Proc. Natl. Acad. Sci. U S A.* 106 (14), 5871–5876. doi:10.1073/pnas.0809524106
- Boras, B., Jones, R. M., Anson, B. J., Arendson, D., Aschenbrenner, L., Bakowski, M. A., et al. (2020). Discovery of a Novel Inhibitor of Coronavirus 3CL Protease as a Clinical Candidate for the Potential Treatment of COVID-19. *bioRxiv*. doi:10.1101/2020.09.12.293498
- Borba, M., de Almeida Val, F., Sampaio, V. S., Alexandre, M. A., Melo, G. C., Brito, M., et al. (2020). Chloroquine Diphosphate in Two Different Dosages as Adjunctive Therapy of Hospitalized Patients with Severe Respiratory Syndrome in the Context of Coronavirus (SARS-CoV-2) Infection: Preliminary Safety Results of a Randomized, Double-Blinded, Phase IIb Clinical Trial (CloroCOVID-19 Study). *MedRxiv*. doi:10.1101/2020.04.07.20056424
- Bounameaux, H., and Camm, A. J. (2014). Edoxaban: an Update on the New Oral Direct Factor Xa Inhibitor. *Drugs* 74 (11), 1209–1231. doi:10.1007/s40265-014-0261-1
- Brojakowska, A., Narula, J., Shimony, R., and Bander, J. (2020). Clinical Implications of SARS-CoV-2 Interaction with Renin Angiotensin System: JACC Review Topic of the Week. *J. Am. Coll. Cardiol.* 75 (24), 3085–3095. doi:10.1016/j.jacc.2020.04.028
- Bull-Otterson, L., Gray, E. B., Budnitz, D. S., Strosnider, H. M., Schieber, L. Z., Courtney, J., et al. (2020). Hydroxychloroquine and Chloroquine Prescribing Patterns by Provider Specialty Following Initial Reports of Potential Benefit for COVID-19 Treatment - United States, January-June 2020. *MMWR Morb Mortal Wkly Rep.* 69 (35), 1210–1215. doi:10.15585/mmwr.mm6935a4
- Butler, C. C., Dorward, J., Yu, L. M., Gbinigie, O., Hayward, G., Saville, B. R., et al. (2021). Azithromycin for Community Treatment of Suspected COVID-19 in People at Increased Risk of an Adverse Clinical Course in the UK (PRINCIPLE): a Randomised, Controlled, Open-Label, Adaptive Platform Trial. *The Lancet* 397, 1063–1074. doi:10.1016/S0140-6736(21)00461-X
- Byon, W., Garonzik, S., Boyd, R. A., and Frost, C. E. (2019). Apixaban: a Clinical Pharmacokinetic and Pharmacodynamic Review. *Clin. Pharmacokinet.* 58 (10), 1265–1279. doi:10.1007/s40262-019-00775-z
- Cai, Q., Yang, M., Liu, D., Chen, J., Shu, D., Xia, J., et al. (2020). Experimental Treatment with Favipiravir for COVID-19: an Open-Label Control Study. *Engineering (Beijing)* 6 (10), 1192–1198. doi:10.1016/j.eng.2020.03.007
- Caly, L., Druce, J. D., Catton, M. G., Jans, D. A., and Wagstaff, K. M. (2020). The FDA-Approved Drug Ivermectin Inhibits the Replication of SARS-CoV-2 *In Vitro*. *Antivir. Res.* 178, 104787. doi:10.1016/j.antiviral.2020.104787
- Caly, L., Wagstaff, K. M., and Jans, D. A. (2012). Nuclear Trafficking of Proteins from RNA Viruses: Potential Target for Antivirals? *Antivir. Res.* 95 (3), 202–206. doi:10.1016/j.antiviral.2012.06.008
- Campbell, W. C. (2012). History of Avermectin and Ivermectin, with Notes on the History of Other Macrocytic Lactone Antiparasitic Agents. *Curr. Pharm. Biotechnol.* 13 (6), 853–865. doi:10.2174/138920112800399095
- Cao, B., Wang, Y., Wen, D., Liu, W., Wang, J., Fan, G., et al. (2020). A Trial of Lopinavir-Ritonavir in Adults Hospitalized with Severe COVID-19. *New Engl. J. Med.* 382, 1787–1799. doi:10.1056/NEJMoa2001282
- Capell, W. H., Barnathan, E. S., Piazza, G., Spyropoulos, A. C., Hsia, J., Bull, S., et al. (2021). Rationale and Design for the Study of Rivaroxaban to Reduce Thrombotic Events, Hospitalization and Death in Outpatients with COVID-19: The PREVENT-HD Study. *Am. Heart J.* 235, 12–23. doi:10.1016/j.ahj.2021.02.001
- Chandwani, A., and Shuter, J. (2008). Lopinavir/ritonavir in the Treatment of HIV-1 Infection: a Review. *Ther. Clin. Risk Manag.* 4 (5), 1023–1033. doi:10.2147/tcrm.s3285
- Chen, C., Huang, J., Cheng, Z., Wu, J., Chen, S., Zhang, Y., et al. (2020). Favipiravir versus Arbidol for COVID-19: a Randomized Clinical Trial. *MedRxiv*. doi:10.1101/2020.03.17.20037432
- Chen, Z., Hu, J., Zhang, Z., Jiang, S., Han, S., Yan, D., et al. (2020). Efficacy of Hydroxychloroquine in Patients with COVID-19: Results of a Randomized Clinical Trial. *MedRxiv*. doi:10.1101/2020.03.22.20040758
- Chu, C. M., Cheng, V. C., Hung, I. F., Wong, M. M., Chan, K. H., Chan, K. S., et al. (2004). Role of Lopinavir/ritonavir in the Treatment of SARS: Initial Virological and Clinical Findings. *Thorax* 59 (3), 252–256. doi:10.1136/thorax.2003.012658
- Colson, P., Rolain, J. M., Lagier, J. C., Brouqui, P., and Raoult, D. (2020). Chloroquine and Hydroxychloroquine as Available Weapons to Fight COVID-19. *Int. J. Antimicrob. Agents* 55 (4), 105932. doi:10.1016/j.ijantimicag.2020.105932
- Cordova, A., Deserno, M., Gelbart, W. M., and Ben-Shaul, A. (2003). Osmotic Shock and the Strength of Viral Capsids. *Biophys. J.* 85 (1), 70–74. doi:10.1016/S0006-3495(03)74455-5
- Dabbous, H. M., Abd-Elsalam, S., El-Sayed, M. H., Sherief, A. F., Ebeid, F. F. S., El Ghafar, M. S. A., et al. (2021). Efficacy of Favipiravir in COVID-19 Treatment: a Multi-center Randomized Study. *Arch. Virol.* 166 (3), 949–954. doi:10.1007/s00705-021-04956-9
- Damle, B., Vourvahis, M., Wang, E., Leaney, J., and Corrigan, B. (2020). Clinical Pharmacology Perspectives on the Antiviral Activity of Azithromycin and Use in COVID-19. *Clin. Pharmacol. Ther.* 108 (2), 201–211. doi:10.1002/cpt.1857
- Davies, M., Osborne, V., Lane, S., Roy, D., Dhanda, S., Evans, A., et al. (2020). Remdesivir in Treatment of COVID-19: A Systematic Benefit-Risk Assessment. *Drug Saf.* 43, 645–656. doi:10.1007/s40264-020-00952-1

- de Abajo, F. J., Rodríguez-Miguel, A., Rodríguez-Martín, S., Lerma, V., and García-Lledó, A. MED-ACE2-COVID19 Study Group (2021). In-hospital Continuation with Angiotensin Receptor Blockers Is Associated with a Lower Mortality Rate Than Continuation with Angiotensin Converting Enzyme Inhibitors in Covid-19 Patients: a Retrospective Cohort Study. *medRxiv*. doi:10.1101/2021.02.01.21250853
- Delang, L., Abdelnabi, R., and Neyts, J. (2018). Favipiravir as a Potential Countermeasure against Neglected and Emerging RNA Viruses. *Antivir. Res.* 153, 85–94. doi:10.1016/j.antiviral.2018.03.003
- Du, L., Kao, R. Y., Zhou, Y., He, Y., Zhao, G., Wong, C., et al. (2007). Cleavage of Spike Protein of SARS Coronavirus by Protease Factor Xa Is Associated with Viral Infectivity. *Biochem. Biophys. Res. Commun.* 359 (1), 174–179. doi:10.1016/j.bbrc.2007.05.092
- Eastman, R. T., Roth, J. S., Brimacombe, K. R., Simeonov, A., Shen, M., Patnaik, S., et al. (2020). Remdesivir: a Review of its Discovery and Development Leading to Emergency Use Authorization for Treatment of COVID-19. *ACS Cent. Sci.* 6 (5), 672–683. doi:10.1021/acscentsci.0c00489
- Echeverría-Esnal, D., Martín-Ontiyuelo, C., Navarrete-Rouco, M. E., De-Antonio Cuscó, M., Ferrández, O., Horcajada, J. P., et al. (2021). Azithromycin in the Treatment of COVID-19: a Review. *Expert Rev. anti-infective Ther.* 19 (2), 147–163. doi:10.1080/14787210.2020.1813024
- Elgazzar, A., Hany, B., Youssef, S. A., Hafez, M., Moussa, H., and eltaweel, A. (2020). Efficacy and Safety of Ivermectin for Treatment and Prophylaxis of COVID-19 Pandemic. *Res. Square*.
- Esper, R. B., da Silva, R. S., Oikawa, F. T. C., Castro, M. M., Razuk-Filho, A., Batista, P. B., et al. (2020). *Empirical Treatment with Hydroxychloroquine and Azithromycin for Suspected Cases of COVID-19 Followed-Up by Telemedicine*. Brazil, São Paulo: Prevent Senior Institute SP, 25.
- Farouk, A., and Salman, S. (2020). Dapsone and Doxycycline Could Be Potential Treatment Modalities for COVID-19. *Med. Hypotheses* 140, 109768. doi:10.1016/j.mehy.2020.109768
- Ferner, R. E., and Aronson, J. K. (2020). Chloroquine and Hydroxychloroquine in Covid-19. *BMJ* 369, m1432. doi:10.1136/bmj.m1432
- Folegatti, P. M., Ewer, K. J., Aley, P. K., Angus, B., Becker, S., Belij-Rammerstorfer, S., et al. (2020). Safety and Immunogenicity of the ChAdOx1 nCoV-19 Vaccine against SARS-CoV-2: a Preliminary Report of a Phase 1/2, Single-Blind, Randomised Controlled Trial. *Lancet* 396 (10249), 467–478. doi:10.1016/S0140-6736(20)31604-4
- Frisk-Holmberg, M., Bergqvist, Y., and Englund, U. (1983). Chloroquine Intoxication. *Br. J. Clin. Pharmacol.* 15 (4), 502–503. doi:10.1111/j.1365-2125.1983.tb01540.x
- Furuta, Y., Komeno, T., and Nakamura, T. (2017). Favipiravir (T-705), a Broad Spectrum Inhibitor of Viral RNA Polymerase. *Proc. Jpn. Acad. Ser. B Phys. Biol. Sci.* 93 (7), 449–463. doi:10.2183/pjab.93.027
- Gao, J., Tian, Z., and Yang, X. (2020). Breakthrough: Chloroquine Phosphate Has Shown Apparent Efficacy in Treatment of COVID-19 Associated Pneumonia in Clinical Studies. *Biosci. Trends* 14, 72–73. doi:10.5582/bst.2020.01047
- Garland, S. G., DeRemer, C. E., Smith, S. M., and Gums, J. G. (2018). Betrixaban: a New Oral Factor Xa Inhibitor for Extended Venous Thromboembolism Prophylaxis in High-Risk Hospitalized Patients. *Ann. Pharmacother.* 52 (6), 554–561. doi:10.1177/1060028018754383
- Gautret, P., Lagier, J. C., Honoré, S., Hoang, V. T., Colson, P., and Raoult, D. (2021). Hydroxychloroquine and Azithromycin as a Treatment of COVID-19: Results of an Open Label Non-randomized Clinical Trial Revisited. *Int. J. Antimicrob. Agents* 57 (1), 106243. doi:10.1016/j.ijantimicag.2020.106243
- Gautret, P., Lagier, J. C., Parola, P., Hoang, V. T., Meddeb, L., Mailhe, M., et al. (2020). Hydroxychloroquine and Azithromycin as a Treatment of COVID-19: Results of an Open-Label Non-randomized Clinical Trial. *Int. J. Antimicrob. Agents* 56 (1), 105949. doi:10.1016/j.ijantimicag.2020.105949
- Geleris, J., Sun, Y., Platt, J., Zucker, J., Baldwin, M., Hripcsak, G., et al. (2020). Observational Study of Hydroxychloroquine in Hospitalized Patients with Covid-19. *N. Engl. J. Med.* 382 (25), 2411–2418. doi:10.1056/NEJMoa2012410
- Gendrot, M., Andreani, J., Jardot, P., Hutter, S., Delandre, O., Boxberger, M., et al. (2020). *In Vitro* antiviral Activity of Doxycycline against SARS-CoV-2. *Molecules* 25 (21), 5064. doi:10.3390/molecules25215064
- Guidicessi, J. R., Noseworthy, P. A., Friedman, P. A., and Ackerman, M. J. (2020). Urgent Guidance for Navigating and Circumventing the QTc-Prolonging and Torsadogenic Potential of Possible Pharmacotherapies for Coronavirus Disease 19 (COVID-19). *Mayo Clin. Proc.* 95 (No. 6), 1213–1221. doi:10.1016/j.mayocp.2020.03.024
- Goldberg, E., Zvi, H. B., Sheena, L., Sofer, S., Krause, I., Sklan, E. H., et al. (2021). A Real-Life Setting Evaluation of the Effect of Remdesivir on Viral Load in COVID-19 Patients Admitted to a Large Tertiary center in Israel. *Clin. Microbiol. Infect.* S1198-743X, 00113. doi:10.1016/j.cmi.2021.02.029
- Good, S. S., Westover, J., Jung, K. H., Zhou, X. J., Moussa, A., La Colla, P., et al. (2021). AT-527, a Double Prodrug of a Guanosine Nucleotide Analog, Is a Potent Inhibitor of SARS-CoV-2 *In Vitro* and a Promising Oral Antiviral for Treatment of COVID-19. *Antimicrob. Agents Chemother.* 65, e02479-20. doi:10.1128/AAC.02479-20
- Gritti, G., Raimondi, F., Ripamonti, D., Riva, I., Landi, F., Alborghetti, L., et al. (2020). IL-6 Signalling Pathway Inactivation with Siltuximab in Patients with COVID-19 Respiratory Failure: an Observational Cohort Study. *medRxiv*. doi:10.1101/2020.04.01.20048561
- Guo, J., Huang, Z., Lin, L., and Lv, J. (2020). Coronavirus Disease 2019 (COVID-19) and Cardiovascular Disease: A Viewpoint on the Potential Influence of Angiotensin-Converting Enzyme Inhibitors/Angiotensin Receptor Blockers on Onset and Severity of Severe Acute Respiratory Syndrome Coronavirus 2 Infection. *J. Am. Heart Assoc.* 9 (7), e016219. doi:10.1161/JAHA.120.016219
- Guo, T., Fan, Y., Chen, M., Wu, X., Zhang, L., He, T., et al. (2020). Cardiovascular Implications of Fatal Outcomes of Patients with Coronavirus Disease 2019 (COVID-19). *JAMA Cardiol.* 5 (7), 811–818. doi:10.1001/jamacardio.2020.1017
- Gupte, V., Hegde, R., Sawant, S., Kalathingal, K., Jadhav, S., Malabade, R., et al. (2021). Efficacy and Safety of Remdesivir in COVID-19 Patients: A Retrospective Analysis of Active Surveillance Database. doi:10.21203/rs.3.rs-366384/v1
- Harrison, C. (2020). Coronavirus Puts Drug Repurposing on the Fast Track. *Nat. Biotechnol.* 38 (4), 379–381. doi:10.1038/d41587-020-00003-1
- Hashim, H. A., Maulood, M. F., Rasheed, A. M., Fatak, D. F., Kabah, K. K., and Abdulamir, A. S. (2020). Controlled Randomized Clinical Trial on Using Ivermectin with Doxycycline for Treating COVID-19 Patients in Baghdad, Iraq. *medRxiv*. doi:10.1101/2020.10.26.20219345
- Heidary, F., and Gharebaghi, R. (2020). Ivermectin: a Systematic Review from Antiviral Effects to COVID-19 Complementary Regimen. *J. Antibiot. (Tokyo)* 73 (9), 593–602. doi:10.1038/s41429-020-0336-z
- Hippisley-Cox, J., Tan, P. S., Coupland, C., Channon, K. M., San Tan, P., Harrison, D. A., et al. (2020). Risk of Severe COVID-19 Disease with ACE Inhibitors and Angiotensin Receptor Blockers: Cohort Study Including 8.3 Million People. *Heart* 106 (19), 1503–1511. doi:10.1136/heartjnl-2020-31748610.1136/heartjnl-2020-318314
- Holshue, M. L., DeBolt, C., Lindquist, S., Lofy, K. H., Wiesman, J., Bruce, H., et al. (2020). First Case of 2019 Novel Coronavirus in the United States. *N. Engl. J. Med.* 382, 929–936. doi:10.1056/NEJMoa2001191
- Hossain, M. J., and Rahman, S. M. A. (2020). Repurposing Therapeutic Agents against SARS-CoV-2 Infection: Most Promising and Neoteric Progress. *Expert Rev. Anti Infect. Ther.*, 1–19. doi:10.1080/14787210.2021.1864327
- Hossen, M. S., Berek, M. A., Jahan, N., and Safiqul Islam, M. (2020). A Review on Current Repurposing Drugs for the Treatment of COVID-19: Reality and Challenges. *SN Compr. Clin. Med.* 2, 1–13. doi:10.1007/s42399-020-00485-9
- Ishaqui, A. A., Khan, A. H., Sulaiman, S. A. S., Alsultan, M. T., Khan, I., and Naqvi, A. A. (2020). Assessment of Efficacy of Oseltamivir-Azithromycin Combination Therapy in Prevention of Influenza-A (H1N1) pdm09 Infection Complications and Rapidity of Symptoms Relief. *Exp. Rev. Resp. Med.* 14 (5), 533–541.
- Iglesias Cubero, G., Rodriguez Reguero, J. J., and Rojo Ortega, J. M. (1993). Restrictive Cardiomyopathy Caused by Chloroquine. *Br. Heart J.* 69 (5), 451–452. doi:10.1136/hrt.69.5.451
- Ison, M. G., and Scheetz, M. H. (2021). Understanding the Pharmacokinetics of Favipiravir: Implications for Treatment of Influenza and COVID-19. *EBioMedicine* 63, 103204. doi:10.1016/j.ebiom.2020.103204
- Jans, D. A., Martin, A. J., and Wagstaff, K. M. (2019). Inhibitors of Nuclear Transport. *Curr. Opin. Cell Biol.* 58, 50–60. doi:10.1016/j.ccb.2019.01.001
- Jie, Z., He, H., Xi, H., and Zhi, Z. (2020). [Expert Consensus on Chloroquine Phosphate for the Treatment of Novel Coronavirus Pneumonia]. *Zhonghua Jie He He Hu Xi Za Zhi.* 43 (3), E019–E188. doi:10.3760/cma.j.issn.1001-0939.2020.0019
- Joo, E. J., Ko, J. H., Kim, S. E., Kang, S. J., Baek, J. H., Heo, E. Y., et al. (2021). Clinical and Virologic Effectiveness of Remdesivir Treatment for Severe Coronavirus Disease 2019 (COVID-19) in Korea: a Nationwide Multicenter Retrospective Cohort Study. *J. Korean Med. Sci.* 36 (11), e83. doi:10.3346/jkms.2021.36.e83

- Joshi, N., and Miller, D. Q. (1997). Doxycycline Revisited. *Arch. Intern. Med.* 157 (13), 1421–1428. doi:10.1001/archinte.157.13.1421
- Juurink, D. N. (2020). Safety Considerations with Chloroquine, Hydroxychloroquine and Azithromycin in the Management of SARS-CoV-2 Infection. *CMAJ* 192 (17), E450–E453. doi:10.1503/cmaj.200528
- Kaka, A. S., MacDonald, R., Greer, N., Vela, K., Duan-Porter, W., Obley, A., et al. (2021). Major Update: Remdesivir for Adults with COVID-19: A Living Systematic Review and Meta-Analysis for the American College of Physicians Practice Points. *Ann. Intern. Med.* 174, 663–672. doi:10.7326/M20-8148
- Kashour, Z., Riaz, M., Garbati, M. A., AlDosary, O., Tlayjeh, H., Gerberi, D., et al. (2021). Efficacy of Chloroquine or Hydroxychloroquine in COVID-19 Patients: a Systematic Review and Meta-Analysis. *J. Antimicrob. Chemother.* 76 (1), 30–42. doi:10.1093/jac/dkaa403
- Keyaerts, E., Li, S., Vijgen, L., Rysman, E., Verbeeck, J., Van Ranst, M., et al. (2009). Antiviral Activity of Chloroquine against Human Coronavirus OC43 Infection in Newborn Mice. *Antimicrob. Agents Chemother.* 53 (8), 3416–3421. doi:10.1128/AAC.01509-08
- Khambholja, K., and Asudani, D. (2020). Potential Repurposing of Favipiravir in COVID-19 Outbreak Based on Current Evidence. *Trav. Med Infect Dis.* 35, 101710. doi:10.1016/j.tmaid.2020.101710
- Khan, Z., Karataş, Y., Ceylan, A. F., and Rahman, H. (2020). COVID-19 and Therapeutic Drugs Repurposing in Hand: the Need for Collaborative Efforts. *Le Pharm. Hospitalier et Clin.* 56, 3–11. doi:10.1016/j.phclin.2020.06.003
- Kim, J., Zhang, J., Cha, Y., Koltz, S., Funt, J., Escalante Chong, R., et al. (2020). Advanced Bioinformatics Rapidly Identifies Existing Therapeutics for Patients with Coronavirus Disease-2019 (COVID-19). *J. Transl. Med.* 18 (1), 257–259. doi:10.1186/s12967-020-02430-9
- Lamontagne, F., Agoritsas, T., Macdonald, H., Leo, Y. S., Diaz, J., Agarwal, A., et al. (2020). A Living WHO Guideline on Drugs for Covid-19. *Bmj* 370, m3379. doi:10.1136/bmj.m3379
- Leclézio, A., Robinson, J., and Banerjee, I. (2020). SARS-CoV-2: ACE Inhibitors, Disastrous or Desirable?. *J. Biomed. Sci.* 7 (1), 40–46. doi:10.3126/jbs.v7i1.29852
- Lepage, M. A., Rozza, N., Kremer, R., and Grunbaum, A. (2021). Safety and Effectiveness Concerns of Lopinavir/ritonavir in COVID-19 Affected Patients: a Retrospective Series. *Clin. Toxicol. (Phila)*, 1–4. doi:10.1080/15563650.2020.1842882
- Li, X. C., Zhang, J., and Zhuo, J. L. (2017). The Vasoprotective Axes of the Renin-Angiotensin System: Physiological Relevance and Therapeutic Implications in Cardiovascular, Hypertensive and Kidney Diseases. *Pharmacol. Res.* 125, 21–38. doi:10.1016/j.phrs.2017.06.005
- Lim, J., Jeon, S., Shin, H. Y., Kim, M. J., Seong, Y. M., Lee, W. J., et al. (2020). Case of the Index Patient Who Caused Tertiary Transmission of COVID-19 Infection in Korea: the Application of Lopinavir/Ritonavir for the Treatment of COVID-19 Infected Pneumonia Monitored by Quantitative RT-PCR. *J. Korean Med. Sci.* 35 (6), e79. doi:10.3346/jkms.2020.35.e79
- Liu, J., Cao, R., Xu, M., Wang, X., Zhang, H., Hu, H., et al. (2020). Hydroxychloroquine, a Less Toxic Derivative of Chloroquine, Is Effective in Inhibiting SARS-CoV-2 Infection *In Vitro*. *Cell Discov.* 6 (1), 16. doi:10.1038/s41421-020-0156-0
- Liu, X., Liu, Y., Chen, K., Yan, S., Bai, X., Li, J., et al. (2021). Efficacy of ACEIs/ARBs vs CCBs on the Progression of COVID-19 Patients with Hypertension in Wuhan: A Hospital-Based Retrospective Cohort Study. *J. Med. Virol.* 93 (2), 854–862. doi:10.1002/jmv.26315
- Logunov, D. Y., Dolzhikova, I. V., Shcheplyakov, D. V., Tukhvatulin, A. I., Zubkova, O. V., Dzharullaeva, A. S., et al. (2021). Safety and Efficacy of an rAd26 and rAd5 Vector-Based Heterologous Prime-Boost COVID-19 Vaccine: an Interim Analysis of a Randomised Controlled Phase 3 Trial in Russia. *The Lancet* 397 (10275), 671–681. doi:10.1016/s0140-6736(21)00234-8
- Lora-Tamayo, J., Maestro, G., Lalueza, A., Rubio-Rivas, M., Paul, G. V., Fernández, F. A., et al. (2021). Early Lopinavir/ritonavir Does Not Reduce Mortality in COVID-19 Patients: Results of a Large Multicenter Study. *J. Infect.* 82, 276–316. doi:10.1016/j.jinf.2021.02.011
- Luo, P., Zheng, J.-l., Liu, Y., Qiu, L., Liu, X.-l., Xue, H.-y., et al. (2021). Effectiveness of Early Treatment of Lopinavir-Ritonavir in Patients with Severe COVID-19: a Case Series. *Clin. Med.* 21 (1), e80–e83. doi:10.7861/clinmed.2020-0348
- Machiels, J. D., Bleeker-Rovers, C. P., Ter Heine, R., Rahamat-Langendoen, J., de Mast, Q., Ten Oever, J., et al. (2020). Reply to Gautret et al: hydroxychloroquine sulfate and azithromycin for COVID-19: what is the evidence and what are the risks? *Int. J. Antimicrob. Agents* 56 (1), 106056. doi:10.1016/j.ijantimicag.2020.106056
- Maione, A., Navaneethan, S. D., Graziano, G., Mitchell, R., Johnson, D., Mann, J. F., et al. (2011). Angiotensin-converting Enzyme Inhibitors, Angiotensin Receptor Blockers and Combined Therapy in Patients with Micro- and Macroalbuminuria and Other Cardiovascular Risk Factors: a Systematic Review of Randomized Controlled Trials. *Nephrol. Dial. Transpl.* 26 (9), 2827–2847. doi:10.1093/ndt/gfq792
- Malek, A. E., Granwehr, B. P., and Kontoyiannis, D. P. (2020). Doxycycline as a Potential Partner of COVID-19 Therapies. *IDCases* 21, e00864. doi:10.1016/j.idcr.2020.e00864
- Maslub, M. G., Radwan, M. A., Mikhail, M. S., and Abdalla, Z. A. (2021). Could Favipiravir Be an Effective and Safe Option for the Management of Covid-19?. *Eur. J. Biomed.* 8 (2), 13–33. doi:10.17605/OSF.IO/SVPHJ
- McCurry, J. (2020). Japanese Flu Drug “Clearly Effective” in Treating Coronavirus, Says China, the Guardian. Available at: <https://www.theguardian.com/world/2020/mar/18/japanese-flu-drug-clearly-effective-in-treating-coronavirus-says-china> (Accessed January 25, 2021).
- Medisetty, M. K., Patel, A., and Pujari, S. (2021). Letter to the editor on “Efficacy and Safety of Favipiravir, an Oral RNA-Dependent RNA Polymerase Inhibitor, in Mild-to-Moderate COVID-19: A Randomized, Comparative, Open-Label, Multicenter, Phase 3 Clinical Trial” by Udwardia Z et al. *Int. J. Infect. Dis.* 105, 722. doi:10.1016/j.ijid.2021.02.035
- Mehra, M. R., Desai, S. S., Kuy, S., Henry, T. D., and Patel, A. N. (2020). Retraction: Cardiovascular Disease, Drug Therapy, and Mortality in Covid-19. *N Engl J Med.* DOI: 10.1056/NEJMoa2007621. *N. Engl. J. Med.* 382 (25), 2582. doi:10.1056/NEJMoa2007621.1056/NEJMc2021225
- Meini, S., Pagotto, A., Longo, B., Vendramin, I., Pecori, D., and Tascini, C. (2020). Role of Lopinavir/Ritonavir in the Treatment of COVID-19: a Review of Current Evidence, Guideline Recommendations, and Perspectives. *Jcm* 9 (7), 2050. doi:10.3390/jcm9072050
- Meybody, Z. A., Owlia, M. B., and Owlia, S. (2021). Efficacy of Doxycycline in Treating COVID-19 Positive Patients: A Case Series. doi:10.21203/rs.3.rs-141875/v3
- Molina, J. M., Delaugerre, C., Le Goff, J., Mela-Lima, B., Ponscarne, D., Goldwirt, L., et al. (2020). No Evidence of Rapid Antiviral Clearance or Clinical Benefit with the Combination of Hydroxychloroquine and Azithromycin in Patients with Severe COVID-19 Infection. *Med. Mal Infect.* 50 (4), 384. doi:10.1016/j.medmal.2020.03.006
- Mothay, D., and Ramesh, K. V. (2020). Binding Site Analysis of Potential Protease Inhibitors of COVID-19 Using AutoDock. *Virusdisease* 31 (2), 1–6. doi:10.1007/s13337-020-00585-z
- Mueck, W., Eriksson, B. I., Bauer, K. A., Borris, L., Dahl, O. E., Fisher, W. D., et al. (2008). Population Pharmacokinetics and Pharmacodynamics of Rivaroxaban-Aan Oral, Direct Factor Xa Inhibitor-In Patients Undergoing Major Orthopaedic Surgery. *Clin. Pharmacokinet.* 47 (3), 203–216. doi:10.2165/00003088-200847030-00006
- Mulangu, S., Dodd, L. E., Davey, R. T., Jr, Tshiani Mbaya, O., Proschan, M., Mukadi, D., et al. (2019). A Randomized, Controlled Trial of Ebola Virus Disease Therapeutics. *N. Engl. J. Med.* 381 (24), 2293–2303. doi:10.1056/NEJMoa1910993
- Narendrakumar, L., Joseph, I., and Thomas, S. (2021). Potential Effectiveness and Adverse Implications of Repurposing Doxycycline in COVID-19 Treatment. *Expert Rev. Anti-infective Ther.* 24, 1–8. doi:10.1080/14787210.2021.1865803
- NCT04351581 (2020). Effects of Discontinuing Renin-Angiotensin System Inhibitors in Patients with COVID-19 (RASCOVID-19). ClinicalTrials.Gov. Available at: <https://www.clinicaltrials.gov/ct2/show/NCT04351581?term=ACEIs%2FARB&cond=COVID19&draw=2&rank=1> (Accessed on January 25, 2021).
- NCT04353596 (2020). Stopping ACE-Inhibitors in COVID-19 (ACEI-COVID), ClinicalTrials.Gov. Available at: <https://www.clinicaltrials.gov/ct2/show/NCT04353596?term=ACEI&cond=COVID19&draw=2&rank=1> (Accessed January 25, 2021).
- NCT04392219 (2020). COVID-19 First in Human Study to Evaluate Safety, Tolerability and Pharmacokinetics of EIDD-2801 in Healthy Volunteers. Available at: <https://clinicaltrials.gov/ct2/show/NCT04392219?term=EIDD-2801&draw=2&rank=1> (Accessed April 8, 2021).

- NCT04396106 (2020). Safety and Efficacy of AT-527 in Subjects with Moderate Coronavirus Disease (COVID-19), ClinicalTrials.Gov. Available at: <https://www.clinicaltrials.gov/ct2/show/NCT04396106?term=AT-527&draw=2&rank=4> (Accessed February 21, 2021).
- NCT04405570 (2020). A Safety, Tolerability and Efficacy of Molnupiravir (EIDD-2801) to Eliminate Infectious Virus Detection in Persons with COVID-19. Available at: <https://clinicaltrials.gov/ct2/show/NCT04405570?term=EIDD-2801&draw=2&rank=3> (Accessed April 8, 2021).
- NCT04523831 (2020). Clinical Trial of Ivermectin Plus Doxycycline for the Treatment of Confirmed COVID-19 Infection, ClinicalTrials.Gov. Available at: <https://clinicaltrials.gov/ct2/show/results/NCT04523831?view=results> (Accessed January 25, 2021).
- NCT04535167 (2020). First-In-Human Study to Evaluate Safety, Tolerability, and Pharmacokinetics Following Single Ascending and Multiple Ascending Doses of PF-07304814 in Hospitalized Participants with COVID-19, ClinicalTrials.Gov. Available at: <https://www.clinicaltrials.gov/ct2/show/NCT04535167?term=PF-07304814&draw=2&rank=1> (Accessed February 21, 2021).
- NCT04627532 (2020). Single Ascending Dose Study of Intravenous Infusion of PF 07304814 in Healthy Adult Participants. Available at: <https://clinicaltrials.gov/ct2/show/NCT04627532?term=PF-07304814&draw=2&rank=2> (Accessed April 8, 2021).
- NCT04709835 (2020). Study to Evaluate the Effects of AT-527 in Non-hospitalized Adult Patients with Mild or Moderate COVID-19. Available at: <https://clinicaltrials.gov/ct2/show/NCT04709835?term=AT-527&draw=2&rank=2> (Accessed April 8, 2021).
- Niaee, M. S., Gheibi, N., Namdar, P., Allami, A., Zolghadr, L., Javadi, A., et al. (2020). Ivermectin as an Adjunct Treatment for Hospitalized Adult COVID-19 Patients: A Randomized Multi-center Clinical Trial. doi:10.21203/rs.3.rs-109670/v1
- Onweni, C. L., Zhang, Y. S., Caulfield, T., Hopkins, C. E., Fairweather, L., and Freeman, W. D. (2020). ACEI/ARB Therapy in COVID-19: the Double-Edged Sword of ACE2 and SARS-CoV-2 Viral Docking. *Crit. Care* 24 (1), 475. doi:10.1186/s13054-020-03195-9
- Orantes, L. D. C., Díaz, J. S. S., Moguel, K. G. P., Sánchez, E. E. A., Rodríguez, O. I. S., Rodríguez, O. I. S., et al. (2021). Oral Anticoagulation with Rivaroxaban as Thromboprophylaxis in Patients Recovered from COVID-19 Pneumonia in Veracruz, Mexico. *Jacoco* 13 (1), 12–15. doi:10.15406/jacoco.2021.13.00463
- Paton, N. I., Lee, L., Xu, Y., Ooi, E. E., Cheung, Y. B., Archuleta, S., et al. (2011). Chloroquine for Influenza Prevention: a Randomised, Double-Blind, Placebo Controlled Trial. *Lancet Infect. Dis.* 11 (9), 677–683. doi:10.1016/S1473-3099(11)70065-2
- Perico, L., Benigni, A., and Remuzzi, G. (2020). Should COVID-19 Concern Nephrologists? Why and to what Extent? the Emerging Impasse of Angiotensin Blockade. *Nephron* 144 (5), 213–221. doi:10.1159/000507305
- Perzborn, E., Roehrig, S., Straub, A., Kubitzka, D., and Misselwitz, F. (2011). The Discovery and Development of Rivaroxaban, an Oral, Direct Factor Xa Inhibitor. *Nat. Rev. Drug Discov.* 10 (1), 61–75. doi:10.1038/nrd3185
- Podder, C. S., Chowdhury, N., Sina, M. I., and Haque, W. M. M. U. (2020). Outcome of Ivermectin Treated Mild to Moderate COVID-19 Cases: a single-centre, Open-Label, Randomised Controlled Study. *IMC J. Med. Sci.* 14 (2), 11–18. doi:10.3329/imcjs.v14i2.52826
- Polack, F. P., Thomas, S. J., Kitchin, N., Absalon, J., Gurtman, A., Lockhart, S., et al. (2020). Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N. Engl. J. Med.* 383 (27), 2603–2615. doi:10.1056/NEJMoa2034577
- Reilly, J. (2020). Treatment Considerations for Coronavirus (COVID-19). *Hosp. Pract.* (1995) 48 (3), 119–120. doi:10.1080/21548331.2020.1754618
- Ridwan, R. (2020). Remdesivir and COVID-19: Justified in Emergency Use Authorization? *Authorea Preprints*. doi:10.22541/au.158999392.29347277
- Rizzo, E. (2020). Ivermectin, Antiviral Properties and COVID-19: a Possible New Mechanism of Action. *Naunyn Schmiedebergs Arch. Pharmacol.* 393, 1153–1156. doi:10.1007/s00210-020-01902-5
- Rothan, H. A., and Byrareddy, S. N. (2020). The Epidemiology and Pathogenesis of Coronavirus Disease (COVID-19) Outbreak. *J. Autoimmun.* 109, 102433. doi:10.1016/j.jaut.2020.102433
- Rynes, R. I. (1997). Antimalarial Drugs in the Treatment of Rheumatological Diseases. *Br. J. Rheumatol.* 36 (7), 799–805. doi:10.1093/rheumatology/36.7.799
- Saha, A., Sharma, A. R., Bhattacharya, M., Sharma, G., Lee, S. S., and Chakraborty, C. (2020). Probable Molecular Mechanism of Remdesivir for the Treatment of COVID-19: Need to Know More. *Arch. Med. Res.* 51 (6), 585–586. doi:10.1016/j.arcmed.2020.05.001
- Saqrane, S., El Mhammedi, M. A., Lahrich, S., Laghrib, F., El Bouabi, Y., Farahi, A., et al. (2021). Recent Knowledge in Favor of Remdesivir (GS-5734) as a Therapeutic Option for the COVID-19 Infections. *J. Infect. Public Health* 14, 655–660. doi:10.1016/j.jiph.2021.02.006
- Sargiacomo, C., Sotgia, F., and Lisanti, M. P. (2020). COVID-19 and Chronological Aging: Senolytics and Other Anti-aging Drugs for the Treatment or Prevention of corona Virus Infection? *Aging (Albany NY)* 12 (8), 6511–6517. doi:10.18632/aging.103001
- Savarino, A., Boelaert, J. R., Cassone, A., Majori, G., and Cauda, R. (2003). Effects of Chloroquine on Viral Infections: an Old Drug against Today's Diseases?. *Lancet Infect. Dis.* 3 (11), 722–727. doi:10.1016/S1473-3099(03)00806-5
- Savarino, A., Di Trani, L., Donatelli, I., Cauda, R., and Cassone, A. (2006). New Insights into the Antiviral Effects of Chloroquine. *Lancet Infect. Dis.* 6 (2), 67–69. doi:10.1016/S1473-3099(06)70361-9
- Shang, J., Wan, Y., Luo, C., Ye, G., Geng, Q., Auerbach, A., et al. (2020). Cell Entry Mechanisms of SARS-CoV-2. *Proc. Natl. Acad. Sci. U S A.* 117 (21), 11727–11734. doi:10.1073/pnas.2003138117
- Shippey, E. A., Wagler, V. D., and Collamer, A. N. (2018). Hydroxychloroquine: An Old Drug with New Relevance. *Cleve Clin. J. Med.* 85 (6), 459–467. doi:10.3949/ccjm.85a.17034
- Sigrist, C. J., Bridge, A., and Le Mercier, P. (2020). A Potential Role for Integrins in Host Cell Entry by SARS-CoV-2. *Antivir. Res.* 177, 104759. doi:10.1016/j.antiviral.2020.104759
- Singh, R., Rathore, S. S., Khan, H., Bhurwal, A., Sheraton, M., Ghosh, P., et al. (2021). Mortality and Severity in COVID-19 Patients on ACEIs & ARBs-A Meta-Regression Analysis. *medRxiv*. doi:10.1101/2021.03.14.21253557
- Singh, T. U., Parida, S., Lingaraju, M. C., Kesavan, M., Kumar, D., and Singh, R. K. (2020). Drug Repurposing Approach to Fight COVID-19. *Pharmacol. Rep.* 72, 1479–1508. doi:10.1007/s43440-020-00155-6
- Smith, T., Bushek, J., LeClaire, A., and Prosser, T. (2020). *COVID-19 Drug Therapy*. Amsterdam, Netherlands: Elsevier.
- Sodhi, M., and Etminan, M. (2020). Therapeutic Potential for Tetracyclines in the Treatment of COVID-19. *Pharmacotherapy* 40 (5), 487–488. doi:10.1002/phar.2395
- Spyropoulos, A. C., Ageno, W., and Barnathan, E. S. (2020). Hospital-based Use of Thromboprophylaxis in Patients with COVID-19. *Lancet* 395 (10234), e75. doi:10.1016/S0140-6736(20)30926-0
- Szolnoky, G. (2020). Further Aspects of Doxycycline Therapy in COVID-19. *Dermatol. Ther.* 33 (6), e13810. doi:10.1111/dth.13810
- Tang, N., Li, D., Wang, X., and Sun, Z. (2020). Abnormal Coagulation Parameters Are Associated with Poor Prognosis in Patients with Novel Coronavirus Pneumonia. *J. Thromb. Haemost.* 18 (4), 844–847. doi:10.1111/jth.14768
- Tang, W., Cao, Z., Han, M., Wang, Z., Chen, J., Sun, W., et al. (2020). Hydroxychloroquine in Patients with Mainly Mild to Moderate Coronavirus Disease 2019: Open Label, Randomised Controlled Trial. *bmj* 369, m1849. doi:10.1136/bmj.m1849
- Tian, C., Li, N., Bai, Y., Xiao, H., Li, S., Ge, Q. G., et al. (2021). Angiotensin Converting Enzymes Inhibitors or Angiotensin Receptor Blockers Should Be Continued in COVID-19 Patients with Hypertension. *World J. Clin. Cases* 9 (1), 47–60. doi:10.12998/wjcc.v9.i1.47
- Touret, F., and de Lamballerie, X. (2020). Of Chloroquine and COVID-19. *Antivir. Res.* 177, 104762. doi:10.1016/j.antiviral.2020.104762
- Touret, F., Gilles, M., Barral, K., Nougairède, A., van Helden, J., Decroly, E., et al. (2020). *In Vitro* screening of a FDA Approved Chemical Library Reveals Potential Inhibitors of SARS-CoV-2 Replication. *Sci. Rep.* 10 (1), 13093–13098. doi:10.1038/s41598-020-70143-6
- Tsuzuki, S., Hayakawa, K., Matsunaga, N., Terada, M., Suzuki, S., Ohtsu, H., et al. (2021). Efficacy of Remdesivir in Japanese Patients Hospitalised with COVID-19: A Large Observational Study Using the COVID-19 Registry Japan. *medRxiv*. doi:10.1101/2021.03.09.21253183
- Udwadia, Z. F., Singh, P., Barkate, H., Patil, S., Rangwala, S., Pendse, A., et al. (2021). Efficacy and Safety of Favipiravir, an Oral RNA-dependent RNA Polymerase Inhibitor, in Mild-To-Moderate COVID-19: A Randomized, Comparative, Open-Label, Multicenter, Phase 3 Clinical Trial. *Int. J. Infect. Dis.* 103, 62–71. doi:10.1016/j.ijid.2020.11.142

- Uçan, A., Çerçi, P., Efe, S., Akgün, H., Özmen, A., Yağmuroğlu, A., et al. (2021). Benefits of Treatment with Favipiravir in Hospitalized Patients for COVID-19: a Retrospective Observational Case-Control Study. doi:10.21203/rs.3.rs-175340/v1
- Vijayvargiya, P., Esquer Garrigos, Z., Castillo Almeida, N. E., Gurram, P. R., Stevens, R. W., and Razonable, R. R. (2020). Treatment Considerations for COVID-19: A Critical Review of the Evidence (Or Lack Thereof). *Mayo Clin. Proc.* 95 (7), 1454–1466. doi:10.1016/j.mayocp.2020.04.027
- Vincent, M. J., Bergeron, E., Benjannet, S., Erickson, B. R., Rollin, P. E., Ksiazek, T. G., et al. (2005). Chloroquine Is a Potent Inhibitor of SARS Coronavirus Infection and Spread. *Viol. J.* 2 (1), 69–10. doi:10.1186/1743-422X-2-69
- Wahl, A., Gralinski, L. E., Johnson, C. E., Yao, W., Kovarova, M., Dinno, K. H., et al. (2021). SARS-CoV-2 Infection Is Effectively Treated and Prevented by EIDD-2801. *Nature* 591, 451–457. doi:10.1038/s41586-021-03312-w
- Waldrop, T., Alsup, D., and McLaughlin, E. C. (2020). Fearing Coronavirus, Arizona Man Dies after Taking a Form of Chloroquine Used to Treat Aquariums, CNN Health. Available at: <https://edition.cnn.com/2020/03/23/health/arizona-coronavirus-chloroquine-death/index.html> (Accessed January 25, 2020).
- Wang, F., Chen, C., Tan, W., Yang, K., and Yang, H. (2016). Structure of Main Protease from Human Coronavirus NL63: Insights for Wide Spectrum Anticoronavirus Drug Design. *Sci. Rep.* 6 (1), 22677. doi:10.1038/srep22677
- Wang, M., Cao, R., Zhang, L., Yang, X., Liu, J., Xu, M., et al. (2020). Remdesivir and Chloroquine Effectively Inhibit the Recently Emerged Novel Coronavirus (2019-nCoV) *In Vitro*. *Cell Res.* 30 (3), 269–271. doi:10.1038/s41422-020-0282-0
- WHO Coronavirus Disease (COVID-19) Dashboard (2020). Available at: <https://COVID19.who.int/> (Accessed January 25, 2021).
- WHO (2020). WHO Recommends against the Use of Remdesivir in COVID-19 Patients. Available at: <https://www.who.int/news-room/feature-stories/detail/who-recommends-against-the-use-of-remdesivir-in-COVID-19-patients> (Accessed January 25, 2021).
- Wouters, O. J., Shadlen, K. C., Salcher-Konrad, M., Pollard, A. J., Larson, H. J., Teerawattananon, Y., et al. (2021). Challenges in Ensuring Global Access to COVID-19 Vaccines: Production, Affordability, Allocation, and Deployment. *The Lancet* 397, 1023–1034. doi:10.1016/S0140-6736(21)00306-8
- Wu, P. E., and Morris, A. M. (2021). Remdesivir for Patients with COVID-19. *CMAJ* 193 (4), E125. doi:10.1503/cmaj.202505
- Wu, R., Wang, L., Kuo, H. D., Shannar, A., Peter, R., Chou, P. J., et al. (2020). An Update on Current Therapeutic Drugs Treating COVID-19. *Curr. Pharmacol. Rep.* 6 (3), 1–15. doi:10.1007/s40495-020-00216-7
- Wysocki, J., Lores, E., Ye, M., Soler, M. J., and Batlle, D. (2020). Kidney and Lung ACE2 Expression after an ACE Inhibitor or an Ang II Receptor Blocker: Implications for COVID-19. *bioRxiv*. doi:10.1101/2020.05.20.106658
- Yang, X., Yu, Y., Xu, J., Shu, H., Xia, J., Liu, H., et al. (2020). Clinical Course and Outcomes of Critically Ill Patients with SARS-CoV-2 Pneumonia in Wuhan, China: a Single-Centered, Retrospective, Observational Study. *Lancet Respir. Med.* 8(5), 475–481. doi:10.1016/S2213-2600(20)30079-5
- Yehualashet, A. S., and Belachew, T. F. (2020). ACEIs and ARBs and Their Correlation with COVID-19: a Review. *Infect. Drug Resist.* 13, 3217–3224. doi:10.2147/IDR.S264882
- Zeeshan Khan Chachar, A., Ahmad Khan, K., Asif, M., Tanveer, K., Khaqan, A., and Basri, R. (2020). Effectiveness of Ivermectin in SARS-CoV-2/COVID-19 Patients. *Int. J. Sci.* 9 (09), 31–35. doi:10.18483/ijsci.2378
- Zhang, G., Wu, Y., Xu, R., and Du, X. (2021). Effects of Renin-Angiotensin-Aldosterone System Inhibitors on Disease Severity and Mortality in Patients with COVID-19: A Meta-Analysis. *J. Med. Virol.* 93 (4), 2287–2300. doi:10.1002/jmv.26695
- Zhang, P., Zhu, L., Cai, J., Lei, F., Qin, J. J., Xie, J., et al. (2020). Association of Inpatient Use of Angiotensin-Converting Enzyme Inhibitors and Angiotensin II Receptor Blockers with Mortality Among Patients with Hypertension Hospitalized with COVID-19. *Circ. Res.* 126 (12), 1671–1681. doi:10.1161/CIRCRESAHA.120.317134
- Zhou, F., Yu, T., Du, R., Fan, G., Liu, Y., Liu, Z., et al. (2020). Clinical Course and Risk Factors for Mortality of Adult Inpatients with COVID-19 in Wuhan, China: a Retrospective Cohort Study. *Lancet* 395 (10229), 1054–1062. doi:10.1016/S0140-6736(20)30566-3
- Zhou, P., Yang, X. L., Wang, X. G., Hu, B., Zhang, L., Zhang, W., et al. (2020). A Pneumonia Outbreak Associated with a New Coronavirus of Probable Bat Origin. *nature* 579 (7798), 270–273. doi:10.1038/s41586-020-2012-7
- Zhou, Y., Agudelo, J., Lu, K., Goetz, D. H., Hansell, E., Chen, Y. T., et al. (2011). Inhibitors of SARS-CoV Entry-Identification Using an Internally-Controlled Dual Envelope Pseudovirus Assay. *Antivir. Res.* 92 (2), 187–194. doi:10.1016/j.antiviral.2011.07.016
- Zhou, Y., Vedantham, P., Lu, K., Agudelo, J., Carrion, R., Jr, Nunneley, J. W., et al. (2015). Protease Inhibitors Targeting Coronavirus and Filovirus Entry. *Antivir. Res.* 116, 76–84. doi:10.1016/j.antiviral.2015.01.011

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Alam, Kamal, Sarker, Zhou, Rahman and Mohamed. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.