

Corrigendum: Ginsenoside Compound K Enhances Fracture Healing *via* **Promoting Osteogenesis and Angiogenesis**

Lingli Ding¹, Song Gu², Bingyu Zhou¹, Min Wang¹, Yage Zhang¹, Siluo Wu¹, Hong Zou³, Guoping Zhao^{4,5,6,7,8,9}*, Zhao Gao¹⁰* and Liangliang Xu¹*

¹Key Laboratory of Orthopaedics and Traumatology, Lingnan Medical Research Center, The First Affiliated Hospital of Guangzhou University of Chinese Medicine, Guangzhou University of Chinese Medicine, Guangzhou, China, ²The First Affiliated Hospital, Guizhou University of Traditional Chinese Medicine, Guiyang, China, ³Engineering Laboratory for Nutrition, Shanghai Institute of Nutrition and Health, Chinese Academy of Sciences, Shanghai, China, ⁴Master Lab for Innovative Application of Nature Products, National Center of Technology Innovation for Synthetic Biology, Tianjin Institute of Industrial Biotechnology, Chinese Academy of Sciences, Tianjin, China, ⁵CAS Key Laboratory of Quantitative Engineering Biology, Shenzhen Institute of Synthetic Biology, Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences, Shanghai Institute of Plant Physiology and Ecology, Chinese Academy of Sciences, Shanghai, China, ⁷Bio-Med Big Data Center, Shanghai Institute of Nutrition and Health, Chinese Academy of Sciences, Shanghai, China, ⁸State Key Laboratory of Genetic Engineering, Department of Microbiology and Immunology, School of Life Sciences, Fudan University, Shanghai, China, ⁹Department of Microbiology, The Chinese University of Hong Kong, Hong Kong, Hong Kong SAR, China, ¹⁰Er Sha Sports Training Center of Guangdong Province, Guangzhou, China

OPEN ACCESS

Edited and reviewed by:

Longhuo Wu, Gannan Medical University, China

*Correspondence:

Guoping Zhao gpzhao@sibs.ac.cn Zhao Gao qiqin1990@163.com Liangliang Xu xull-2016@gzucm.edu.cn

Specialty section:

This article was submitted to Ethnopharmacology, a section of the journal Frontiers in Pharmacology

Received: 25 May 2022 Accepted: 30 May 2022 Published: 16 June 2022

Citation:

Ding L, Gu S, Zhou B, Wang M, Zhang Y, Wu S, Zou H, Zhao G, Gao Z and Xu L (2022) Corrigendum: Ginsenoside Compound K Enhances Fracture Healing via Promoting Osteogenesis and Angiogenesis. Front. Pharmacol. :952598. doi: 10.3389/fphar.2022.952598 Keywords: CK, fracture healing, Wnt/β -catenin, osteogenesis, angiogenesis

A Corrigendum on

Ginsenoside Compound K Enhances Fracture Healing via Promoting Osteogenesis and Angiogenesis

by Ding L, Gu S, Zhou B, Wang M, Zhang Y, Wu S, Zou H, Zhao G, Gao Z and Xu L (2022). Front. Pharmacol. 13:855393. doi: 10.3389/fphar.2022.855393

In the original article, there was a mistake in **Figure 3A** as published. **Figure 3A** is a schematic diagram of the time points of animal modeling and samples, in which the model and time points were incorrectly marked. The corrected **Figure 3** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Ding, Gu, Zhou, Wang, Zhang, Wu, Zou, Zhao, Gao and Xu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.



FIGURE 3 [CK accelerated the progression of tracture healing. (A) Schematic illustration of time points of animal modeling and sample collection. (B) Representative 3-dimensional micro-CT images of femurs in each group. (C–E) Quantitative analysis of parameters, including CV, BMD, and BV/TV, *p < 0.05, compared with the PBS group, n = 3 (F–G) Biomechanical properties of the fractured bones by 3-point bending test, *p < 0.05, compared with the PBS group, n = 3 (F–G)