



OPEN ACCESS

EDITED BY

Jian Gao,
Shanghai Children's Medical Center,
China

REVIEWED BY

Yong Wang,
Beijing University of Chinese Medicine,
China
Jun Kang,
Tianjin University, China
Shulan Su,
Nanjing University of Chinese Medicine,
China

*CORRESPONDENCE

Xianchun Duan,
ahtcmdxc@163.com
Weidong Chen,
wdchen@ahtcm.edu.cn

[†]These authors have contributed equally to this work and share first authorship

SPECIALTY SECTION

This article was submitted to
Inflammation Pharmacology,
a section of the journal
Frontiers in Pharmacology

RECEIVED 27 May 2022

ACCEPTED 27 June 2022

PUBLISHED 12 August 2022

CITATION

Wang N, Chu F, Zhang L, Fei C, Yu C,
Xue S, Wang Y, Fang L, Peng D, Duan X
and Chen W (2022), Taohong siwu
decoction attenuates AIM2 and NLRC4
inflammasomes by ameliorates
deoxyribonucleic acid damage after
ischemic stroke.
Front. Pharmacol. 13:954867.
doi: 10.3389/fphar.2022.954867

COPYRIGHT

© 2022 Wang, Chu, Zhang, Fei, Yu, Xue,
Wang, Fang, Peng, Duan and Chen. This
is an open-access article distributed
under the terms of the [Creative
Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/).
The use, distribution or reproduction in
other forums is permitted, provided the
original author(s) and the copyright
owner(s) are credited and that the
original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution
or reproduction is permitted which does
not comply with these terms.

Taohong siwu decoction attenuates AIM2 and NLRC4 inflammasomes by ameliorates deoxyribonucleic acid damage after ischemic stroke

Ni Wang^{1,2,3,4,5†}, Furui Chu^{1,2,3,4,5†}, Lijuan Zhang^{1,2}, Changyi Fei^{1,2},
Chao Yu^{1,2}, Sujun Xue^{1,2}, Yongzhong Wang¹, Ling Fang⁶,
Daiyin Peng^{2,3,4}, Xianchun Duan^{1,2,3,4,5*} and Weidong Chen^{2,3,4*}

¹Department of Pharmacy, The First Affiliated Hospital of Anhui University of Traditional Chinese Medicine, Hefei, China, ²School of Pharmacy, Anhui University of Chinese Medicine, Hefei, China, ³MOE-Anhui Joint Collaborative Innovation Center for Quality Improvement of Anhui Genuine Chinese Medicinal Materials, Hefei, China, ⁴Anhui Province Key Laboratory of Chinese Medicinal Formula, Hefei, China, ⁵Key Laboratory of Xin'an Medicine (Anhui University of Chinese Medicine), Ministry of Education, Hefei, China, ⁶Department of Pharmacy, The First Affiliated Hospital of Anhui Medical University, Hefei, China

Taohong siwu decoction (THSWD) has been shown to have a therapeutic effect on ischemic strokes (IS). However, it is not clear to us whether THSWD reduces deoxyribonucleic acid (DNA) damage after stroke and reduces the inflammatory response caused by the damage. Therefore, we constructed an IS model (I/R) in rats and performed oxygen-glucose deprivation/reoxygenation (OGD/R) on BV2 cells. Then ELISA, immunofluorescence staining, immunohistochemistry staining, and RT-qPCR were performed to detect the expressions of absent in melanoma 2 (AIM2), NLRC4, and Caspase-1 inflammasomes and other inflammatory factors. Experimental stroke causes DNA damage, and we found that the aforementioned inflammasomes as well as inflammatory factors were significantly inhibited after treatment with THSWD by comparing the model group with the model administration group. In addition, we examined the expression of AIM2, NLRC4, and Caspase-1 in BV2 cells of OGD/R and found that the expression of the aforementioned inflammasomes was significantly decreased in OGD/R by administration of THSWD-containing serum. Our data suggest that THSWD can reduced DNA damage after stroke as well as the inflammatory response caused by the damage.

KEYWORDS

Taohong Siwu Decoction, ischemic stroke, AIM2, NLRC4, inflammasomes

Abbreviations: THSWD, taohong siwu decoction; IS, ischemic strokes; I/R, ischemic stroke model; OGD/R, oxygen-glucose deprivation/reoxygenation; DNA, deoxyribonucleic acid; AIM2, Absent in melanoma 2; DSB, double-strand break

Introduction

Ischemic stroke (IS) is caused by the occlusion of the arteries in the brain, causing brain damage and inflammatory response (Pluta et al., 2021). Brain damage caused by IS includes excitatory toxicity, oxidative stress, neuroinflammation, apoptosis, and other phenomena (Sekeljic et al., 2012; Radenovic et al., 2020). Damaged brain areas get progressively worse as blood flow decreases (Majid, 2014). The main principles of treatment for IS include dissolving blood clots and restoring blood supply, but blood recirculation can cause cerebral ischemia-reperfusion injury (Omoto et al., 2022). Among them, deoxyribonucleic acid (DNA) damage and inflammatory response are considered to be important pathological mechanisms of cerebral ischemia-reperfusion injury (Dasdelen et al., 2021; Franke et al., 2021).

Oxidation-induced secondary damage occurs during IS, resulting in oxidative damage to DNA (Lorente et al., 2021). Too much production and release of pro-inflammatory cytokines in response to DNA damage severely affect the ability of cells to regenerate. Among them, NLRP3 and AIM2 inflammasomes are the main inflammasomes involved in DNA damage and associated cytokine release (Cinat et al., 2021). Inflammasomes have now been shown to be key mediators in triggering inflammation after IS, and it is a novel multimeric protein complex (Mohamed et al., 2015). NLRC4 is a member of the NLR family, which activates caspases, forms inflammasomes, and participates in the regulation of inflammatory responses (Sundaram and Kanneganti, 2021). Absent in melanoma 2 (AIM2) is derived from damaged cells and can trigger an inflammatory response by recruiting caspase-1 (Zhang et al., 2020). It has been shown that AIM2 inflammasome and NLRC4 inflammasome expression is upregulated after IS. AIM2 inflammasome leads to brain damage and cognitive impairment after chronic stroke in mice (Habib et al., 2020).

Taohong Siwu Decoction (THSWD) was first recorded in “Yizong Jinjian” written by Wu Qian. It is a classic formula for activating blood circulation and resolving blood stasis, with the effect of nourishing blood circulation, eliminating blood stasis, and creating new blood (Xia et al., 2021). Zhang et al. (2018) explained the mechanism of THSWD intervention in acute blood stasis model rats based on liquid chromatography with quadrupole time-of-flight mass spectrometry (LC/Q-TOF-MS) of urine metabolomics, and they found that the intervention mechanism may be mainly related to

the regulation of lipid metabolism and amino acid metabolism. It has been shown that THSWD reduces the activation level of NLRP3 inflammatory vesicles in MCAO/R rats, downregulates GSDMD, and inhibits cellular scorching (Wang et al., 2020). Using network pharmacology, Pan et al. (2022) conducted a preliminary exploration of the mechanism of action of THSWD for IS and found that THSWD reversed the high expression levels of C1qb, C1qc, C3ar1, C5ar1, and Cfd proteins and inhibited the inflammatory response after cerebral ischemia. Our group’s previous study showed that THSWD could reverse the expression of inflammation in IS and reduced the pathological manifestations (Wang et al., 2021). However, it is not clear to us whether THSWD reduces DNA damage after stroke and reduces the inflammatory response caused by the damage. Experimental stroke causes DNA damage. We observed DNA damage by gavage of THSWD in stroke rats and assessed the effect of THSWD on AIM2 and NLRC4 inflammasomes vesicles in rat brains after stroke. In addition, to provide ideas for further studies of THSWD, we evaluated the expression and regulation of AIM2 and NLRC4 inflammasomes in BV2 cells after oxygen-glucose deprivation/reoxygenation by THSWD-containing serum.

Materials and methods

Herbal medicine and animals

The specific information on herbs is shown in Table 1. According to the proportion of herbs in Table 1, the medicinal materials were first extracted with 10 times the amount of water for 2 h and filtered, and the filtrate was stored. The filtrate was extracted with eight times the amount of water for 1.5 h, filtered, and then combined twice and concentrated (Wang et al., 2021). Fifty male Sprague–Dawley (SD) rats were obtained from the Experimental Animal Center of Anhui University of Traditional Chinese Medicine. The rats were confined in a room with appropriate temperature and humidity and given feed and water on a daily basis. All animal experiments were approved by the Animal Experiment Ethics Committee of Anhui University of Chinese Medicine (license number: LLSC20160336).

TABLE 1 Constituents of THSWD.

Components	Part used	Proportion	Batch number
Bai Shao (BS) (<i>Paeonia lactiflora</i> Pall.)	Dried root	3	17050301
Shu Dihuang (SDH) [<i>Rehmannia glutinosa</i> (Gaertn.) DC.]	Dried root	4	17042501
Dang Gui (DG) (<i>Angelica sinensis</i> (Oliv.) Diels)	Dried root	3	16070501
Chuan Xiong (CX) (<i>Conioselinum anthriscoides</i> ‘Chuanxiong’)	Dried rhizome	2	17061601
Tao Ren (TR) [<i>Prunus persica</i> (L.) Batsch.]	Dried ripe seed	3	17033101
Hong Hua (HH) (<i>Carthamus tinctorius</i> L.)	Dried flower	2	17041401

TABLE 2 Neurological function score.

Score	Symptoms seen in rats
0	Moved normally
1	The left front paw could not be successfully extended
2	Turned to the left when crawling
3	The rat's body was skewed to the left when crawling
4	Unconscious, unable to crawl

Animal model establishment and grouping

After 1 week of rats rearing, based on the literature by Longa et al. (1989), a rat middle cerebral artery occlusion model (I/R) was constructed in SD rats. The specific operations are as follows: we made a hole with scissors 1 cm from the bifurcation of the common carotid artery in the rat; inserted a fish line into the hole, approximately 18–22 mm; and partially withdrew the line after 2 h to restore blood flow and then sutured the wound. The animals were scored for neurological function within 12 h after the establishment of the animal model, and the rats were randomly divided into the model group and the drug administration group. A Zea Longa 5-point scale was used for neurological function scores. The specific scored criteria are shown in Table 2. Longa scores of 1–3 were considered appropriate for inclusion in the study.

Rats were assigned into three groups with 10 rats each: Sham, I/R, and I/R + THSWD. In the I/R + THSWD group, 9 g/(kg-d) of THSWD was continuously administered intragastrically for 7 days, and in the remaining groups, the same amount of normal saline was administered intragastrically.

Preparation of hippocampal cell suspension

The hippocampus was removed from the rat brain and put into a Petri dish containing PBS solution, and then, the hippocampus was transferred to a 1.5 ml PE tube and digested with 0.25% trypsin 1 ml in a 37°C water bath for 10 min. After the digestion was finished, serum was added, filtered through 300 mesh nylon mesh, and centrifuged at 1000 rpm for 5 min, and the precipitate was cells.

Taohong siwu decoction drug-contained serum intervention


SD rats were assigned into two groups with 10 rats each: the normal group and the drug-containing serum group. In the drug-contained serum group, 9 g/(kg-d) of THSWD was continuously administered intragastrically twice a day for 7 days, and in the normal group, the same amount of normal saline was administered intragastrically. Anesthetized rats had their blood taken from the abdominal aorta, centrifuged, and passed through the filter membrane.

Oxygen-glucose deprivation/reoxygenation model

BV2 cells were maintained in a humidified environment at 37°C with 5% CO₂. The cells were cultured in Dulbecco's modified Eagle medium (DMEM, Gibco Biotech, United States) supplemented with 10% fetal bovine serum (FBS, DMEM, Gibco Biotech, United States) and 0.5% penicillin–streptomycin (PS, DMEM, Gibco Biotech, United States). EDTA (DMEM, Gibco Biotech, United States)

TABLE 3 Primer sequence.

Primer		Sequence	Product length
Gapdh	Forward	CTCCACTCACGGCAAATTC AAC	147
	Reverse	GTAGACTCCACGACATACTCAGC	
Polg1	Forward	GATGAATGGGCCTACCTTGA	66
	Reverse	TGGGGTCTGTCTTCTACAGC	
Tert	Forward	CTAGTCATGTGTCAAGACCCTCTT	110
	Reverse	GCCAGCACGTTTCTCTCGTT	
Dloop1	Forward	CCCTTCCCCATTGGTCT	45
	Reverse	TGGTTTACGGAGGATGG	
Dloop2	Forward	CAGTCATAAACTCTTCTCT	70
	Reverse	CGGAGGATGGTAGATTAAT	
IL-1 β	Forward	AGTTGACGGACCCAAA	104
	Reverse	TCTTGTTGATGTGCTGCTG	
IL-18	Forward	GGAGGGTTGTGTCCAG	62
	Reverse	AATACAGGCGAGGTCATCA	



treatment was used to detach cells for splitting. During the logarithmic growth phase of BV2 cells, the cell medium was replaced with DMEM without glucose (90,113, Solarbio, Beijing, China). Cells were placed in anoxic chambers containing 5% CO₂ and 95% N₂, which were placed in an incubator for 1, 2, 3, 4, 6, or 8 h to establish an OGD model. At the end of OGD, the medium was replaced with high-glucose DMEM for subsequent culture in an aerobic incubator for 12 h to establish the OGD/R model in BV2 cells. Cells in the control group were always cultured under normal conditions. THSWD-contained serum was dissolved in water at various concentrations before co-incubation with OGD/R-induced BV2 cells. Based on literature reports, to explore the effective concentration of THSWD-contained serum that could improve the viability of BV2 cells, OGD/R-induced BV2 cells were co-incubated with 0, 2.5, 5, 10, 15, and 20% THSWD-contained serum, and viability was detected via MTT assay. Therefore, the drug concentration that improves the vitality of BV2 cells is selected. Cells in OGD/R + THSWD group were co-incubated with THSWD rat serum, and cells in the control (CN) group were co-

incubated with the same volume of the corresponding solvent.

Cell viability assay

The BV2 cells (1×10^4 /well) were seeded into 96-well plates for assessment of viability with an MTT kit (Beyotime, Shanghai, China).

Immunohistochemistry

The fixed rat brain tissue was paraffin-embedded, and sections were made. Then, the following were performed: dewaxing, hydration, antigen repair, closure with normal serum, addition of γ -H2AX antibody (Proteintech, 10856-1-AP, 1:200), staining using DAB, staining again where the color turns blue, gradient dehydration with alcohol, making clear blocked slices, and finally observation (400 \times , ischemic area or corresponding area) under a microscope.

Immunofluorescence stainings

Fixed samples were incubated with PBS containing 0.5% Triton X-100 for 20 min. PBS was added and washed three times, followed by blocking with 5% BSA for 30 min. Sections were then washed, and primary antibodies (AIM2, A3356, 1:100, Abclonal, CN; Caspase-1, 22915-1-AP, 1:100, Proteintech, CN; NLRC4, A7382, 1:100, Abclonal, CN) were added and then incubated overnight at 4°C. A secondary antibody (Cy3 GoatAntiRabbit IgG (H+L) (AS007, 1:200, Abclonal, CN) or FITC Goat AntiRabbit IgG (H+L) (AS011, 1:200, Abclonal, CN)) was added and incubated for 50 min at room temperature (23°C \pm 2°C), and cell nuclei were stained with DAPI.

Enzyme-linked immunosorbent assay

Blood was collected from the rat's abdominal aorta and left at room temperature for 30 min, and the supernatant was centrifuged to detect IL-1 β and IL-18 using an ELISA kit.

Quantitative real-time PCR

Total RNA was extracted from each group of rat brain tissue according to the instructions of the EZ-10 Total RNA Mini-Preps Kit, reverse transcribed into cDNA, and analyzed using the 2^{- $\Delta\Delta$ Ct} method for relative quantification. The primer sequences are shown in Table 3.

Detection of cytosolic deoxyribonucleic acid

After the I/R and ODG/R ended, total DNA and cytosolic DNA were extracted from hippocampal and BV2 cells and assayed based on the literature by Yang et al. (2021). The primers for gDNA are Tert and Plog1, and the primers for mtDNA are Dloop1 and Dloop2.

Statistical analysis

SPSS and GraphPad software were used to analyze the data. Normal distributed measures were expressed as mean \pm standard deviation, and one-way ANOVA and *t*-test were used. $p < 0.05$ was statistically significant.

Results

Neurological functional scores

After the rats were modeled, all rats in all groups except the Sham group had significant neurological deficits, indicating successful modeling, as shown in Figure 1.

Taohong siwu decoction-contained serum increased the viability of BV2 cells following induction by oxygen-glucose deprivation/reoxygenation

To establish a suitable OGD/R model of BV2 cells, BV2 cells were first exposed to OGD for 1, 2, 3, 4, 5, 6, or 8 h and then

reoxygenated for 12 h. The MTT assay results (Figure 2A) revealed that OGD 3 h/R 12 h significantly decreased cell viability, whereas OGD 4 h/R 12 h, OGD 5 h/R 12 h OGD 6 h/R 12 h, and OGD 8 h/R 12 h more significantly decreased viability, suggesting excessive cell injury. In addition, OGD 5 h/R 12 h could inhibit nearly 50% cell viability, and we chose it as the best condition. To explore the effective concentration of THSWD-contained serum that could improve the viability of BV2 cells, OGD/R-induced BV2 cells were co-incubated with 0, 2.5, 5, 10, 15, and 20% THSWD-contained serum, and viability was detected via MTT assay (Figure 2B). Among the tested concentrations, cell viability was most obviously improved with 10% THSWD-contained serum. Therefore, we chose 10% THSWD-contained serum as an effective drug concentration for follow-up experiments.

Taohong siwu decoction reduced the expression of absent in melanoma 2, NLRC4, and Caspase-1 in the ischemic stroke model

Experimental stroke causes DNA damage, where AIM2 inflammasomes are the main inflammasomes involved in DNA damage, and NLRC4 can form inflammasomes involved in regulating host immune and inflammatory responses (Cui et al., 2000; Guo et al., 2018). AIM2 is activated by the cleavage of procaspase-1, which is cleaved to caspase-1, which is closely associated with the inflammatory form of cell death (Sun and Scott, 2016). The results are shown in Figure 3. By comparison, it was found that the expression levels of AIM2, NLRC4, and Caspase-1 were significantly inhibited in the I/R + THSWD group after the gavage of THSWD to stroke rats.




FIGURE 2

Inhibition rate and different concentrations of THSWD-containing serum. Notes: (A) inhibition rate (B) cell viability.




FIGURE 3

Expression of absent in melanoma 2 (AIM2), NLR4, and caspase-1 inflammasomes in animal models. Notes: # $p < 0.01$ vs. Sham group; ## $p < 0.01$ vs. I/R group.

Taohong siwu decoction reduced the expression of absent in melanoma 2, NLR4, and Caspase-1 in oxygen-glucose deprivation/reoxygenation-induced BV2 cells

To further evaluate the inhibitory effects of THSWD on AIM2 and NLR4 DNA recognizers and their downstream caspase-1 inflammatory factors, we constructed an OGD/R for BV2 cells and administered THSWD-containing serum for immunofluorescence assays. We found that the expression levels of AIM2, NLR4, and Caspase-1 were higher in the OGD/R group

than in the CN group. After administration of THSWD-containing serum, we found that the OGD/R + THSWD group had a significant inhibitory effect on the high expression levels of AIM2, NLR4, and Caspase-1, as shown in Figure 4.

Taohong siwu decoction reduced the release of inflammatory factors from the ischemic stroke model

Activated caspase-1 is able to activate IL-1 β and IL-18 inflammatory factors and induce their release, which in turn






FIGURE 4

Expression of AIM2, NLRP4 and caspase-1 inflammasomes in cellular models. Notes: [#] $p < 0.01$ vs. Sham group; ^{##} $p < 0.01$ vs. I/R group.

recruits other inflammatory cells and amplifies the inflammatory response (Broz and Dixit, 2016). We used ELISA to detect IL-1 β and IL-18 inflammatory factors, and the results showed that the levels of IL-1 β and IL-18 were significantly increased in the I/R group compared with those in the Sham group and decreased in the I/R + THSWD group compared with those in the I/R group, as shown in Figure 5. THSWD can prevent the release of the above indicators.

Taohong siwu decoction reduced the release of inflammatory factors from oxygen-glucose deprivation/reoxygenation-induced BV2 cells

To examine whether THSWD-contained serum could inhibit the inflammatory response of OGD/R-induced BV2 cells, qPCR was performed to detect levels of several major inflammatory



factors. The results demonstrated that the release of IL-1 β and IL-18 were significantly increased in the OGD/R group compared with those in the CN group but significantly decreased in the OGD/R + THSWD-contained serum group compared with those in the OGD/R group after treatment, as shown in Figure 6. These findings suggest that THSWD-contained serum prevented the inflammatory response of BV2 cells following induction by OGD/R.

Reduce deoxyribonucleic acid damage

γ -H2AX is a specific biomarker for characterizing DNA damage, especially DNA double-strand breaks (DSBs) (Lin et al., 2019). Although experimental stroke produces DNA damage, we established a rat stroke model and used immunohistochemistry for the detection of γ -H2AX, a marker of DNA damage. We found that the expression of γ -H2AX was clearly observed to be higher in the I/R group than in the Sham group, and after THSWD treatment, the expression of γ -H2AX was significantly inhibited in the I/R + THSWD group, as shown in Figure 7. Suggesting that THSWD can reduce stroke-induced DNA damage. To further prove that THSWD can reduce DNA damage, DNA was extracted from both animal and cellular models and examined via RT-qPCR, and the results are shown in Figures 8A,B. It can be seen that DNA damage was reduced after the administration of THSWD and THSWD-containing serum.

Discussion

In the present study, we demonstrated that THSWD attenuated mitochondrial DNA and nuclear DNA damage during stroke; reduced AIM2, NLRP4, and Caspase-1 inflammasomes; and inhibited IL-1 β and IL-18 inflammatory factor release after IS. In particular, THSWD may reduce the neuroinflammatory response after cerebral ischemia by ameliorating DNA damage after IS and inhibiting inflammatory somatization.

Injured cells can release free DNA such as mitochondrial DNA and nuclear DNA, which induce an inflammatory response by binding to receptors (Kanou et al., 2021). Inadequate blood flow supply to the brain enhances oxidative stress, resulting in DNA damage (Li et al., 2018). Experimental stroke causes DNA damage, a common precursor event for nerve cell death (Cui et al., 2000). Therefore, DNA damage has become a major focus of stroke research. γ -H2AX is a specific biomarker for DNA damage, especially DNA DSB, and the number of focal points formed by γ -H2AX corresponds to the number of DSB. It has been suggested that IS leads to a significant increase in γ -H2AX injury (Lin et al., 2019). We detected the expression of γ -H2AX in the I/R model and the accumulation of cytoplasmic DNA in I/R and OGD/R cells. It was found that γ -H2AX decreased significantly after THSWD intragastric administration in rats with IS. In the I/R and OGD/R models, there was DNA damage, and damage was reduced after administration of THSWD and THSWD-containing serum.

In most cases, activated NLRs and ALRs recruit a two-component protein called apoptosis-associated spot protein that can induce the protein hydrolysis of pro-IL-1 β and pro-IL-18 (Man and Kanneganti, 2015). AIM2 contains two typical domains of the hIN-200 protein family structurally, one of which is the N-terminal thermoprotein domain Pyrin-domain and the




FIGURE 7
Expression of γ -H2AX in each group. Notes: [#] $p < 0.01$ vs. Sham group; ^{**} $p < 0.05$ vs. I/R group.




FIGURE 8
DNA testing. Notes: (A) animal models (B) cellular models. A ^{*} $p < 0.05$, [#] $p < 0.01$ vs. Sham group; ^{**} $p < 0.05$ vs. I/R group. B ^{*} $p < 0.05$, [#] $p < 0.01$ vs. CN group; ^{**} $p < 0.05$, ^{##} $p < 0.01$ vs. oxygen-glucose deprivation/reoxygenation (OGD/R) group.

other is the C-terminal HIN domain. When abnormal double-stranded DNA is present in the cytoplasm, the HIN domain of AIM2 binds to the cytoplasm double-stranded DNA in a sequence-independent manner. AIM2 can be activated by procaspase-1 cleavage (Ravichandran and Heneka, 2021). Activated caspase-1 cleaves pro-inflammatory cytokines such as pro-IL-1 β and pro-IL-18 into mature forms of IL-1 β and IL-18 and induces their release, which recruits other inflammatory cells and amplifies the inflammatory response (Broz and Dixit, 2016; Sun and Scott, 2016).

It has been shown that cerebrospinal fluid induces the activation of neurons by AIM2 inflammasomes in patients with traumatic brain injury, suggesting that AIM2 may play a potential pathogenic role in neuronal disease (Adamczak et al., 2014). In addition, the deleterious effect of AIM2 on ischemic brain injury has also been shown in rodent models of stroke (Denes et al., 2015). NLRC4 is both an NLR protein and an inflammasome activator that interacts directly with procaspase-1, which in turn mediates the cleavage of IL-1 β and IL-18 precursors, converting them into mature inflammatory factors and ultimately creating a “waterfall effect” that leads to a severe inflammatory response (Duncan and Canna, 2018). Recent studies have further highlighted the role of AIM2 and NLRC4 in post-ischemic pathophysiology (Heinisch et al., 2022). It was found that cerebral ischemia-reperfusion injury induces the activation of AIM2 and NLRC4 inflammasomes, promotes the maturation and release of inflammatory factors, amplifies the inflammatory response, and aggravates brain injury (Li et al., 2020). We established an animal stroke model and gave THSWD treatment. We detected the expression of three inflammasomes, AIM2, NLRC4, and Caspase-1 using immunofluorescence and found that the expression of the three inflammasomes was significantly decreased in stroke rats after THSWD administration. Then, we examined the expression and regulation of AIM2, NLRC4, and Caspase-1 inflammasomes in BV2 cells after OGD/R by THSWD-containing serum and found that THSWD-containing serum significantly inhibited the high expression levels of AIM2, NLRC4, and Caspase-1. Next, we examined the expression of IL-1 β and IL-18 inflammatory factors in the I/R model and the OGD/R model using ELISA and RT-qPCR and found a significant increase in the model group and a significant decrease in the administered group, indicating that THSWD can inhibit IL-1 β and IL-18 inflammatory factors.

Conclusion

In summary, our results suggest that THSWD may be effective in the treatment of IS by reducing DNA damage after IS, inhibiting the expression of AIM2, NLRC4, and Caspase-1 inflammasomes and the release of IL-1 β and IL-18 inflammatory factors. To provide an experimental and theoretical basis for further study of THSWD treatment of IS.

Data availability statement

The original contributions presented in the study are included in the article/supplementary material; further inquiries can be directed to the corresponding authors.

Ethics statement

The animal study was reviewed and approved by the Animal Experiment Ethics Committee of Anhui University of Chinese Medicine (license number: LLSC20160336).

Author contributions

XD, WC, and DP conceived and designed the study. FC, LZ, and CF performed the *in vitro* experiments. NW wrote the manuscript, and CY, SX, YW, and LF provided ideas for the experimental design and modified the manuscripts to ensure the integrity of the entire experimental design. All authors read and approved the final manuscript.

Funding

This research was supported by the National Natural Science Fund Regional Innovation and Development Joint Fund Project (No. U19A2009), National Natural Science Foundation of China (No. 82074059), Anhui University Collaborative Innovation Project (No. GXXT-2019-043), the Anhui Provincial College Natural Science Research Key Project (No.KJ2019A0466, No. KJ2020A0409), Excellent and Top Talents Program in Colleges and Universities (No. gxyq2019034), Anhui Provincial Key Laboratory of Traditional Chinese Medicine Compounds (No. 2019AKLCMF03), and Key Laboratory of Xin'an Medicine (Anhui University of Chinese Medicine), Ministry of Education, NO. 2022XAYX07.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors, and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Adamczak, S. E., de Rivero Vaccari, J. P., Dale, G., Brand, F. J., 3rd, Nonner, D., Bullock, M. R., et al. (2014). Pyroptotic neuronal cell death mediated by the AIM2 inflammasome. *J. Cereb. Blood Flow. Metab.* 34 (4), 621–629. doi:10.1038/jcbfm.2013.236
- Broz, P., and Dixit, V. M. (2016). Inflammasomes: Mechanism of assembly, regulation and signalling. *Nat. Rev. Immunol.* 16 (7), 407–420. doi:10.1038/nri.2016.58
- Cinat, D., Coppes, R. P., and Barazzuol, L. (2021). DNA damage-induced inflammatory microenvironment and adult stem cell response. *Front. Cell Dev. Biol.* 9, 729136. doi:10.3389/fcell.2021.729136
- Cui, J., Holmes, E. H., Greene, T. G., and Liu, P. K. (2000). Oxidative DNA damage precedes DNA fragmentation after experimental stroke in rat brain. *Faseb J.* 14 (7), 955–967. doi:10.1096/fasebj.14.7.955
- Dasdelen, D., Solmaz, M., Menevse, E., Mogulkoc, R., Baltaci, A. K., Erdogan, E., et al. (2021). Increased apoptosis, tumor necrosis factor- α , and DNA damage attenuated by 3', 4'-dihydroxyflavonol in rats with brain Ischemia-reperfusion. *Indian J. Pharmacol.* 53 (1), 39–49. doi:10.4103/ijp.IJP_727_20
- Denes, A., Coutts, G., Lénárt, N., Cruickshank, S. M., Pelegrin, P., Skinner, J., et al. (2015). AIM2 and NLR4 inflammasomes contribute with ASC to acute brain injury independently of NLRP3. *Proc. Natl. Acad. Sci. U. S. A.* 112 (13), 4050–4055. doi:10.1073/pnas.1419090112
- Duncan, J. A., and Canna, S. W. (2018). The NLR4 inflammasome. *Immunol. Rev.* 281 (1), 115–123. doi:10.1111/imr.12607
- Franke, M., Bieber, M., Kraft, P., Weber, A. N. R., Stoll, G., Schuhmann, M. K., et al. (2021). The NLRP3 inflammasome drives inflammation in ischemia/reperfusion injury after transient middle cerebral artery occlusion in mice. *Brain Behav. Immun.* 92, 223–233. doi:10.1016/j.bbi.2020.12.009
- Guo, Q., Wu, Y., Hou, Y., Liu, Y., Liu, T., Zhang, H., et al. (2018). Cytokine secretion and pyroptosis of thyroid follicular cells mediated by enhanced NLRP3, NLRP1, NLR4, and AIM2 inflammasomes are associated with autoimmune thyroiditis. *Front. Immunol.* 9, 1197. doi:10.3389/fimmu.2018.01197
- Habib, P., Harms, J., Zendedel, A., Beyer, C., and Slowik, A. (2020). Gonadal hormones E2 and P mitigate cerebral ischemia-induced upregulation of the AIM2 and NLR4 inflammasomes in rats. *Int. J. Mol. Sci.* 21 (13), E4795. doi:10.3390/ijms21134795
- Heinisch, O., Zeyen, T., Goldmann, T., Prinz, M., Huber, M., Jung, J., et al. (2022). Erythropoietin abrogates post-ischemic activation of the NLRP3, NLR4, and AIM2 inflammasomes in microglia/macrophages in a TAK1-dependent manner. *Transl. Stroke Res.* 13 (3), 462–482. doi:10.1007/s12975-021-00948-8
- Kanou, T., Nakahira, K., Choi, A. M., Yeung, J. C., Cypel, M., Liu, M., et al. (2021). Cell-free DNA in human *ex vivo* lung perfusate as a potential biomarker to predict the risk of primary graft dysfunction in lung transplantation. *J. Thorac. Cardiovasc. Surg.* 162 (2), 490–499. e492. doi:10.1016/j.jtcvs.2020.08.008
- Li, P., Stetler, R. A., Leak, R. K., Shi, Y., Li, Y., Yu, W., et al. (2018). Oxidative stress and DNA damage after cerebral ischemia: Potential therapeutic targets to repair the genome and improve stroke recovery. *Neuropharmacology* 134 (Pt B), 208–217. doi:10.1016/j.neuropharm.2017.11.011
- Li, Q., Cao, Y., Dang, C., Han, B., Han, R., Ma, H., et al. (2020). Inhibition of double-strand DNA-sensing cGAS ameliorates brain injury after ischemic stroke. *EMBO Mol. Med.* 12 (4), e11002. doi:10.15252/emmm.201911002
- Lin, K. C., Chen, K. H., Wallace, C. G., Chen, Y. L., Ko, S. F., Lee, M. S., et al. (2019). Combined therapy with hyperbaric oxygen and melatonin effectively reduce brain infarct volume and preserve neurological function after acute ischemic infarct in rat. *J. Neuropathol. Exp. Neurol.* 78 (10), 949–960. doi:10.1093/jnen/nlz076
- Longa, E. Z., Weinstein, P. R., Carlson, S., and Cummins, R. (1989). Reversible middle cerebral artery occlusion without craniectomy in rats. *Stroke* 20 (1), 84–91. doi:10.1161/01.str.20.1.84
- Lorente, L., Martín, M. M., González-Rivero, A. F., Pérez-Cejas, A., Abreu-González, P., Ramos, L., et al. (2021). DNA and RNA oxidative damage are associated to mortality in patients with cerebral infarction. *Med. Intensiva* 45 (1), 35–41. doi:10.1016/j.medin.2019.07.008
- Majid, A. (2014). Neuroprotection in stroke: Past, present, and future. *ISRN Neurol.* 2014, 515716. doi:10.1155/2014/515716
- Man, S. M., and Kanneganti, T. D. (2015). Regulation of inflammasome activation. *Immunol. Rev.* 265 (1), 6–21. doi:10.1111/imr.12296
- Mohamed, I. N., Ishrat, T., Fagan, S. C., and El-Remessy, A. B. (2015). Role of inflammasome activation in the pathophysiology of vascular diseases of the neurovascular unit. *Antioxid. Redox Signal.* 22 (13), 1188–1206. doi:10.1089/ars.2014.6126
- Omoto, A. C. M., do Carmo, J. M., Nelson, B., Flynn, E., Aitken, N., Wang, Z., et al. (2022). Chronic central nervous system leptin infusion improves cardiac function and metabolism after ischemia/reperfusion injury. *FASEB J.* 36 (Suppl. 1). doi:10.1096/fasebj.2022.36.S1.R3740
- Pan, L., Peng, C., Wang, L., Li, L., Huang, S., Fei, C., et al. (2022). Network pharmacology and experimental validation-based approach to understand the effect and mechanism of Taohong Siwu Decoction against ischemic stroke. *J. Ethnopharmacol.* 294, 115339. doi:10.1016/j.jep.2022.115339
- Pluta, R., Januszewski, S., and Czuczwar, S. J. (2021). The role of gut microbiota in an ischemic stroke. *Int. J. Mol. Sci.* 22 (2), E915. doi:10.3390/ijms22020915
- Radenovic, L., Nenadic, M., Ułamek-Kozioł, M., Januszewski, S., Czuczwar, S. J., Andjus, P. R., et al. (2020). Heterogeneity in brain distribution of activated microglia and astrocytes in a rat ischemic model of Alzheimer's disease after 2 years of survival. *Aging (Albany NY)* 12 (12), 12251–12267. doi:10.18632/aging.103411
- Ravichandran, K. A., and Heneka, M. T. (2021). Inflammasome activation in neurodegenerative diseases. *Essays Biochem.* 65 (7), 885–904. doi:10.1042/ebc20210021
- Sekeljic, V., Batavejic, D., Stamenkovic, S., Ułamek, M., Jabłoński, M., Radenovic, L., et al. (2020). Cellular markers of neuroinflammation and neurogenesis after ischemic brain injury in the long-term survival rat model. *Brain Struct. Funct.* 217 (2), 411–420. doi:10.1007/s00429-011-0336-7
- Sun, Q., and Scott, M. J. (2016). Caspase-1 as a multifunctional inflammatory mediator: Noncytokine maturation roles. *J. Leukoc. Biol.* 100 (5), 961–967. doi:10.1189/jlb.3MR0516-224R
- Sundaram, B., and Kanneganti, T. D. (2021). Advances in understanding activation and function of the NLR4 inflammasome. *Int. J. Mol. Sci.* 22 (3), 1048. doi:10.3390/ijms22031048
- Wang, M., Liu, Z., Hu, S., Duan, X., Zhang, Y., Peng, C., et al. (2020). Taohong Siwu decoction ameliorates ischemic stroke injury via suppressing pyroptosis. *Front. Pharmacol.* 11, 590453. doi:10.3389/fphar.2020.590453
- Wang, N., Fei, C., Chu, F., Huang, S., Pan, L., Peng, D., et al. (2021). Taohong Siwu decoction regulates cell necrosis and neuroinflammation in the rat middle cerebral artery occlusion model. *Front. Pharmacol.* 12, 732358. doi:10.3389/fphar.2021.732358
- Xia, W., Hu, S., Wang, M., Xu, F., Han, L., Peng, D., et al. (2021). Exploration of the potential mechanism of the Tao Hong Si Wu Decoction for the treatment of postpartum blood stasis based on network pharmacology and *in vivo* experimental verification. *J. Ethnopharmacol.* 268, 113641. doi:10.1016/j.jep.2020.113641
- Yang, Y., Wu, M., Cao, D., Yang, C., Jin, J., Wu, L., et al. (2021). ZBP1-MLKL necroptotic signaling potentiates radiation-induced antitumor immunity via intratumoral STING pathway activation. *Sci. Adv.* 7 (41), eabf6290. doi:10.1126/sciadv.abf6290
- Zhang, M. J., Zhao, Q. C., Xia, M. X., Chen, J., Chen, Y. T., Cao, X., et al. (2020). The HDAC3 inhibitor RGFP966 ameliorated ischemic brain damage by downregulating the AIM2 inflammasome. *Faseb J.* 34 (1), 648–662. doi:10.1096/fj.201900394RRR
- Zhang, X., Li, P., Hua, Y., Ji, P., Yao, W., Ma, Q., et al. (2018). Urinary metabolomics study the mechanism of Taohong Siwu Decoction intervention in acute blood stasis model rats based on liquid chromatography coupled to quadrupole time-of-flight mass spectrometry. *J. Chromatogr. B Anal. Technol. Biomed. Life Sci.* 1074–1075, 51–60. doi:10.1016/j.jchromb.2017.12.035