Check for updates

OPEN ACCESS

EDITED AND REVIEWED BY Shu Wang, Beijing Tiantan Hospital, Capital Medical University, China

*CORRESPONDENCE Da-Yong Xia, ⊠ xiady1981@163.com Xiang-Sheng Zhang, ⊠ zhangxssp@163.com

RECEIVED 06 December 2024 ACCEPTED 03 July 2025 PUBLISHED 14 July 2025

CITATION

Zhang X-H, Peng L, Zhang J, Dong Y-P, Wang C-J, Liu C, Xia D-Y and Zhang X-S (2025) Correction: Berberine ameliorates subarachnoid hemorrhage injury via induction of Sirtuin 1 and inhibiting HMGB1/Nf-KB pathway. *Front. Pharmacol.* 16:1540873. doi: 10.3389/fphar.2025.1540873

COPYRIGHT

© 2025 Zhang, Peng, Zhang, Dong, Wang, Liu, Xia and Zhang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Correction: Berberine ameliorates subarachnoid hemorrhage injury via induction of Sirtuin 1 and inhibiting HMGB1/Nf-κB pathway

Xiang-Hua Zhang¹, Lei Peng¹, Jing Zhang¹, Yi-Peng Dong¹, Cheng-Jun Wang¹, Cang Liu¹, Da-Yong Xia^{2*} and Xiang-Sheng Zhang^{1*}

¹Department of Neurosurgery, Beijing Friendship Hospital, Capital Medical University, Beijing, China, ²Department of Neurosurgery, The First Affiliated Hospital of Wannan Medical College (Yijishan Hospital of Wannan Medical College), Wuhu, China

KEYWORDS

subarachnoid hemorrhage, inflammation, berberine, HMGB1, Sirtuin 1

A Correction on

Berberine ameliorates subarachnoid hemorrhage injury via induction of Sirtuin 1 and inhibiting HMGB1/Nf- κ B pathway

by Zhang X-H, Peng L, Zhang J, Dong Y-P, Wang C-J, Liu C, Xia D-Y and Zhang X-S (2020). Front. Pharmacol. 11:1073. doi: 10.3389/fphar.2020.01073

There was a mistake in Figure 4A as published, concerning the NF-kB p65 protein band. It appears that the protein band was improperly cropped during the figure preparation. Furthermore, the Western blot image included in Figure 4A does not align with the complete Western blot image. The corrected Figure 4 appears below.

The original version of this article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.



FIGURE 4

Effects of berberine treatment on SIRT1/HMGB1/TLR4/Myd88/Nf- κ B signaling pathway after SAH. (A–F) Western blot assay (A) and quantification for the expression of SIRT1 (B), HMGB1 (C), TLR4 (D), Myd88 (E), and Nf- κ B p65 (F) in the indicated groups. n = 6 per group. (G) Representative photomicrographs of immunofluorescence staining for Nf- κ B p65. Bars represent the mean \pm SD. **P* < 0.05.