



OPEN ACCESS

EDITED AND REVIEWED BY
Stefania Tacconelli,
University of Studies G. d'Annunzio Chieti and
Pescara, Italy

*CORRESPONDENCE

Shu Han,
✉ han00shu@zju.edu.cn

[†]These authors have contributed equally to
this work

RECEIVED 13 April 2025

ACCEPTED 30 April 2025

PUBLISHED 13 May 2025

CITATION

Chen H, Fu X, Jiang J and Han S (2025)
Corrigendum: C16 peptide promotes vascular
growth and reduces inflammation in a
neuromyelitis optica model.
Front. Pharmacol. 16:1611115.
doi: 10.3389/fphar.2025.1611115

COPYRIGHT

© 2025 Chen, Fu, Jiang and Han. This is an
open-access article distributed under the terms
of the [Creative Commons Attribution License](#)
(CC BY). The use, distribution or reproduction in
other forums is permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication in this
journal is cited, in accordance with accepted
academic practice. No use, distribution or
reproduction is permitted which does not
comply with these terms.

Corrigendum: C16 peptide promotes vascular growth and reduces inflammation in a neuromyelitis optica model

Haohao Chen^{1†}, Xiaoxiao Fu^{2†}, Jinzhan Jiang¹ and Shu Han^{2*}

¹Medical Molecular Biology Laboratory, School of Medicine, Jinhua Polytechnic, Jinhua, China, ²Institute of Anatomy and Cell Biology, Medical College, Zhejiang University, Hangzhou, China

KEYWORDS

C16 peptide, receptor tyrosine kinase Tie2, PI3K/Akt pathways, $\alpha\text{v}\beta 3$ integrin, neuromyelitis optica (NMO)

A Corrigendum on

C16 peptide promotes vascular growth and reduces inflammation in a neuromyelitis optica model

by Chen H, Fu X, Jiang J and Han S (2019). *Front. Pharmacol.* 10:1373. doi: 10.3389/fphar.2019.01373

In the published article, there was an error in **Supplementary Figure S1** as published. The magnification of subfigure B and G of **Supplementary Figure S1** is different from others. The corrected **Supplementary Figure S1** and its caption “Immunofluorescence images of Tie2 expression in transverse sections through the anterior horn of the lumbar SC in (A–I) normal and NMO rats treated with (C, G) C16, (D, H) Tie2 KI + C16 and (E, I) LY294002 + C16 at 3 and 8 weeks P.I. Scale bar = 100 μm . (J) Plot of Tie2+ blood vessels. a, $P < 0.05$ versus normal group; b, $P < 0.05$ versus the vehicle-treated group; c, $P < 0.05$ versus the C16-treated group; d, $P < 0.05$ versus Tie2 KI + C16 group. Tie2 loss in the ON of vehicle-treated NMO rats is indicated by the arrow in B, F.” has been updated in the original article.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.