



OPEN ACCESS

EDITED BY

Mohamed Abdo Rizk,
Mansoura University, Egypt

REVIEWED BY

Isabella Salzer,
Medical University of Vienna, Austria
Andy R. Eugene,
Osawatomie State Hospital, United States

*CORRESPONDENCE

Bader Alyahya,
✉ balyahya@ksu.edu.sa

RECEIVED 22 May 2025

ACCEPTED 21 July 2025

PUBLISHED 30 July 2025

CITATION

Alyahya B, Alalshaikh A, Almohawes M, Alnowiser M, Alsuliman O, Alrefaei R, Alaidarous S, Alnahdi M, Tamur S, Alfaifi M, Al Deeb M and Al Aseri ZA (2025) The clinical value of screening for acetaminophen in all patients with intentional overdose or altered mental status suspected to be secondary to overdose.

Front. Pharmacol. 16:1633548.

doi: 10.3389/fphar.2025.1633548

COPYRIGHT

© 2025 Alyahya, Alalshaikh, Almohawes, Alnowiser, Alsuliman, Alrefaei, Alaidarous, Alnahdi, Tamur, Alfaifi, Al Deeb and Al Aseri. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

The clinical value of screening for acetaminophen in all patients with intentional overdose or altered mental status suspected to be secondary to overdose

Bader Alyahya ^{1*}, Abdulaziz Alalshaikh ¹,
Mohammed Almohawes ¹, Mosaed Alnowiser ¹,
Omar Alsuliman ², Rand Alrefaei ², Sarah Alaidarous ²,
Maha Alnahdi ¹, Shadi Tamur ³, Musa Alfaifi ⁴,
Mohammed Al Deeb ⁵ and Zohair A. Al Aseri ^{1,6,7}

¹Emergency Medicine Department, College of Medicine, King Saud University, Riyadh, Saudi Arabia,

²College of Medicine, King Saud University, Riyadh, Saudi Arabia, ³Department of Pediatrics, College of Medicine, Taif University, Taif, Saudi Arabia, ⁴Department of Emergency Medicine, Armed Forces Hospital, Khams Mushait, Saudi Arabia, ⁵Department of Emergency Medicine, College of Medicine, King Saud Bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia, ⁶Department of Clinical Sciences, College of Medicine and Riyadh Hospital, Dar Al Uloom University, Riyadh, Saudi Arabia, ⁷Adult Critical Care, Therapeutic Deputyship, Ministry of Health, Riyadh, Saudi Arabia

Background: Acetaminophen (APAP) is commonly coingested in cases of suicide or intoxication because it is widely available, effectively analgesic and antipyretic, and it is often combined with other medications, such as opioids and antihistamines. APAP overdose often causes no symptoms or nonspecific symptoms in the first 12–24 h after ingestion. Delayed diagnosis is associated with a reduced response to antidote and sometimes liver failure and mortality. However, ordering unnecessary test is not cost-effective specially if it is mostly negative, and empirical therapy is associated with significant cost and possible adverse effects.

Methods: This single-center retrospective study was conducted at King Saud University Medical City (KSUMC) in Riyadh, Saudi Arabia. Our population included all patients who presented to the emergency department with intentional drug overdose or altered mental status (AMS) suspected to be related to an overdose between June 2015 and January 2024 but with a history that was not suggestive of APAP overdose. Medical records were reviewed for patient information on demographic data, overdose details, and documentation of the clinical features of toxicity. All the subjects were kept anonymous; we used code numbers as identifiers. The data collected by the investigators were entered into an Excel worksheet in an encrypted format.

Results: A total of 2914 patients were screened for acetaminophen (APAP) levels and 1517 met our inclusion criteria. Fourteen (0.9%) patients had detectable levels (>10 µg/mL) despite a negative history. Three (0.2%) patients had levels above 100 µg/mL and were treated with N-acetylcysteine (NAC).

Conclusion: Our study revealed that a small number of patients who presented with intentional overdose but denied APAP ingestion or AMS suspected to be due

to overdose had a positive APAP level. But, Given the serious consequences of APAP toxicity we cannot recommend stopping the screening for APAP specially in high-risk suicidal patients. A larger multicenter study is recommended to identify those high-risk patients. We also found deviation from the current guidelines regarding NAC administration in patients with positive APAP level when the time of ingestion is unknown.

KEYWORDS

acetaminophen level, acetaminophen toxicity, Saudi Arabia, acetaminophen screening, silent paracetamol overdose

1 Introduction

Acetaminophen (APAP) overdose has been recognized as one of the most common toxicities in Saudi Arabia and worldwide (Alghafees et al., 2022; Chidiac et al., 2023). Easy accessibility, over-the-counter availability, and low cost are some of the main factors leading to this issue. APAP is widely used as an effective analgesic and antipyretic agent (Chidiac et al., 2023). APAP toxicity has been identified as a major cause of fulminant liver failure. Despite being considered a safe medication, APAP is often used for self-harm and attempted suicide (Yoon et al., 2016).

Diagnosing APAP toxicity heavily relies on the patient's history and serum concentration, as the initial presentation may be vague, and patients are often asymptomatic in the first 24 h (Rowden et al., 2006). Owing to the fear of its serious complications, many physicians acquire a serum level, even if the patient denies taking it (Walls et al., 2022). A number of studies have reported that hardly any patient to have a positive acetaminophen level without any suggestive history among those presenting with intentional drug overdose or an altered mental status (AMS) likely related to poisoning (Bentur et al., 2011; Ashbourne et al., 1989; Sporer and Khayam-Bashi, 1996; Ala et al., 2019; Dargan et al., 2001; Hartington et al., 2002; Eldridge and Holstege, 2006). However, many of these studies recommend the use of an APAP level in that population given its low cost and the significant consequences of liver toxicity and, in some cases, mortality (Ashbourne et al., 1989; Hartington et al., 2002; Eldridge, 2007; Graham et al., 2006). However, the test is not always available and sometimes physicians request the test but then do not follow the existing consensus guidelines regarding treatment (Dart et al., 2023). Also, in peripheral hospitals, when APAP screening is not available, patients may need to be transferred a long distance just to obtain an APAP level or be treated empirically.

The necessity of routine measurement of APAP levels in patients with intentional drug overdose or AMS is a topic of debate in the medical community. Some studies suggest that routine measurement of APAP levels in these patients may not significantly affect the outcome (Ala et al., 2019; Dargan et al., 2001). Several studies have explored the issue of unnecessary ordering of diagnostic tests in various healthcare settings and studied emergency physicians' perceptions of medically unnecessary investigations. They reported that physicians often

order these tests because of concerns about malpractice and patient satisfaction rather than clinical necessity (Alalshaikh et al., 2022; Kanzaria et al., 2015).

An extensive literature review revealed no studies in our region evaluating the need for APAP level in patients with intentional drug overdose who denied APAP ingestion or those with AMS (therefore cannot provide history) suspected to be secondary to a drug overdose.

2 Methods

We conducted a single-center retrospective observational study at King Saud University Medical City (KSUMC) in Riyadh, Saudi Arabia, reviewing 2914 cases from 1 June 2015, to 1 January 2024, as our electronic health record system (Esihi) was implemented in 2015. This study assessed the clinical implications of APAP serum levels in patients presenting to the emergency department with suspected drug overdose. The study included all patients whose APAP level was ordered for them and then categorized into individuals who either denied APAP ingestion or those whose ingestion status was uncertain due to AMS or unreliable histories.

Patients under 14 years of age (as patients 14 years and above are seen in the adult emergency department), documented accidental APAP ingestion, those admitted to taking an APAP overdose, patients who received IV APAP in the hospital, or cases where documentation was inadequate were excluded.

We categorized APAP serum levels into two thresholds: a serum concentration of more than 10 µg/mL was considered a positive level (Dart et al., 2023), and a very high level was defined as any level exceeding 100 µg/mL.


Data were extracted from electronic health records and included whether N-acetylcysteine (NAC) was administered during the emergency visit and focused on patient demographics, overdose specifics, APAP serum levels, and liver toxicity markers, including transaminases and PT/INR levels.

To maintain subject anonymity, we used code numbers as identifiers, and all collected data were stored in an encrypted Excel sheet accessible only to the authors. This ensured strict confidentiality throughout the data extraction and analysis process.

2.1 Statistical analysis

The data were extracted and revised in an Excel sheet. Statistical analysis was conducted via the Statistical Program for Social

Abbreviations: APAP, N-acetyl-p-aminophenol (acetaminophen); AMS, Altered mental status; KSUMC, King Saud University Medical City; NAC, N-acetylcysteine; µg/mL, microgram/mL.



Sciences (SPSS) computer program (version 26.0; Armonk, NY, United States). Categorical variables are presented as numbers and percentages. Continuous, nonnormally distributed variables are reported as the means (SDs), medians, and interquartile ranges (IQRs). The chi-square test and Fisher's exact test were used to compare categorical variables. P-values less than 0.05 were considered statistically significant.

3 Results

As shown in Figure 1, 2914 patients presented to the ED of KSUMC between 1 June 2015, and 1 January 2024, and had APAP level measured. Of these, 1397 did not meet the inclusion criteria and were excluded.

A total of 1517 patients underwent acetaminophen (APAP) level screening and met our inclusion criteria. Males constituted 51.7% of the population. The median (IQR) age was 26 years. Approximately 42.5% of the studied patients had a history of psychiatric illness, whereas 23.9% had a history of street drug use. Notably, 8.5% patients had both a history of psychiatric illness and street drug use. Most of the included patients (84.5%) presented with intentional overdose but denied APAP ingestion, and the remainder presented with AMS suspected to be secondary to an overdose or poisoning. (Table 1).

Most included patients presented with negative APAP levels ≤ 10 $\mu\text{g/mL}$ ($n = 1503$, 99.1%). Fourteen patients had positive APAP levels; eleven patients (0.7%) had APAP levels between 11 and 46.5 $\mu\text{g/mL}$, whereas three patients (0.2%) had very high levels exceeding 100 $\mu\text{g/mL}$. One patient presented with a level of >300 $\mu\text{g/mL}$ and one had a level of 151 $\mu\text{g/mL}$, both are considered very high and fall above the treatment threshold regardless of the time of ingestion. Patients with APAP levels >100 $\mu\text{g/mL}$ ($n = 3$, 0.2%) received NAC treatment (Table 2). Table 3 summarizes the clinical characteristics of three individuals with markedly elevated acetaminophen (APAP) levels.

Patients with AMS who were not able to report APAP ingestion were more likely to present with detectable levels than were those who reported negative APAP ingestion (2.6% vs 0.6%, $P = 0.013$). No other factors, including age, sex, or history of having a psychiatric illness or street drug use, were significantly associated with detectable APAP levels (Table 4).

4 Discussion

Intentional drug overdose is a common presentation to emergency departments. Screening for acetaminophen overdose is often part of universal testing in all poisoned patients. APAP testing is not always available, and patients may need to be transferred to a different healthcare facility and sometimes to a different city to obtain the APAP level or receive empirical treatment. There is always a risk when transferring patients, especially at night, and there is also a significant cost associated with that (Hernandez-Boussard et al., 2017; Yu et al., 2022; Mueller et al., 2019). The main objective of this study was to assess the clinical value of ordering APAP levels in patients presenting with an intentional drug overdose when they deny APAP ingestion or those with AMS suspected to be secondary to a drug overdose.

Our results support the findings of previous studies on this topic, as only 0.9% of the studied population was found to have a positive APAP level (>10 $\mu\text{g/mL}$), and only three of the 1517 patients (0.19%) had very high APAP levels and were treated with NAC. However, based on the standard practice which is supported by a recent consensus guideline by Dart et al., all 14 patients should probably have undergone NAC treatment because the time of ingestion is unknown (Dart et al., 2023). Ashbourne and colleagues demonstrated that among 486 patients who presented to their urban hospital with intentional drug overdose, seven patients had unrecognized acetaminophen overdose. However, only one patient was potentially hepatotoxic level and needed to be treated with NAC (Ashbourne et al., 1989).

Lucanie et al. evaluated the incidence of APAP overdose among patients presenting with suicidal ingestion. Among the 320 documented patients who denied acetaminophen exposure, 23 patients had levels above 10 $\mu\text{g/mL}$. NAC was given to twelve patients, seven of whom were identified to have toxic levels. Meanwhile, the other five had non-toxic or uninterpretable levels (Lucanie et al., 2002). Dargan and colleagues screened 411 patients who had a history of either drug overdose ($n = 296$) or AMS ($n = 115$). They reported that four patients (3.5%) in the AMS group had positive levels and were treated with NAC. All those who denied APAP ingestion were not found to have detectable levels (136 of 296) (Dargan et al., 2001). In contrast, several other studies have failed to find any patient with a potentially toxic level (Sporer and Khayam-Bashi, 1996; Ala et al., 2019; Hartington et al., 2002; Graham et al., 2006; Chan et al., 1995; Garner, 1996).

Among the three patients with very high levels, patient no. 1 was a 17-year-old male known to have sickle cell disease on regular oral folic acid, hydroxyurea, and oral opioids. He was brought by an ambulance after his mother found him lying on the floor. She was uncertain whether he ingested any medication or not, and no history could be obtained from the altered patient. He was found to have an APAP level >300 $\mu\text{g/mL}$.

TABLE 1 Demographics and clinical characteristics of patients with APAP levels.

Variables		Number	Percentage
Sex	Male	785	51.7
	Female	732	48.3
Age	Mean (SD)	29.98 (13.177)	
	Median (IQR)	26 (15)	
Marital status	Single	1122	74
	Married	395	26
History of psychological illness	Yes	645	42.5
	No	872	57.5
Type of psychological illness	Depression	293	19.3
	Psychosis	232	15.3
	Bipolar	70	4.6
	Anxiety	50	3.3
Previous suicide attempt	Yes	258	17
	No	1259	83
History of Street Drug Use	Yes	362	23.9
	No	1155	76.1
Type of street drug used	Amphetamine	152	10
	Benzodiazepines	97	6.4
	Pregabalin	73	4.8
	Cocaine	30	1.9
	Not documented	12	0.8
Presentation	Suspected or confirmed intentional overdose with negative history of APAP ingestion	1282	84.5
	AMS	235	15.5

Abbreviations: AMS: altered mental status; APAP: N-acetyl-p-aminophenol (acetaminophen).

TABLE 2 APAP level at presentation and management (n = 1517).

Variables		Number	Percentage
APAP level (µg/mL)	Very high level, likely Toxic (>100)	3	0.2
	Potentially toxic (>10–60)	11	0.7
	Undetectable (≤10)	1503	99.1
N-acetylcysteine treatment	Yes	3	0.2
	No	1514	99.8

Abbreviation: APAP: N-acetyl-p-aminophenol (acetaminophen).

Among the other two patients, patient no. 2 was a 25-year-old female known to have general anxiety disorder and borderline personality disorder who presented to the emergency department with her brother following the ingestion of 20–30 pills of 5-6 different medications. She was able to recall two of these medications (esitalopram and lamotrigine). Later, she was found to have an APAP level of 114 µg/mL, and treatment with NAC was initiated. Patient no. 3 was a 14-year-old female with a history of

previous suicidal ingestion. She was brought in by her father after he found her unconscious with wrist-cutting marks. She denied any history of ingestion. Her APAP level was found to be 151 µg/mL and received NAC treatment. None of the three patients developed acute liver injury during their hospital stay, and all three patients achieved good recovery.

Camilleri deduced in his study that a history of APAP overdose is moderately reliable and more reliable than a history of street drug

TABLE 3 Patient characteristics among individuals with markedly elevated APAP level.

Variables	Patient no. 1	Patient no. 2	Patient no. 3
Sex	Male	Female	Female
Age	17	14	23
History of psychological illness	No	No	Yes
History of Street Drug Use	No	No	No
Presentation	AMS	Suspected intentional overdose with negative history of APAP ingestion	Suspected intentional overdose with negative history of APAP ingestion
APAP level	>300	151	114.2

Abbreviations: AMS: altered mental status; APAP: N-acetyl-p-aminophenol (acetaminophen).

TABLE 4 Risk factors for having high APAP levels.

Variables		APAP level		Total	P Value
		Detectable (n = 14)	Undetectable (n = 1503)		
Age median (IQR)		23 (9)	26 (15)	1517	0.140
Sex	Female	7 (1.0%)	725 (99.0%)	732	0.895
	Male	7 (0.9%)	778 (99.1%)	785	
Weight (Kg)	Mean weight	74.5	76.2	1517	0.538
History of psychiatry illness	Yes	3 (0.5%)	642 (99.5%)	645	0.109
	No	11 (1.3%)	861 (98.7%)	872	
History of street drug use	Yes	5 (1.4%)	357 (98.6%)	362	0.342
	No	9 (0.8%)	1146 (99.2%)	1155	
Presentation	Suspected or confirmed intentional overdose with negative history of APAP ingestion	8 (0.6%)	1274 (99.4%)	1282	0.013
	AMS	6 (2.6%)	229 (97.4%)	235	

Abbreviations: AMS: altered mental status; APAP: N-acetyl-p-aminophenol (acetaminophen).

misuse (Camill and eri, 2015). Despite the rare incidences of potentially toxic APAP levels, the majority of the studies mentioned above recommended screening all patients with suspected drug overdose regardless of whether the history was suggestive. Bentur et al. concluded that overdosed patients who deny both APAP and multidrug ingestion can be considered reliable for APAP history. Therefore, they proposed that routine APAP screening in patients who denied both APAP and multidrug ingestions might not be needed, especially in healthcare centers with limited resources or where the APAP level is not readily available (Bentur et al., 2011). However, in our study, two of the fourteen patients with positive results presented with single drug ingestion and denied APAP ingestion.

4.1 Limitations

This study has several limitations. First, it was conducted at a single center, which may limit the generalizability of the findings.

Additionally, since the study was based in an urban healthcare center, the results may not accurately represent the situation in peripheral hospitals or the impact of patient transfers. This study’s retrospective design carries inherent limitations. Specifically, it is susceptible to biases related to data availability, accuracy, and potential selection bias.

4.2 Conclusion

Our study revealed that a small number of patients who presented with intentional overdose but denied APAP ingestion or AMS suspected to be due to overdose had a positive APAP level. But, Given the serious consequences of APAP toxicity we cannot recommend stopping the screening for APAP specially in high-risk suicidal patients. A larger multicenter study is recommended to identify those high-risk patients. We also found deviation from the current guidelines regarding NAC administration in patients with positive APAP level when the time of ingestion is unknown.

Data availability statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

Ethics statement

The studies involving humans were approved by King Saud University Institutional Review Board. The studies were conducted in accordance with the local legislation and institutional requirements. Written informed consent for participation was not required from the participants or the participants' legal guardians/next of kin in accordance with the national legislation and institutional requirements.

Author contributions

BA: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Project administration, Resources, Software, Supervision, Validation, Visualization, Writing – original draft, Writing – review and editing. AA: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Project administration, Software, Supervision, Validation, Writing – original draft, Writing – review and editing. MhA: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Project administration, Software, Supervision, Validation, Writing – original draft, Writing – review and editing. MsA: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Project administration, Software, Supervision, Validation, Writing – original draft, Writing – review and editing. OA: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Project administration, Software, Supervision, Validation, Writing – original draft, Writing – review and editing. RA: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Project administration, Software, Supervision, Validation, Writing – original draft, Writing – review and editing. SA: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Project administration, Software, Supervision, Validation, Writing – original draft, Writing – review and editing. MaA: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Project administration, Software, Supervision, Validation, Writing – original draft, Writing – review and editing. ST: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Project administration, Software, Supervision, Validation, Writing – original draft, Writing – review and editing. MuA: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Project administration, Software, Supervision, Validation, Writing – original draft, Writing – review and editing.

References

Ala, A., Vahdati, S. S., Rigi, G. B., Habibollahi, P., Tajodini, S., and Jannati, K. (2019). Acetaminophen serum levels in patients with unknown poisoning and loss of consciousness. *Adv. Biosci. Clin. Med.* 7 (2), 20–24. doi:10.7575/aiaac.abcm.v7n.2p.20

and editing. MoA: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Project administration, Software, Supervision, Validation, Writing – original draft, Writing – review and editing. ZA: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Project administration, Software, Supervision, Validation, Writing – original draft, Writing – review and editing.

Funding

The author(s) declare that financial support was received for the research and/or publication of this article. The funding for this research was obtained from the Department of Emergency Medicine, College of Medicine, King Saud University. The funding body had no role in the design of the study; in the data collection, analysis, and interpretation; or in the writing of the manuscript.

Acknowledgments

We would like to thank King Saud University Medical City. We would also like to thank Evidence Clinical Research Solutions for their contribution to the data analysis. Moreover, we would like to thank all the people who participated in the research for their patience and kindness.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Generative AI statement

The author(s) declare that no Generative AI was used in the creation of this manuscript.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Alalshaikh, A., Alyahya, B., Almohawes, M., Alnowiser, M., Ghandour, M., Alyousef, M., et al. (2022). Emergency medicine physicians' views on providing unnecessary management in the emergency department. *Open Access Emerg. Med.* 14, 183–193. doi:10.2147/OAEM.S341709

- Alghafees, M. A., Abdulmomen, A., Eid, M., Alhussin, G. I., Alosaimi, M. Q., Alduhaimi, G. S., et al. (2022). Poisoning-related emergency department visits: the experience of a Saudi high-volume toxicology center. *Ann. Saudi Med.* 42 (1), 36–44. doi:10.5144/0256-4947.2022.36
- Ashbourne, J. F., Olson, K. R., and Khayam-Bashi, H. (1989). Value of rapid screening for acetaminophen in all patients with intentional drug overdose. *Ann. Emerg. Med.* 18 (10), 1035–1038. doi:10.1016/s0196-0644(89)80925-4
- Bentur, Y., Lurie, Y., Tamir, A., Keyes, D. C., and Basis, F. (2011). Reliability of history of acetaminophen ingestion in intentional drug overdose patients. *Hum. Exp. Toxicol.* 30 (1), 44–50. doi:10.1177/0960327110366784
- Camilleri, R. (2015). A meta-analysis of the reliability of the history in suspected poisoning. *J. Emerg. Med.* 48 (6), 679–684. doi:10.1016/j.jemermed.2014.12.067
- Chan, T. Y. K., Chan, A. Y. W., Ho, C. S., and Critchley, JAJH (1995). The clinical value of screening for paracetamol in patients with acute poisoning. *Hum. Exp. Toxicol.* 14 (2), 187–189. doi:10.1177/096032719501400205
- Chidiac, A. S., Buckley, N. A., Noghrehchi, F., and Cairns, R. (2023). Paracetamol (acetaminophen) overdose and hepatotoxicity: mechanism, treatment, prevention measures, and estimates of burden of disease. *Expert Opin. Drug Metab. Toxicol.* 19 (5), 297–317. doi:10.1080/17425255.2023.2223959
- Dargan, P. I., Ladhani, S., and Jones, A. L. (2001). Measuring plasma paracetamol concentrations in all patients with drug overdose or altered consciousness: does it change outcome? *Emerg. Med. J.* 18 (3), 178–182. doi:10.1136/emj.18.3.178
- Dart, R. C., Mullins, M. E., Matoushek, T., Ruha, A. M., Burns, M. M., Simone, K., et al. (2023). Management of acetaminophen poisoning in the US and Canada: a consensus statement. *JAMA Netw. Open* 6 (8), e2327739. doi:10.1001/jamanetworkopen.2023.27739
- Eldridge, D. L. (2007). New trends in diagnostic investigations in poisoned patients. *Drug Discov. Today Technol.* 4 (3-4), e103–e107. doi:10.1016/j.ddtec.2008.11.002
- Eldridge, D. L., and Holstege, C. P. (2006). Utilizing the laboratory in the poisoned patient. *Clin. Lab. Med.* 26 (1), 13–30. doi:10.1016/j.cll.2006.01.003
- Garner, A. (1996). Is routine screening for paracetamol in intentional drug ingestion of clinical value? *Emerg. Med. (N Y)*. 8 (2), 61–64. doi:10.1111/j.1442-2026.1996.tb00254.x
- Graham, C. A., Irons, A. J., and Munro, P. T. (2006). Paracetamol and salicylate testing: routinely required for all overdose patients? *Eur. J. Emerg. Med.* 13 (1), 26–28. doi:10.1097/00063110-200602000-00006
- Hartington, K., Hartley, J., and Clancy, M. (2002). Measuring plasma paracetamol concentrations in all patients with drug overdoses; development of a clinical decision rule and clinicians willingness to use it. *Emerg. Med. J.* 19 (5), 408–411. doi:10.1136/emj.19.5.408
- Hernandez-Boussard, T., Davies, S., McDonald, K., and Wang, N. E. (2017). Interhospital facility transfers in the United States: a nationwide outcomes study. *J. Patient Saf.* 13 (4), 187–191. doi:10.1097/PTS.0000000000000148
- Kanzaria, H. K., Hoffman, J. R., Probst, M. A., Caloyer, J. P., Berry, S. H., and Brook, R. H. (2015). Emergency physician perceptions of medically unnecessary advanced diagnostic imaging. *Acad. Emerg. Med.* 22 (4), 390–398. doi:10.1111/acem.12625
- Lucanie, R., Chiang, W. K., and Reilly, R. (2002). Utility of acetaminophen screening in unsuspected suicidal ingestions. *Vet. Hum. Toxicol.* 44 (3), 171–173.
- Mueller, S. K., Fiskio, J., and Schnipper, J. (2019). Interhospital transfer: transfer processes and patient outcomes. *J. Hosp. Med.* 14 (8), 486–491. doi:10.12788/jhm.3192
- Rowden, A. K., Norvell, J., Eldridge, D. L., and Kirk, M. A. (2006). Acetaminophen poisoning. *Clin. Lab. Med.* 26 (1), 49–65. doi:10.1016/j.cll.2006.01.004
- Sporer, K. A., and Khayam-Bashi, H. (1996). Acetaminophen and salicylate serum levels in patients with suicidal ingestion or altered mental status. *Am. J. Emerg. Med.* 14 (5), 443–446. doi:10.1016/S0735-6757(96)90146-1
- Walls, R., Hockberger, R., Gausche-Hill, M., Erickson, T. B., and Wilcox, S. R. (2022). *Rosen's emergency medicine - concepts and clinical practice E-book: 2-volume set*. Elsevier Health Sciences. Available online at: <https://books.google.com.sa/books?id=HMx1EAAAQBAJ>.
- Yoon, E., Babar, A., Choudhary, M., Kutner, M., and Pyrsopoulos, N. (2016). Acetaminophen-induced hepatotoxicity: a comprehensive update. *J. Clin. Transl. Hepatol.* 4 (2), 131–142. doi:10.14218/JCTH.2015.00052
- Yu, A., Jordan, S. R., Gilmartin, H., Mueller, S. K., Holliman, B. D., and Jones, C. D. (2022). "Our hands are tied until your doctor gets here": nursing perspectives on inter-hospital transfers. *J. Gen. Intern. Med.* 37 (7), 1729–1736. doi:10.1007/s11606-021-07276-5