



Do Changes in Hemodynamic Parameters Depend Upon Length of Sleep Deprivation? Comparison Between Subjects With Normal Blood Pressure, Prehypertension, and Hypertension

Joanna Słomko^{1*}, Monika Zawadka-Kunikowska¹, Sławomir Kujawski¹, Jacek J. Klawe¹, Małgorzata Tafil-Klawe², Julia L. Newton³ and Paweł Zalewski¹

¹ Department of Hygiene, Epidemiology and Ergonomics, Faculty of Health Sciences, Nicolaus Copernicus University in Toruń, Toruń, Poland, ² Department of Human Physiology, Faculty of Medicine, Nicolaus Copernicus University in Toruń, Toruń, Poland, ³ Institute for Cellular Medicine, The Medical School, Newcastle University, Newcastle upon Tyne, United Kingdom

OPEN ACCESS

Edited by:

Sara Montagnese,
Università degli Studi di Padova, Italy

Reviewed by:

Alireza Mani,
University College London,
United Kingdom
Teresa Seccia,
Università degli Studi di Padova, Italy

*Correspondence:

Joanna Słomko
jslomko@cm.umk.pl
orcid.org/0000-0002-8503-3172

Specialty section:

This article was submitted to
Chronobiology,
a section of the journal
Frontiers in Physiology

Received: 13 June 2018

Accepted: 11 September 2018

Published: 28 September 2018

Citation:

Słomko J, Zawadka-Kunikowska M, Kujawski S, Klawe JJ, Tafil-Klawe M, Newton JL and Zalewski P (2018) Do Changes in Hemodynamic Parameters Depend Upon Length of Sleep Deprivation? Comparison Between Subjects With Normal Blood Pressure, Prehypertension, and Hypertension. *Front. Physiol.* 9:1374. doi: 10.3389/fphys.2018.01374

The main objective of the study was to analyze the impact of sleep deprivation upon hemodynamic and autonomic parameters in subjects with normal blood pressure (BP) compared to prehypertension and hypertension at 24, 28, and 32 h of total sleep deprivation (TSD). Thirty volunteers, healthy men with current medical tests indicating the absence of disease took part in the study. After physical examination (basic neurological, clinical examination, echocardiography and doppler ultrasound of the renal arteries, evaluation of the autonomic nervous system) subjects were divided into three groups: I – normotensive, II – pre-hypertensive, III – hypertensive (age: 31.2 ± 2.1 vs. 33.5 ± 2.7 vs. 36.8 ± 2.7 years, $p > 0.05$; BMI: 25.2 ± 0.8 vs. 29.0 ± 1.5 vs. 26.4 ± 1.0 kg/m², $p > 0.05$). Hemodynamic and autonomic parameters were automatically measured at rest and in a tilted position with a Task Force Monitor. The Task Force Monitor consists of electrocardiography, impedance cardiography, oscillometric, and continuous BP measurement. Mixed models with random effects was applied in order to analyze the parameters' dependence on the time and the group of patients. One-way ANOVA or Kruskal–Wallis test were used to detect differences between normotensive, pre-hypertensive and hypertensive groups in each time point. In the pre-hypertensive group 28-h TSD resulted in increased vagal outflow [changes in high frequency heart rate (HR) variability, $p = 0.0189$], as evidenced by decreased HR ($p = 0.0293$). Moreover after 24-h TSD and 28-h TSD we observed changes in BP parameters. In hypertensive group, the most important changes in hemodynamic parameters: systolic blood pressure (sBP, $p = 0.0031$), diastolic blood pressure (dBP, $p = 0.0136$), cardiac output (CO, $p = 0.0439$) and changes in HR ($p = 0.0063$) after tilt test were observed after 32-h TSD. In conclusion, our results show that changes in hemodynamic parameters during sleep deprivation depend on the baseline BP and duration of TSD. What is important,

both groups reported a decrease of sBP and dBP during the TSD (pre-hypertensive group after 24, 28-h TSD; hypertensive group after 32-h TSD). In our opinion, this is the first study which considers three homogenous groups in terms of gender: only men, during different points of acute TSD: 24, 28, and 32 h of TSD in laboratory condition.

Keywords: hypertension, cardiovascular, autonomic nervous system, sleep deprivation, blood pressure

INTRODUCTION

Sleep deprivation (SD) is a growing health problem and is related with changes in lifestyle behaviors and an increased prevalence of sleep disorders (e.g., insomnia, obstructive sleep apnea) and neurological disorders. Epidemiological studies suggest a link between short sleep duration and an increased incidence of cardiovascular diseases occurring via the activation of various pathways: neural autonomic control changes, altered inflammatory response or increased oxidative stress (Tobaldini et al., 2013b, 2014, 2017). Current literature on the impact of SD upon the function of the autonomic nervous system are not conclusive. Holmes et al. (2002) indicate that 30 h of SD lowers activity of sympathetic autonomic control manifested by a decrease in heart rate (HR), with no change in parasympathetic autonomic nervous system activity. Other authors have observed that after 24-h SD, HR and blood pressure (BP) were higher compared to baseline in healthy subjects. Zhong et al. (2005) claimed that 36 h of SD increases activity of sympathetic autonomic nervous system function, with a reduction in activity of parasympathetic function part and a decrease in baroreflex sensitivity. A few studies found no changes in the autonomic profile induced by chronic SD (Muentner et al., 2000). SD studies of hypertensive subjects have shown significant increases in BP and sympathetic nervous system function after SD (Lusardi et al., 1999). What's important, compared to acute deprivation, chronic SD studies are more difficult to perform and results of these studies are not homogeneous; the studies differ in terms of study group characteristics, experimental settings, and experimental timetables.

The main objective of the study was to analyze the impact of total sleep deprivation (TSD) upon hemodynamic and autonomic parameters in subjects with normal BP, prehypertension, and hypertension at different periods of TSD. In our opinion, this is the first study which applies three homogenous groups in terms of gender: only men, during different point of acute TSD: 24, 28, and 32 h of TSD in laboratory condition.

MATERIALS AND METHODS


Study Group

Thirty volunteers, healthy men with current medical tests indicating the absence of disease (including routine laboratory tests) took part in the study. In addition to giving their voluntary consent to participation in the study, the main enrollment criteria included sex: male, no co-morbidity, no reported sleep disorders (Pittsburgh Sleep Quality Index < 5; Buysse et al., 1989), no extreme chronotype, [ratings between

14 and 21 points on the morning-evening M/E questionnaire (Torsvall and Akerstedt, 1980)]. Exclusion criteria consisted of factors that could possibly modify the response to SD: shift work, caffeine, alcohol, drugs dependence, participation in sports at competitive level, alcohol consumption within 12 h before the test, receiving any medication/supplements during the study and potential disorders of the cardiovascular system observed during the test. All potential study participants were questioned about their sleep quality, life habits and health state. Pre-test health state assessment of subjects included: basic neurological, clinical examination, evaluation of the autonomic nervous system using the Autonomic Symptom Profile (Suarez et al., 1999). After physical examination subjects were divided into three groups: I – normotensive, II – pre-hypertensive, III – hypertensive (age: 31.2 ± 2.1 years vs. 33.5 ± 2.7 years vs. 36.8 ± 2.7 years, $p > 0.05$; BMI: 25.2 ± 0.8 kg/m² vs. 29.0 ± 1.5 kg/m² vs. 26.4 ± 1.0 kg/m², $p > 0.05$) (Figure 1). Prehypertension and primary hypertension was found based on a physical examination, the collected data included symptoms that may indicate the secondary hypertension, the presence of risk factors, comorbidities, family interviews of hypertension and cardiovascular and renal diseases, and the objective one, i.e., the two measurements of BP in the sitting position during two different visits, echocardiography and doppler ultrasound of the renal arteries. The study was approved by the Ethics Committee, Ludwik Rydygier Memorial Collegium Medicum in Bydgoszcz, Nicolaus Copernicus University, Toruń; written informed consent was obtained from all of the participants.

Total Sleep Deprivation (TSD) Protocol

One week prior to the study (baseline week), the participants were requested to abstain from excessive caffeine (one caffeine-containing beverage per day at most) and alcohol consumption (<5 alcoholic beverages per week). Subjects arrived at the laboratory at 08:30 a.m. after their typical sleep at home the night before (total sleep time, TST = 432.6 ± 61.2 min). TSD began at 9 a.m. and finished at 5 p.m. after 32 h of continuous wakefulness. The examination was performed in the chronobiology laboratory while maintaining constant conditions (constant routine, temperature 22°C, humidity 60%, light < 10 lx). Light intensity control is crucial due to the known differences in HR variability (HRV) between dim and bright light conditions during sleep deprivation. Subjects ate the same meals at the same time of the day (8:00, 12:00, 15:00, 19:30). Water (100 ml) was administered at hourly intervals during the protocol. Reading, writing, talking, and playing games were allowed during the experiment. To make sure that patients did not fall asleep they were cared for by trained personnel. During the TSD, two investigators were present in the



laboratory, rotations of 12 h were organized. Additionally, the device Actigraph GT3X (Actigraph, Pensacola, FL, United States) was used during the experiment to monitor subjects' sleep deprivation (Santos-Lozano et al., 2013).

Cardiovascular Measurements

Hemodynamic [HR, BP, stroke volume (SV), cardiac output (CO), total peripheral resistance (TPR)] and autonomic parameters [low frequency (LF), high frequency (HF)] were automatically measured at rest (15 min after stabilization of the signals) and in a tilted position (10 min after stabilization of the signals) with a Task Force Monitor – TFM (CNS Systems, Gratz, Austria). Task Force Monitor® is designed for non-invasive measurements of hemodynamic parameters and consist of electrocardiography (ECG), impedance cardiography (ICG), oscillometric (oscBP), and continuous (contBP) blood pressure measurement (Fortin et al., 1998, 2006; Buszko et al., 2017).

Measurements of cardiovascular system parameters were obtained at 4 points during the study: 9 a.m. (baseline) on the first day of the study, 9 a.m. (24-h TSD), 1 p.m. (28-h TSD), 5 p.m. (32-h TSD) of the second day.

Statistical Analyses

Mixed models with random effects based on two-way ANOVA was applied in order to analyze the parameters' dependence on

the time and the group of patients. The Maximum likelihood method was applied for estimating variance parameters. The analyses were performed with R version 3.5.0 (R: library lme) and Matlab 2017b.

One-way ANOVA or Kruskal–Wallis test were used to detect differences between groups considering the group effect on cardiovascular parameters and HRV. If significant differences were observed by ANOVA, *post hoc* pair-wise comparisons were performed using Tukey's test. When significant differences were found in Kruskal–Wallis test, a multiple comparison of rank sum test was performed. The level of significance for all tests was set at $p < 0.05$. All calculations were conducted with STATISTICA 13.0 PL statistical package (StatSoft).

RESULTS

To assess the effect of time and the groups of patients in discussed parameters we applied mixed models with random effects. The estimation of the model for hemodynamic and spectral parameters were presented in the **Tables 1–3** and **Figures 2–5**. A significant changes were found after 28-h TSD in HR at the pre-hypertensive and hypertensive groups. 24-h and 28-h TSD influence BP parameter – significant changes in sBP and dBP were observed after 24-h and 28-h TSD in the pre-hypertensive and after 24-h (only sBP) and 32-h TSD in hypertensive

TABLE 1 | Hemodynamic parameters' dependence on the time and the group of patients.

Parameter	Variable	Value	SE	p-value	
HR	Intercept	52.9	3.1	0.0001	
	Group II vs. group I	9.2	4.9	ns	
	Group III vs. group I	13.1	4.6	0.0087	
	24-h TSD vs. baseline	0.2	2.8	ns	
	28-h TSD vs. baseline	6.6	2.8	0.0238	
	32-h TSD vs. baseline	5.1	2.8	ns	
	Group II: 24-h TSD	-5.5	4.5	ns	
	Group III: 24-h TSD	-1.8	4.2	ns	
	Group II: 28-h TSD	-10.0	4.5	0.0293	
	Group III: 28-h TSD	-9.6	4.2	0.0252	
	Group II: 32-h TSD	-6.5	4.5	ns	
	Group III: 32-h TSD	-8.9	4.2	0.0378	
	sBP	Intercept	117.2	2.7	0.0001
		Group II vs. group I	19.3	4.2	0.0001
Group III vs. group I		31.3	3.9	0.0001	
24-h TSD vs. baseline		1.7	2.7	ns	
28-h TSD vs. baseline		2.2	2.7	ns	
32-h TSD vs. baseline		3.6	2.7	ns	
Group II: 24-h TSD		-8.7	4.3	0.0453	
Group III: 24-h TSD		-8.7	4.0	0.0329	
Group II: 28-h TSD		-9.4	4.3	0.0304	
Group III: 28-h TSD		-3.5	4.0	ns	
Group II: 32-h TSD		-7.2	4.3	ns	
Group III: 32-h TSD		-12.2	4.0	0.0031	
dBP		Intercept	72.0	1.9	0.0001
		Group II vs. group I	15.5	2.9	0.0001
	Group III vs. group I	23.8	2.9	0.0001	
	24-h TSD vs. baseline	2.7	1.8	0.0001	
	28-h TSD vs. baseline	2.4	1.8	ns	
	32-h TSD vs. baseline	2.1	1.8	ns	
	Group II: 24-h TSD	-6.9	2.9	0.0188	
	Group III: 24-h TSD	-4.6	2.7	ns	
	Group II: 28-h TSD	-8.8	2.9	0.003	
	Group III: 28-h TSD	-3.7	2.7	ns	
	Group II: 32-h TSD	-5.2	2.9	ns	
	Group III: 32-h TSD	-6.7	2.7	0.0136	
	SV	Intercept	124.9	7.1	0.0001
		Group II vs. group I	-16.1	11.1	ns
Group III vs. group I		-31.5	10.5	0.0056	
24-h TSD vs. baseline		7.5	4.1	ns	
28-h TSD vs. baseline		-4.1	4.1	ns	
32-h TSD vs. baseline		1.6	4.1	ns	
Group II: 24-h TSD		-10.6	6.5	ns	
Group III: 24-h TSD		-2.8	6.1	ns	
Group II: 28-h TSD		0.1	6.4	ns	
Group III: 28-h TSD		4.3	6.0	ns	
Group II: 32-h TSD		-8.0	6.5	ns	
Group III: 32-h TSD		-8.3	6.0	ns	
CO		Intercept	6.2	0.5	0.0001
		Group II vs. group I	-0.1	0.8	ns
	Group III vs. group I	-0.5	0.8	ns	

(Continued)

TABLE 1 | Continued

Parameter	Variable	Value	SE	p-value
TPR	24-h TSD vs. baseline	0.4	0.5	ns
	28-h TSD vs. baseline	0.7	0.5	ns
	32-h TSD vs. baseline	0.7	0.5	ns
	Group II: 24-h TSD	-1.2	0.8	ns
	Group III: 24-h TSD	-0.3	0.7	ns
	Group II: 28-h TSD	-1.3	0.8	ns
	Group III: 28-h TSD	-0.9	0.7	ns
	Group II: 32-h TSD	-1.3	0.8	ns
	Group III: 32-h TSD	-1.7	0.5	0.0439
	Intercept	1113.7	97.1	0.0001
	Group II vs. group I	262.7	153.5	ns
	Group III vs. group I	432.8	144.0	0.0057
	24-h TSD vs. baseline	-40.8	71.6	ns
	28-h TSD vs. baseline	37.6	71.6	ns
32-h TSD vs. baseline	-77.9	71.6	ns	
Group II: 24-h TSD	118.9	113.3	ns	
Group III: 24-h TSD	-54.8	106.2	ns	
Group II: 28-h TSD	-43.6	113.3	ns	
Group III: 28-h TSD	11.9	106.2	ns	
Group II: 32-h TSD	139.9	113.3	ns	
Group III: 32-h TSD	194.2	106.2	ns	

Group I – normotensive, group II – pre-hypertensive, group III – hypertensive. TSD, total sleep deprivation. Bolded values = values which are statistically significant, $p < 0.05$.

TABLE 2 | Spectral parameters' dependence on the time and the group of patients.

Parameter	Variable	Value	SE	p-value	
LF-RRI	Intercept	1391.0	332.6	0.0001	
	Group II vs. group I	311.6	525.9	ns	
	Group III vs. group I	-642.8	493.3	ns	
	24-h TSD vs. baseline	-76.5	360.3	ns	
	28-h TSD vs. baseline	399.1	360.3	ns	
	32-h TSD vs. baseline	-387.2	360.3	ns	
	Group II: 24-h TSD	-547.1	569.7	ns	
	Group III: 24-h TSD	126.8	534.4	ns	
	Group II: 28-h TSD	-760.2	569.7	ns	
	Group III: 28-h TSD	-142.1	534.4	ns	
	Group II: 32-h TSD	-51.4	569.7	ns	
	Group III: 32-h TSD	570.1	534.4	ns	
	HF-RRI	Intercept	1189.4	583.6	0.0448
		Group II vs. group I	290.7	922.8	ns
Group III vs. group I		-752.4	865.7	ns	
24-h TSD vs. baseline		-77.8	656.5	ns	
28-h TSD vs. baseline		1572.4	656.5	0.0189	
32-h TSD vs. baseline		-150.1	656.5	ns	
Group II: 24-h TSD		-228.3	1037.9	ns	
Group III: 24-h TSD		239.4	973.7	ns	
Group II: 28-h TSD		-1987.4	1037.9	0.0591	
Group III: 28-h TSD		-1401.3	973.7	ns	
Group II: 32-h TSD		161.1	1037.9	ns	
Group III: 32-h TSD		509.9	973.7	ns	

Bolded values = values which are statistically significant, $p < 0.05$.

TABLE 3 | Magnitude of changes (delta) induced by tilt test on hemodynamic parameters' dependence on the time and the group of patients.

Parameter	Variable	Value	SE	p-value
Δ HR	Intercept	17.1	4.0	0.0001
	Group II vs. group I	-6.4	6.4	ns
	Group III vs. group I	-12.1	6.0	0.0525
	24-h TSD vs. baseline	-0.8	3.9	ns
	28-h TSD vs. baseline	-2.7	3.9	ns
	32-h TSD vs. baseline	40.9	3.9	0.0001
	Group II: 24-h TSD	-0.2	6.2	ns
	Group III: 24-h TSD	-1.0	5.8	ns
	Group II: 28-h TSD	4.4	6.2	ns
	Group III: 28-h TSD	3.5	5.8	ns
	Group II: 32-h TSD	9.1	6.2	ns
	Group III: 32-h TSD	16.3	5.8	0.0063
	Δ sBP	Intercept	18.7	8.2
Group II vs. group I		-8.3	12.9	ns
Group III vs. group I		-19.7	12.2	ns
24-h TSD vs. baseline		3.2	2.9	ns
28-h TSD vs. baseline		2.7	2.9	ns
32-h TSD vs. baseline		3.8	2.9	ns
Group II: 24-h TSD		6.4	4.6	ns
Group III: 24-h TSD		-0.2	4.3	ns
Group II: 28-h TSD		1.9	4.6	ns
Group III: 28-h TSD		-4.2	4.3	ns
Group II: 32-h TSD		0.7	4.6	ns
Group III: 32-h TSD		-0.4	4.3	ns
Δ dBP		Intercept	25.9	6.5
	Group II vs. group I	-10.9	10.3	ns
	Group III vs. group I	-18.6	9.7	ns
	24-h TSD vs. baseline	3.1	2.6	ns
	28-h TSD vs. baseline	2.8	2.5	ns
	32-h TSD vs. baseline	3.6	2.5	ns
	Group II: 24-h TSD	6.3	4.0	ns
	Group III: 24-h TSD	0.7	3.8	ns
	Group II: 28-h TSD	1.7	4.0	ns
	Group III: 28-h TSD	-2.1	3.8	ns
	Group II: 32-h TSD	0.2	4.0	ns
	Group III: 32-h TSD	-1.5	3.8	ns

Bolded values = values which are statistically significant, p < 0.05.

group. CO changes were observed in hypertensive group, after 32-h TSD. Only 28-h TSD influence autonomic parameters – that changes were observed in HF-RRI in pre-hypertensive group.

Differences between normotensive, pre-hypertensive and hypertensive groups in each time point were presented in the **Tables 4–6**. *Post hoc* analysis revealed significant differences in baseline between normotensive and hypertensive in HR, sBP, dBP, SV, TPR, HF-RRI, between normotensive and pre-hypertensive in sBP, dBP; 24-h SD between normotensive and hypertensive in HR, sBP, dBP, SV, TPR; 28-h SD between normotensive and hypertensive in sBP, dBP, mBP, HF-RRI and between pre-hypertensive and hypertensive in sBP, dBP; 32-h SD between normotensive and hypertensive in sBP, dBP, SV, TPR and between normotensive and pre-hypertensive in dBP and TPR.

DISCUSSION

The major finding of the present study is the evidence that variability of hemodynamic parameters depends upon sleep deprivation timetables – the most important changes in pre-hypertensive group were after 28-h TSD and in hypertensive group after 32-h TSD. First, in the pre-hypertensive group 28-h TSD resulted in increased vagal outflow (manifested in HF-RRI changes), as evidenced by decreased HR. Moreover after 24-h TSD and 28-h TSD we observed changes in BP parameters. Second, in the hypertensive group the most important changes in hemodynamic parameters: BP parameters, CO and changes in HR after tilt test were observed after 32-h TSD. Third, after 28-h TSD BP parameter differences between groups were observed not only between normotensive and hypertensive but also between the pre-hypertensive and hypertensive group. Moreover in the pre-hypertensive and hypertensive group we observed significant changes in tilt test: they had lower sBP increase after tilt in comparison to normotensive group. The observed changes may indicate less-effective regulatory mechanisms in these groups. Thus, we might speculate that chronic sleep deprivation might slowly lead to a pathological condition.

There are only a limited number of studies on the effect of sleep deprivation on autonomic function in subjects with primary/essential hypertension. The main response of the cardiovascular and autonomic nervous system to TSD in hypertensive patients is increased sympathetic nervous activity during the night and the following morning, leading to increased BP and HR (Lusardi et al., 1999; Klawe et al., 2002; Tafil-Klawe et al., 2007). More often in the literature is described the influence of stress factors on the homeostasis of the human body. Sleep deprivation, sleep fragmentation or shift work are an important stress factor that is associated with disruption of the natural circadian rhythm. Esler et al. (1991) noticed that higher rates of norepinephrine spillover into the cerebrovascular circulation after experimental laboratory stressor were in patients with essential hypertension than in healthy subjects, suggesting an underlying increase in central nervous system norepinephrine turnover may be the basis for the increased sympathetic outflow. In primary hypertension (particularly in younger patients), a differentiated activation of the sympathetic outflow to the heart and kidneys is present, based on measurements of norepinephrine spillover to plasma (Esler et al., 1991). Moreover, other studies show that during the mental stress test, the BP response was significantly greater in subjects with positive than with negative family histories (Widgren et al., 1992).

Despite numerous studies, the causes and mechanisms of hypertension have not been fully understood. Authors suggest that the central nervous system mechanisms underlying this disease and the channels which allow these mechanisms are funneled to the peripheral autonomic nervous system and trigger this cardiovascular disorder.

Several pathologies including hypertension, diabetes and obstructive sleep apnea have been associated with changes in carotid bodies activity (Conde et al., 2017). Increase in carotid sinus nerve activity drives excitation in medullary




FIGURE 2 | HR parameters' dependence on the time and the group of patients. Mixed models with random effects based on two-way ANOVA indicate that there was a significant effect of time ($p = 0.0238$), group ($p = 0.0087$) and both group \times time (II group after 28-h TSD; III group after 28 and 32-h TSD). Interestingly, both groups reported a decrease of HR during the TSD compared to baseline.




FIGURE 3 | CO parameters' dependence on the time and the group of patients. Mixed models with random effects based on two-way ANOVA indicate that there was a significant effect of time \times group ($p = 0.0439$). After 32-h TSD CO was significantly lower in the III group compared to baseline.

presympathetic pathways and is integrated in the brain stem to induce cardiorespiratory compensatory responses. Augmented chemoreceptor tonic drive and hypoxic ventilatory response were observed in young men with arterial hypertension. In the young normotensive subjects with a family background of hypertension and in the young spontaneously hypertensive rats, an increased sensitivity to chemoreceptor stimulation is seen before the onset of hypertension (Trzebski et al., 1982; Tafil-Klawe et al., 1985, Klawe et al., 1991). These observations

suggest that peripheral chemoreceptor overactivity plays a causal role in the development of hypertension. Increase in BP after sleep deprivation, increase in sympathetic activity following increased tonic drive from arterial chemoreceptor, in young men with normal baroreceptor activity activate arterial baroreceptor, causing reflex response – increase in vagal activity. In older patients with hypertension a decrease in baroreceptor sensitivity (resetting of baroreflex) is observed. One possible mechanism is adaptation of the baroreflex to prolonged stimulation (peripheral




FIGURE 4 | sBP parameters' dependence on the time and the group of patients. Mixed models with random effects based on two-way ANOVA indicate that there was a significant effect of group ($p = 0.0001$) and both group \times time (II group after 24, 28-h TSD; III group after 32-h TSD). What is important, both groups reported a decrease of sBP during the TSD compared to baseline.




FIGURE 5 | dBP parameters' dependence on the time and the group of patients. Mixed models with random effects based on two-way ANOVA indicate that there was a significant effect of group ($p = 0.0001$) and both group \times time (II group after 24, 28-h TSD; III group after 32-h TSD). Interestingly, both groups reported a decrease of dBP during the TSD compared to baseline.

or central). Our pre-hypertensive and hypertensive subjects were younger, so the cardiovascular response to sleep deprivation might be considered as the baroreceptor's prevention to increase BP – vagal activity.

The following physiological events, regulating sympathetic activity, should be considered in pre-hypertension and hypertension: augmented peripheral chemoreceptor drive, attenuation of hypoxic chemosensitivity with age, decrease

in baroreceptor reactivity in hypertension and in the course of biological aging, interaction between baro- and chemoreflexes (decrease in baroreceptor sensitivity in hypoxic conditions). The process of adaptation of these receptor's sensitivity, different stages of development of hypertension and family background of hypertension show a mosaic of different probable cardiovascular responses to physiological stimuli.

TABLE 4 | Hemodynamic parameters at baseline and after 24-h, 28-h, 32-h SD in normotensive, pre-hypertensive, and hypertensive groups in resting conditions.

Parameter	Time	Group I mean ± SD	Group II mean ± SD	Group III mean ± SD	I vs. II <i>p</i> -value	I vs. III <i>p</i> -value	II vs. III <i>p</i> -value
HR	Baseline	52.9 ± 6.0	62.2 ± 13.0	66.1 ± 9.0	ns	0.0101	ns
	24-h TSD	53.2 ± 6.5	56.8 ± 11.8	64.5 ± 9.7	ns	0.0262	ns
	28-h TSD	59.5 ± 19.7	58.7 ± 10.6	63.0 ± 7.2	ns	ns	ns
	32-h TSD	58 ± 8.7	60.7 ± 12.4	62.2 ± 8.2	ns	ns	ns
sBP	Baseline	116.2 ± 4.0	137.5 ± 2.1	148.5 ± 9.2	0.0352	0.0001	ns
	24-h TSD	118.9 ± 7.8	129.6 ± 14.0	141.5 ± 8.4	ns	0.0004	ns
	28-h TSD	119.3 ± 7.6	129.3 ± 12.6	147.2 ± 11.2	ns	0.0001	0.0059
	32-h TSD	120.8 ± 7.2	133.0 ± 8.4	140.0 ± 13.2	ns	0.0006	ns
dBP	Baseline	72.0 ± 5.7	87.5 ± 4.7	95.8 ± 4.6	0.0168	0.0001	ns
	24-h TSD	74.7 ± 6.3	83.3 ± 12.6	93.9 ± 3.9	ns	0.0002	ns
	28-h TSD	74.4 ± 5.4	81.1 ± 8.4	94.5 ± 5.5	ns	0.0001	0.0009
	32-h TSD	74.2 ± 6.5	84.4 ± 8.1	91.2 ± 4.3	0.0091	0.0001	ns
SV	Baseline	124.9 ± 25.0	108.9 ± 35.4	93.5 ± 15.5	ns	0.0258	ns
	24-h TSD	132.5 ± 25.1	105.8 ± 30.9	98.2 ± 11.4	ns	0.0036	ns
	28-h TSD	120.8 ± 26.0	104.9 ± 32.3	93.7 ± 16.5	ns	ns	ns
	32-h TSD	126.5 ± 20.8	102.5 ± 34.3	86.7 ± 13.3	ns	0.002	ns
CO	Baseline	6.6 ± 1.6	6.6 ± 2.0	6.1 ± 1.1	ns	ns	ns
	24-h TSD	7.1 ± 1.7	5.8 ± 1.4	6.3 ± 1.0	ns	ns	ns
	28-h TSD	7.3 ± 3.6	5.9 ± 1.4	5.9 ± 1.2	ns	ns	ns
	32-h TSD	7.3 ± 1.7	6.0 ± 1.7	5.4 ± 1.0	ns	0.015	ns
TPR	Baseline	1113.7 ± 315.6	1376.4 ± 388.6	1546.5 ± 340.8	ns	0.0317	ns
	24-h TSD	1072.9 ± 302.5	1454.6 ± 435.7	1450.8 ± 251.9	ns	0.0374	ns
	28-h TSD	1151.3 ± 378.7	1370.4 ± 295.9	1596.1 ± 407.3	ns	ns	ns
	32-h TSD	1035.7 ± 277.6	1438.4 ± 381.3	1662.8 ± 250.0	0.0319	0.0003	ns

Bolded values = values which are statistically significant, *p* < 0.05.

TABLE 5 | Spectral parameters at baseline and after 24-h, 28-h, 32-h SD in normotensive, pre-hypertensive and hypertensive groups in resting conditions.

Parameter	Time	Group I mean ± SD	Group II mean ± SD	Group III mean ± SD	I vs. II <i>p</i> -value	I vs. III <i>p</i> -value	II vs. III <i>p</i> -value
LF-RR1	Baseline	1391.1 ± 838.5	1702.6 ± 2404.5	748.2 ± 846.5	ns	ns	ns
	24-h TSD	1314.6 ± 1208.0	1079.0 ± 840.0	798.6 ± 791.0	ns	ns	ns
	28-h TSD	1790.2 ± 1300.1	1341.6 ± 1178.7	1005.3 ± 1032.9	ns	ns	ns
	32-h TSD	1003.8 ± 810.0	1264.0 ± 1036.8	931.1 ± 1023.5	ns	ns	ns
HF-RR1	Baseline	1189.4 ± 786.9	1480.1 ± 2975.1	436.9 ± 514.5	ns	0.0418	ns
	24-h TSD	1111.5 ± 1148.7	1173.9 ± 1922.9	598.5 ± 560.8	ns	ns	ns
	28-h TSD	2761.8 ± 4638.3	1065.1 ± 1634.4	608.1 ± 643.9	ns	0.0183	ns
	32-h TSD	1039.3 ± 1016.8	1491.1 ± 266.6	796.9 ± 872.9	ns	ns	ns

Bolded values = values which are statistically significant, *p* < 0.05.

Our results suggest an increase in parasympathetic dominance after TSD, which may be related to the modulation regulatory and compensatory mechanisms underlying the pathomechanism of the primary hypertension.

The current literature suggesting changes in cardiovascular and autonomic functioning during sleep deprivation is ambiguous. This is most likely because of the experimental timetables (length of SD, hours of SD per day, and total protocol duration), study population characteristics or experimental setting (laboratory conditions vs. real life models) different – often opposite reactions – are observed. Sleep deprivation does not have a commonly accepted definition including the minimum sleep length, it is not considered a disorder, therefore in the available literature the authors use a different

value: most often 24–60 h. Several experimental protocols have been carried out to assess the changes induced SD in healthy subjects. One night of sleep deprivation affects cardiovascular autonomic response causing an increase of cardiac sympathetic modulation and reduce the amplitude of cardiovascular autonomic response to gravitational stimulus. Moreover authors observed that SD modifies the immune pro-inflammatory profile, increasing the plasmatic levels of IFN- γ (Tobaldini et al., 2013a). The opposite results were obtained by Vaara et al. (2009) which suggest that 60-h TSD resulted in increased vagal outflow, as evidenced by decreased HR.

In SD studies, selection of participants to the study group seems to be significant, where the differences resulting from

TABLE 6 | Magnitude of changes (delta) induced by tilt test on hemodynamic parameters at baseline and after 24-h, 28-h, 32-h SD in normotensive, pre-hypertensive, and hypertensive groups.

Parameter	Time	Group I mean \pm SD	Group II mean \pm SD	Group III mean \pm SD	I vs. II <i>p</i> -value	I vs. III <i>p</i> -value	II vs. III <i>p</i> -value
Δ HR	Baseline	17.1 \pm 6.1	10.6 \pm 4.0	11.7 \pm 7.9	ns	ns	ns
	24-h TSD	16.3 \pm 5.7	9.6 \pm 5.0	11.2 \pm 8.9	ns	ns	ns
	28-h TSD	18.7 \pm 7.0	12.3 \pm 6.8	13.0 \pm 7.5	ns	ns	ns
	32-h TSD	16.6 \pm 7.8	9.8 \pm 3.9	14.0 \pm 8.0	ns	ns	ns
Δ sBP	Baseline	18.7 \pm 4.1	10.4 \pm 9.4	14.4 \pm 11.3	ns	ns	ns
	24-h TSD	22.0 \pm 9.0	20.0 \pm 8.4	16.2 \pm 9.4	ns	ns	ns
	28-h TSD	21.4 \pm 6.5	15.0 \pm 5.0	13.6 \pm 7.3	ns	0.0457	ns
	32-h TSD	22.5 \pm 4.7	14.9 \pm 5.6	16.7 \pm 6.6	0.0266	ns	ns
Δ dBP	Baseline	26.0 \pm 5.3	15.0 \pm 6.1	19.3 \pm 4.4	0.0011	0.0421	ns
	24-h TSD	29.0 \pm 9.1	24.4 \pm 9.6	22.6 \pm 10.1	ns	ns	ns
	28-h TSD	28.8 \pm 6.3	19.3 \pm 4.2	20.7 \pm 5.7	0.0022	0.032	ns
	32-h TSD	29.6 \pm 5.7	18.9 \pm 6.7	20.7 \pm 4.2	0.006	0.0094	ns

Bolded values = values which are statistically significant, $p < 0.05$.

gender seem to be the most important. For this reason, the results of our research included a homogeneous group in terms of gender, only men. Findings from the Whitehall II cohort showed gender-specific associations between sleep deprivation and hypertension. Short duration of sleep (<5 h) was associated with higher risks of hypertension only among women (Cappuccio et al., 2007). Carter et al. (2012) suggest that mechanisms underlying the acute hypertensive response to SD differ in men and women and that neural (downward shift of the sympathetic baroreflex operating point in men) and non-neural (reduced testosterone in men) mechanisms may be underlying these sex differences. Experimental settings of SD studies also have a significant influence on the study result: study carried out in shift work (night work conditions) often associated with the occurrence of environmental stress, or in experimental (laboratory) conditions. Pagani et al. (2009) suggest that one night sleep deprivation without stress or disturbances, does not lead to increased arterial pressure values or to changes in autonomic or baroreflex profiles that could conceivably favor hypertension development. Moreover they suggest that previous observations of increased arterial pressure variability levels after one night SD might rather reflect the disturbing effect of psychological stress on autonomic arousal (Pagani et al., 2009).

The results of our research taking place in laboratory show that one night sleep deprivation is too short a time to induce significant changes in hemodynamic parameters in hypertensive patients. However, in pre-hypertensive the BP changes were significant after 24-h TSD. Also, the choice of the test evaluating of the autonomic nervous system function seems to be important. The tilt test used in this experiment is characterized by a diversified response depending on the sex of the subjects – women tended to have lower tilt-table tolerance associated with a smaller splanchnic vasoconstrictor reserve than men (Jarvis et al., 2010). Comparison of the effects of orthostatic stimulation after sleep deprivation are rarely performed. Tobaldini et al. (2013a) suggest that increase of cardiac sympathetic modulation and the reduction of cardiac

vagal control induced by tilt test were significantly lowered after 26-h sleep deprivation and this suggests that one night of SD impairs the capability of the physicians to correctly respond to stressor stimuli.

In summary, based on the published evidence, acute SD is able to modify hemodynamic control and autonomic cardiovascular regulation in pre-hypertensive and hypertensive patients.

One limitation of this study was the relatively small number of participants. Most of the available SD studies were conducted on small groups, which is related to the specificity of sleep research and their high cost. The lack of an objective sleep measure, such as polysomnographic recording before the experiment (on the baseline week) and laboratory measurements (such as the aldosterone/renin ratio) must be considered as a limitation of the study. Moreover primary aldosteronism was not included in the screening of secondary hypertension.

However, the strength of the present study is twofold: first, in our opinion this is the first study which compares three homogenous groups: normotensive, pre-hypertensive and hypertensive during different time points of sleep deprivation. Secondly, the results of this study confirm that variability of hemodynamic and autonomic parameters depends upon baseline value of BP and sleep deprivation timetables. Future research is needed in standardized conditions with large-scale studies to clarify the detrimental effects of chronic SD in normotensive and hypertensive patients.

AUTHOR CONTRIBUTIONS

JS and PZ contributed conception and design of the study and wrote the first draft of the manuscript. JS, MZ-K, and SK organized the database. MZ-K performed the statistical analysis. JK, MT-K, and JN wrote the sections of the manuscript. All authors contributed to manuscript revision, read and approved the submitted version.

REFERENCES

- Buszko, K., Piątkowska, A., Kozłuk, E., and Opolski, G. (2017). Entropy in investigation of vasovagal syndrome in passive head up tilt test. *Entropy* 19:236. doi: 10.3390/e19050236
- Buyse, D., Reynolds, C. F. III, Monk, T. H., Berman, S. R., and Kupfer, D. J. (1989). The Pittsburgh sleep quality index: a new instrument for psychiatric practice and research. *Psychiatry Res.* 28, 193–213. doi: 10.1016/0165-1781(89)90047-4
- Cappuccio, F. P., Stranges, S., Kandala, N., Miller, M. A., Taggart, F. M., Kumari, M., et al. (2007). Gender-specific association of short sleep duration with prevalent and incident hypertension: the Whitehall II Study. *Hypertension* 50, 693–700. doi: 10.1161/HYPERTENSIONAHA.107.095471
- Carter, J. R., Durocher, J. J., Larson, R. A., DellaVella, J. P., and Yang, H. (2012). Sympathetic neural response to 24-hour sleep deprivation in humans: sex differences. *Am. J. Physiol. Heart Circ. Physiol.* 302, H1991–H1997. doi: 10.1152/ajpheart.01132.2011
- Conde, S. V., Monteiro, E. C., and Sacramento, J. F. (2017). Purines and carotid body: new roles in pathological conditions. *Front. Pharmacol.* 8:913. doi: 10.3389/fphar.2017.00913
- Esler, M., Ferrier, C., Lambert, G., Eisenhofer, G., Cox, H., and Jennings, G. (1991). Biochemical evidence of sympathetic hyperactivity in human hypertension. *Hypertension* 17, III29–III35. doi: 10.1161/01.HYP.17.4_Suppl. III29
- Fortin, J., Klinger, T., Wagner, C., Sterner, H., Madritsch, C., and Grullenberg, R. (1998). “The task force monitor – a non-invasive beat-to-beat monitor for hemodynamic and autonomic function of the human body,” in *Proceedings of the 20th Annual International Conference of the IEEE Engineering in Medicine and Biology Society*, Austin, TX.
- Fortin, J., Marte, W., and Grullenberg, R. (2006). Continuous non-invasive blood pressure monitoring using concentrically interlocking control loops. *Comput. Biol. Med.* 36, 941–957. doi: 10.1016/j.compbiomed.2005.04.003
- Holmes, A., Burgess, H. J., and Dawson, D. (2002). Effects of sleep pressure on endogenous cardiac autonomic activity and body temperature. *J. Appl. Physiol.* 92, 2578–2584. doi: 10.1152/jappphysiol.01106.2001
- Jarvis, S. S., Florian, J. P., Curren, M. J., and Pawelczyk, J. A. (2010). Sex differences in vasoconstrictor reserve during 70 deg head-up tilt. *Exp. Physiol.* 95, 184–193. doi: 10.1113/expphysiol.2009.048819
- Klawe, J. J., Tafil-Klawe, M., Sikorski, W., and Drzewiecka, B. (2002). Age-related circadian variations of carotid cardiac baroreflex in sleep deprived men. *J. Physiol. Pharmacol.* 53(Suppl. 1):44.
- Lusardi, P., Zoppi, A., Preti, P., Pesce, R. M., Piazza, E., and Fogari, R. (1999). Effects of insufficient sleep on blood pressure in hypertensive patients: a 24-h study. *Am. J. Hypertens.* 12, 63–68. doi: 10.1016/S0895-7061(98)00200-3
- Muenter, N. K., Watenpaugh, D. E., Wasmund, W. L., Wasmund, S. L., Maxwell, S. A., and Smith, M. L. (2000). Effect of sleep restriction on orthostatic cardiovascular control in humans. *J. Appl. Physiol.* 88, 966–972. doi: 10.1152/jappl.2000.88.3.966
- Pagani, M., Pizzinelli, P., Traon, A. P., Ferreri, C., Beltrami, S., Bareille, M. P., et al. (2009). Hemodynamic, autonomic and baroreflex changes after one night sleep deprivation in healthy volunteers. *Auton. Neurosci.* 145, 76–80. doi: 10.1016/j.autneu.2008.10.009
- Santos-Lozano, A., Santin-Medeiros, F., Cardon, G., Torres-Luque, G., Bailon, R., Bergmeir, C., et al. (2013). Actigraph GT3X: validation and determination of physical activity intensity cut points. *Int. J. Sports Med.* 34, 975–982. doi: 10.1055/s-0033-1337945
- Suarez, G. A., Opfer-Gehrking, T. L., Offord, K. P., Atkinson, E. J., O'Brien, P. C., and Low, P. A. (1999). The autonomic symptom profile a new instrument to assess autonomic symptoms. *Neurology* 52, 523–528. doi: 10.1212/WNL.52.3.523
- Tafil-Klawe, M., Klawe, J. J., Złomańczuk, P., Szczepańska, B., Sikorski, W., and Śmietanowski, M. (2007). Daily changes in cardiac and vascular blood pressure components during breath holding episodes in obstructive sleep apnea patients after day-shift and night-shift work. *J. Physiol. Pharmacol.* 58(Suppl. 5), 685–690.
- Tafil-Klawe, M., Raschke, F., Becker, H., Hein, H., Stoohs, R., Kublik, A., et al. (1991). Investigations of arterial baro- and chemoreflexes in patients with arterial hypertension and obstructive sleep apnea syndrome. *Sleep Health Risk* 319–334. doi: 10.1007/978-3-642-76034-1_38
- Tafil-Klawe, M., Trzebski, A., Klawe, J., and Palko, T. (1985). Augmented chemoreceptor reflex tonic drive in early human hypertension and in normotensive subjects with family background of hypertension. *Acta Physiol. Pol.* 36, 51–58.
- Tobaldini, E., Cogliati, C., Fiorelli, E., Nunziata, V., Wu, M., Prado, M., et al. (2013a). One night on-call: sleep deprivation affects cardiac autonomic control and inflammation in physicians. *Eur. J. Int. Med.* 24, 664–670. doi: 10.1016/j.ejim.2013.03.011
- Tobaldini, E., Costantino, G., Solbiati, M., Cogliati, C., Kara, T., Nobili, L., et al. (2017). Sleep, sleep deprivation, autonomic nervous system and cardiovascular diseases. *Neurosci. Biobehav. Rev.* 74, 321–329. doi: 10.1016/j.neubiorev.2016.07.004
- Tobaldini, E., Nobili, L., Strada, S., Casali, K. R., Braghiroli, A., and Montano, N. (2013b). Heart rate variability in normal and pathological sleep. *Front. Physiol.* 4:294. doi: 10.3389/fphys.2013.00294
- Tobaldini, E., Pecis, M., and Montano, N. (2014). Effects of acute and chronic sleep deprivation on cardiovascular regulation. *Arch. Ital. Biol.* 152, 103–110. doi: 10.12871/000298292014235
- Torsvall, L., and Akerstedt, T. (1980). A diurnal type scale. Construction, consistency and validation in shift work. *Scand. J. Work Environ. Health* 6, 283–290. doi: 10.5271/sjweh.2608
- Trzebski, A., Tafil, M., Zoltowski, M., and Przybylski, J. (1982). Increased sensitivity of the arterial chemoreceptor drive in young men with mild hypertension. *Cardiovasc. Res.* 3, 163–172. doi: 10.1093/cvr/16.3.163
- Vaara, J., Kyrolainen, H., Koivu, M., Tulppo, M., and Finni, T. (2009). The effect of 60-h sleep deprivation on cardiovascular regulation and body temperature. *Eur. J. Appl. Physiol.* 105, 439–444. doi: 10.1007/s00421-008-0921-5
- Widgren, B. R., Wikstrand, J., Berglund, G., and Andersson, O. K. (1992). Increased response to physical and mental stress in men with hypertensive parents. *Hypertension* 20, 606–611. doi: 10.1161/01.HYP.20.5.606
- Zhong, X., Hilton, J., Gates, G. J., Jelic, S., Stern, Y., Bartels, M. N., et al. (2005). Increased sympathetic and decreased parasympathetic cardiovascular modulation in normal humans with acute sleep deprivation. *J. Appl. Physiol.* 98, 2024–2032. doi: 10.1152/jappphysiol.00620.2004

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Slomko, Zawadka-Kunikowska, Kujawski, Klawe, Tafil-Klawe, Newton and Zalewski. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.