



A New Fitness Test of Estimating VO_{2max} in Well-Trained Rowing Athletes

Wei Dong Gao^{1,2}, Olli-Pekka Nuutila³, Hai Bo Fang¹, Qian Chen¹ and Xi Chen^{2*}

¹ Zhejiang Institute of Sports Science, Hangzhou, China, ² School of Sports Science, Wenzhou Medical University, Wenzhou, China, ³ Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland

Background: This study was designed to investigate the validity of maximal oxygen consumption (VO_{2max}) estimation through the Firstbeat fitness test (FFT) method when using submaximal rowing and running programs for well-trained athletes.

Methods: Well-trained flatwater rowers ($n = 45$, 19.8 ± 3.0 years, 184 ± 8.7 cm, 76 ± 12.9 kg, and 58.7 ± 6.0 mL·kg⁻¹·min⁻¹) and paddlers ($n = 45$, 19.0 ± 2.5 years, 180 ± 7.7 cm, 74 ± 9.4 kg, and 59.9 ± 4.8 mL·kg⁻¹·min⁻¹) completed the FFT and maximal graded exercise test (GXT) programs of rowing and running, respectively. The estimated VO_{2max} was calculated using the FFT system, and the measured VO_{2max} was obtained from the GXT programs. Differences between the estimated and measured VO_{2max} values were analyzed to assess the accuracy and agreement of the predictions. Equations from the previous study were also used to predict the VO_{2max} in the submaximal programs to compare the accuracy of prediction with the FFT method.

Results: The FFT method was in good agreement with the measured VO_{2max} in both groups based on the intraclass correlation coefficients (>0.8). Additionally, the FFT method had considerable accuracy in VO_{2max} estimation as the mean absolute percentage error ($\leq 5.0\%$) and mean absolute error (< 3.0 mL·kg⁻¹·min⁻¹) were fairly low. Furthermore, the FFT method seemed more accurate in the estimation of VO_{2max} than previously reported equations, especially in the rowing test program.

Conclusion: This study revealed that the FFT method provides a considerably accurate estimation of VO_{2max} in well-trained athletes.

Keywords: maximal aerobic capacity, flatwater rowers, flatwater paddlers, submaximal fitness test, treadmill running, rowing

INTRODUCTION

Maximal oxygen consumption (VO_{2max}) is defined as the maximal capacity of the pulmonary, cardiovascular, and muscular systems to deliver and utilize oxygen, which can reflect an individual's cardiorespiratory fitness (Saltin and Strange, 1992; Bassett and Howley, 1997; Levine, 2008). Measurement of VO_{2max} provides important outcomes for both physical performance and health status in general (Brink-Elfegoun et al., 2007), and it is frequently used to assess the aerobic capacity of professional or amateur athletes (Shephard, 2009). Therefore, VO_{2max} is often used in

OPEN ACCESS

Edited by:

Stefanos Volianitis,
Qatar University, Qatar

Reviewed by:

William Sheel,
University of British Columbia,
Canada
Steffen Held,
German Sport University Cologne,
Germany

*Correspondence:

Xi Chen
chenab004@126.com

Specialty section:

This article was submitted to
Exercise Physiology,
a section of the journal
Frontiers in Physiology

Received: 28 April 2021

Accepted: 10 June 2021

Published: 02 July 2021

Citation:

Gao WD, Nuutila O-P, Fang HB,
Chen Q and Chen X (2021) A New
Fitness Test of Estimating VO_{2max}
in Well-Trained Rowing Athletes.
Front. Physiol. 12:701541.
doi: 10.3389/fphys.2021.701541

endurance sports to provide training and athletes' performance information to coaches (Midgley et al., 2009; Sartor et al., 2013).

$\text{VO}_{2\text{max}}$ can be measured through direct methods with a metabolic gas measurement system, with the athlete performing a maximal graded exercise test (GXT) until exhaustion. This is regarded as the gold standard as it can obtain an accurate value of $\text{VO}_{2\text{max}}$ (Beltz et al., 2016). However, as it is time-consuming and there is an economic burden, the use of direct measurement methods is limited. In addition, the exhausting exercise program affects the training arrangement of the season (Montgomery et al., 2009; Tanner and Gore, 2013; Riebe et al., 2018). From this perspective, indirect methods of estimating $\text{VO}_{2\text{max}}$ based on the submaximal exercise program seem to be a good choice for athletes or teams, and these can frequently be used during the training season.

Previous studies have reported several indirect methods for estimating $\text{VO}_{2\text{max}}$ in athletes based on running programs (Tsiaras et al., 2010; Marsh, 2012; Matabuena et al., 2018). Marsh (2012) found a four-stage incremental running program estimating $\text{VO}_{2\text{max}}$ well, and the equation was fairly accurate [standard error of estimate (SEE) = 3.98–4.08 $\text{mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$; $r = 0.642\text{--}0.646$], as recommended by the American College of Sports Medicine. However, this equation is more suitable for males than for females, as the correlation data were conducted in male athletes. Other studies also reported submaximal $\text{VO}_{2\text{max}}$ predictive equations [(SEE) = 2.52–3.51 $\text{mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$; $r = 0.85\text{--}0.91$] for the amateur exercise population (Larsen et al., 2002; Vehrs et al., 2007).

Klusiewicz et al. (2016) developed a predictive equation ($r = 0.711$) based on the PWC170 obtained from a submaximal rowing program (Klusiewicz and Faff, 2003), which was later validated to assess the aerobic fitness of rowers. The equation based on the rowing programs had not reached an accuracy similar to that from the equation based on the running programs, with correlation coefficients of 0.55 in male rowers (Klusiewicz and Faff, 2003) and R^2 values of 0.79 and 0.64 in male and female rowers, respectively, Klusiewicz et al. (2016). Thus, the present study was designed to develop a new indirect method of $\text{VO}_{2\text{max}}$ estimation for both male and female rowing athletes.

Recently, a new system [Firstbeat fitness test (FFT)] was used for the indirect estimation of $\text{VO}_{2\text{max}}$. The estimated $\text{VO}_{2\text{max}}$ value is automatically generated after collecting heart rate (HR) data from a configurable test program (rowing or running) using Firstbeat sports software (Firstbeat, Jyväskylä, Finland). Studies have revealed that the Firstbeat $\text{VO}_{2\text{max}}$ estimation system is valid in nonathletic populations (Kraft and Roberts, 2017; Kyröläinen et al., 2018; Anderson et al., 2019; Passler et al., 2019). To the best of our knowledge, no study has reported the accuracy of this system in estimating $\text{VO}_{2\text{max}}$ in well-trained athletes. Therefore, this study was designed to investigate the accuracy of $\text{VO}_{2\text{max}}$ estimation based on the FFT system when using submaximal rowing and running programs. In addition, this study aimed to evaluate the cross-validation of previous $\text{VO}_{2\text{max}}$ predictive equations in both submaximal running (Marsh, 2012) and rowing programs (Klusiewicz et al., 2016), providing more information on the $\text{VO}_{2\text{max}}$ estimated by the FFT system to coaches and sports scientists. The hypothesis for this study was that the

FFT method could provide accurate $\text{VO}_{2\text{max}}$ estimation in well-trained athletes, which would verify a new accurate option for estimating $\text{VO}_{2\text{max}}$ in athletic population.

MATERIALS AND METHODS

The FFT system was used to estimate the participants' $\text{VO}_{2\text{max}}$ values, which were also compared to the $\text{VO}_{2\text{max}}$ values from the direct method measurement of the GXT programs. Additionally, a cross-validation design was used to evaluate the validation of the $\text{VO}_{2\text{max}}$ estimation when compared to other classical predictive equations based on submaximal rowing (Klusiewicz et al., 2016) and running programs (Marsh, 2012).

Participants

A total of 90 well-trained athletes were recruited from Zhejiang Water Sports Training Center and divided into two groups based on the sports items (i.e., rowers and paddlers): 45 flatwater rowers (23 males and 22 females) in the ROW group and 45 flatwater paddlers (29 males and 16 females) in the RUN group (participant characteristics are shown in **Table 1**). This study was conducted according to the guidelines in the Declaration of Helsinki and was approved by the ethical committee of Zhejiang Institute of Sports Science (ZJSS201909162). All participants were informed of the details of the entire program and signed an approved informed consent document.

Procedures


Figure 1 shows the flow diagram of this study. First, submaximal FFT programs were performed, the ROW group athletes performed a submaximal incremental rowing test and the RUN group athletes performed a submaximal running test. Every athlete's HR (Firstbeat, Jyväskylä, Finland) was collected during the programs, and the estimated $\text{VO}_{2\text{max}}$ value was obtained from FFT program. After 3–5 days of recovery from the submaximal program, the participants performed a maximal graded rowing (ROW group athletes) or running program (RUN group athletes) to obtain the measured $\text{VO}_{2\text{max}}$ value using a breath-by-breath metabolic measurement system, which was regarded as the golden standard test of $\text{VO}_{2\text{max}}$ (Fletcher et al., 2013). All experimental tests were performed at the same time frame during the regular training period (9:00–12:00, 15:00–18:00) in a quiet and air-conditioned laboratory (temperature, 18–23°C; humidity, 40–60%). The participants were asked to avoid heavy load training 7 days before the tests and during the recovery days, as well as abstain from caffeine and alcohol for 24 h before testing.

FFT Program

The ROW group athletes performed a submaximal incremental rowing test on the rowing ergometer (model E, Concept 2, Morrisville, VT, United States), while the RUN group athletes performed a submaximal running test on the treadmill (H/P/Cosmos, Nussdorf, Germany). According to previous studies (Marsh, 2012; Klusiewicz et al., 2016), a multistage incremental test program with up to 85% maximal HR (HR_{max}) was suggested to provide reliable predicted $\text{VO}_{2\text{max}}$.

TABLE 1 | Descriptive characteristics of the participants.

	ROW group			RUN group		
	Male (n = 23)	Female (n = 22)	All (n = 45)	Male (n = 29)	Female (n = 16)	All (n = 45)
Age (years)	20.7 ± 3.6	19.0 ± 2.1	19.8 ± 3.0	18.7 ± 2.3	19.5 ± 3.0	19.0 ± 2.5
Height (cm)	190.7 ± 5.4	176.6 ± 4.7	184 ± 8.7	184.0 ± 4.6	171.4 ± 5.0	180 ± 7.7
Body mass (kg)	84.5 ± 3.6	67.0 ± 8.9	76 ± 12.9	78.7 ± 7.0	65.7 ± 7.0	74 ± 9.4
Training experience (years)	4.7 ± 3.1	3.9 ± 2.1	4.3 ± 2.6	3.7 ± 1.9	4.6 ± 2.2	4.0 ± 2.1

**FIGURE 1** | Experimental tests diagram.

To determine the appropriate intensity for the submaximal programs, a pilot study of 10 athletes (five rowers and five paddlers) was conducted to obtain the physiological responses for different stages, especially the last stage on the rowing ergometer or running treadmill. The submaximal incremental exercise test program consisting of four 3-min rowing exercises (an initial workload of 160 W in male athletes and 120 W in female athletes) and treadmill running (an initial running speed of 9 km/h in male athletes and 8 km/h in female athletes) was designed based on the pilot study and then performed by the ROW and RUN group athletes, respectively. Detailed information on the two programs is shown in **Supplementary Table 1**. The Firstbeat HR chest belt was used to record the HR during the entire test program. All rowers completed FFT rowing program and paddlers completed FFT running program. Finally, the Firstbeat Sports software (version 4.7.3.1, Jyväskylä, Finland) automatically produced estimated VO_{2max} values based on the collected HR data of the submaximal FFT program.

GXT Program

The ROW group athletes performed a maximal incremental rowing test program (Tanner and Gore, 2013) on the Concept II rowing ergometer, with an initial workload of 160 W in males and 120 W in females. The RUN group athletes performed the Bruce running program on a treadmill (Riebe et al., 2018). Detailed information on the two maximal programs is shown in **Supplementary Table 2**. A breath-by-breath metabolic measurement system (Quark PFT Ergo, Cosmed, Rome, Italy) was used to record the respiratory gas information throughout the whole GXT procedure in both the ROW and RUN groups. The system was calibrated in advance by following the manufacturer's instructions. During the testing process, the rating of perceived exertion (RPE) on the Borg scale (6–20) and the HR (SZ990, Cosmed, Rome, Italy) were also recorded at the

end of each stage manually. The achievement of VO_{2max} was identified when meeting at least two of the three following criteria (Howley et al., 1995): (1) achievement of the oxygen consumption plateau with an increasing workload, (2) respiratory exchange ratio (RER) reached ≥ 1.10 , and (3) the HR reaching within 10 beats of the age-adjusted HR_{max} upon using the equation, $220 - \text{age}$. The value of the measured VO_{2max} was defined as the highest 30-s average value of VO_2 measured during GXT (Midgley et al., 2007).

Statistical Analysis

All data were presented as mean \pm standard deviation. The Shapiro–Wilk test was performed to test the normality of the outcome variables. Then Pearson's correlation between the estimated VO_{2max} values from the submaximal programs and the measured VO_{2max} values from the direct method using the GXT program was performed to assess the correlation magnitude and coefficient of determination (R^2). To assess the accuracy of the estimation, the mean absolute percentage error (MAPE) and mean absolute error were calculated. The intraclass correlation coefficient (ICC) was used to determine the agreement between the estimated VO_{2max} values and measured VO_{2max} values. Furthermore, the Bland-Altman plot was used to investigate the level of agreement with the 95% limits of agreement (Bland and Altman, 1986). All statistical analyses were performed using IBM SPSS Statistics software for Windows (version 24 IBM, Armonk, NY, United States), and statistical significance was set at $P < 0.05$.

RESULTS

Accuracy of VO_{2max} Estimation From FFT

Table 2 shows the VO_{2max} values measured by GXT programs and the HR, RPE, and RER in the last stage of the GXT

TABLE 2 | Results of the maximal graded exercise test.

	ROW group			RUN group		
	Male (n = 23)	Female (n = 22)	All (n = 45)	Male (n = 29)	Female (n = 16)	All (n = 45)
HR at VO _{2max} (bpm)	189.0 ± 4.5	190.8 ± 5.3	189.9 ± 4.9	198.0 ± 5.2	193.3 ± 8.5	196.3 ± 6.9
RPE at VO _{2max} (6–20)	18.2 ± 1.1	18.2 ± 0.9	18.2 ± 1.0	18.9 ± 0.8	18.8 ± 1.1	18.9 ± 0.9
RER at VO _{2max}	1.13 ± 0.1	1.11 ± 0.1	1.12 ± 0.1	1.13 ± 0.1	1.08 ± 0.1	1.11 ± 0.1
VO _{2max} (mL·kg ⁻¹ ·min ⁻¹)	60.7 ± 5.9	56.7 ± 5.5	58.7 ± 6.0	62.3 ± 3.4	55.6 ± 4.0	59.9 ± 4.8

HR, heart rate; RPE, rating of perceived exertion on Borg scale 6–20; RER, respiratory exchange ratio; and VO_{2max}, maximal oxygen consumption.

TABLE 3 | The correlations and differences between the estimated VO_{2max} from FFT and the measured VO_{2max} from GXT.

Gender	Estimated VO _{2max} (mL·kg ⁻¹ ·min ⁻¹)	Measured VO _{2max} (mL·kg ⁻¹ ·min ⁻¹)	CE (mL·kg ⁻¹ ·min ⁻¹)	t	r	ICC	MAE (mL·kg ⁻¹ ·min ⁻¹)	MAPE (%)
ROW								
Male (n = 23)	62.5 ± 6.0	60.7 ± 5.9	1.8 ± 3.6	2.429*	0.798*	0.736*	3.0 ± 2.6	5.3
Female (n = 22)	57.5 ± 6.3	56.7 ± 5.5	0.7 ± 3.3	1.047	0.851*	0.841*	2.7 ± 2.0	4.8
All (n = 45)	60.0 ± 6.0	58.7 ± 6.0	1.3 ± 3.5	2.501*	0.834*	0.818*	2.9 ± 2.3	5.0
RUN								
Male (n = 29)	63.2 ± 4.8	62.3 ± 3.4	0.9 ± 2.6	1.838	0.851*	0.787*	2.2 ± 1.6	3.5
Female (n = 16)	58.0 ± 4.4	55.6 ± 4.0	2.4 ± 2.4	3.902*	0.837*	0.727*	2.7 ± 2.0	5.0
All (n = 45)	61.3 ± 5.3	59.9 ± 4.8	1.4 ± 2.6	3.624*	0.868*	0.834*	2.4 ± 1.7	4.1

CE, constant error, arithmetic mean of the difference between estimated and measured VO_{2max}; t, t value from paired sample t-test; r, pearson correlation coefficient; ICC, intraclass correlation coefficient; MAE, mean absolute error; and MAPE, mean absolute percentage error.

*Statistically significant ($P \leq 0.05$).

program, and **Table 3** shows the estimated VO_{2max} from the FFT and the analysis of correlations and differences between the two VO_{2max} values. The results showed that the estimated VO_{2max} was significantly overestimated in both ROW [constant error (CE) = 1.3 ± 3.5 mL·kg⁻¹·min⁻¹, t [44] = 2.501, and $P = 0.016$] and RUN (CE = 1.4 ± 2.6 mL·kg⁻¹·min⁻¹, t [44] = 3.624, and $P < 0.001$) submaximal test programs (**Table 3**). However, the results of the ICC revealed that the estimated VO_{2max} from the FFT had a good level of agreement with the directly measured VO_{2max} from the GXT in both the ROW (0.818, $P < 0.001$) and RUN groups (0.834, $P < 0.001$). Additionally, the results also showed a fairly low MAPE in both the ROW and RUN groups (ROW, 5.0%; RUN, 4.1%), which also verified that the FFT was considerably accurate in VO_{2max} estimation. Furthermore, linear regression plots demonstrated a good predictive model of the FFT method in both the rowing ($R^2 = 0.695$, $P < 0.001$, and SEE = 3.35 mL·kg⁻¹·min⁻¹; **Figure 2A**) and running submaximal programs ($R^2 = 0.753$, $P < 0.001$, and SEE = 2.43 mL·kg⁻¹·min⁻¹; **Figure 2B**).



Level of Agreement Between the Estimated VO_{2max} From the FFT and the Directly Measured VO_{2max} From the GXT Program

The Bland-Altman plots also demonstrated an agreement between the estimated VO_{2max} from the FFT and the directly measured VO_{2max} from the GXT program (**Figure 3**). The findings revealed that the estimated VO_{2max} had fairly low mean

differences (bias) in both the ROW (**Figure 3A**) and RUN (**Figure 3B**) groups (bias: 1.29 mL·kg⁻¹·min⁻¹ for rowing and 1.42 mL·kg⁻¹·min⁻¹ for running). Furthermore, the FFT rowing program had a larger range of bias than that in the running program when estimating VO_{2max} from the FFT [upper to lower limits of agreement (ULoA-LLoA): 13.56 mL·kg⁻¹·min⁻¹ vs. 10.32 mL·kg⁻¹·min⁻¹, respectively].

Comparison Between the FFT Method and Previous Predictive Equations

We then examined previous equations that predict VO_{2max} based on Klusiewicz et al.'s (2016) and Marsh's (2012) studies using the HR and PWC170 (the equations and the predicted VO_{2max} are shown in **Table 4**) and compared them to the directly measured VO_{2max} from the GXT program. It was found that although Eq. 1 had a fairly accurate prediction of VO_{2max} in the rowing program, the FFT method had better accuracy and lower error terms for the overall ROW group (CE = 2.9 ± 2.3 mL·kg⁻¹·min⁻¹ and ICC = 0.818), as well as in both male (CE = 3.0 ± 2.6 mL·kg⁻¹·min⁻¹ and ICC = 0.736) and female (CE = 2.7 ± 2.0 mL·kg⁻¹·min⁻¹ and ICC = 0.841) subgroups. In addition, in the RUN group, Eqs 2, 3 had similar validity coefficients ($r = 0.655$ and 0.671, respectively) and level of agreement (ICC = 0.615 and 0.586, respectively) when the athletes performed the running program. However, in the female subgroup, neither Eq. 2 nor 3 showed a significant correlation with directly measured VO_{2max} from the GXT program. Moreover, Eq. 2 had a nonsignificant



difference [$t(44) = 1.238, P > 0.05$] and lower predictive errors ($CE = 0.7 \pm 3.7 \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$, $MAPE = 4.8\%$) than did Eq. 3.

DISCUSSION

This study aimed to investigate the accuracy of the FFT method for VO_{2max} estimation using submaximal programs, such as rowing or running, in well-trained athletes. Good levels of agreement between the estimated VO_{2max} from the FFT and the measured VO_{2max} from the GXT and $< 10\%$ MAPE were observed in the current study, which met the criteria suggested by Nelson et al. (2016). Additionally, the coefficient of determination (**Figure 2**) indicated that the FFT method accounted for 69.5% and 75.3% of the variance of VO_{2max} in the ROW and RUN groups, respectively, suggesting that the FFT method estimated VO_{2max} well. Furthermore, the

results also illustrated that FFT methods were more accurate in predicting VO_{2max} than the previous predictive equations when using the same submaximal programs in well-trained athletes. These findings indicate that the FFT method can be a fairly accurate option for obtaining VO_{2max} in well-trained athletes. Such submaximal tests can be more widely applied in the sports setting, such as in an individualized training or intervention approach, and during a repeated baseline testing setting.

Analysis of Estimated VO_{2max} Using the FFT Method

Novelty Analysis

Previous studies have developed several equations for predicting VO_{2max} when using submaximal programs in rowing or running exercise (Lakomy and Lakomy, 1993; Vehrs et al.,

TABLE 4 | Descriptive examination of the correlations and differences between other indirect methods and the measured VO_{2max} .

Methods	Reference	Predicted VO_{2max} ($mL \cdot kg^{-1} \cdot min^{-1}$)	CE ($mL \cdot kg^{-1} \cdot min^{-1}$)	<i>t</i>	<i>r</i>	ICC	MAE ($mL \cdot kg^{-1} \cdot min^{-1}$)	MAPE (%)
Equation 1	Klusiewicz et al., 2016							
	Male (<i>n</i> = 23)	65.7 ± 6.1	5.0 ± 5.6	4.310*	0.568*	0.427*	6.0 ± 4.5	10.3
	Female (<i>n</i> = 22)	59.5 ± 5.9	2.8 ± 3.5	3.701*	0.807*	0.723*	3.9 ± 2.3	6.9
	All (<i>n</i> = 45)	62.7 ± 6.7	3.9 ± 4.8	5.513*	0.721*	0.604*	5.0 ± 3.8	8.6
Equation 2	Marsh, 2012							
	Male (<i>n</i> = 29)	62.1 ± 2.8	-0.2 ± 2.8	-0.333	0.602*	0.600*	2.2 ± 1.7	3.5
	Female (<i>n</i> = 16)	57.8 ± 2.6	2.2 ± 4.5	1.954	0.125	0.099	3.6 ± 3.8	6.9
	All (<i>n</i> = 45)	60.6 ± 3.4	0.7 ± 3.7	1.238	0.655*	0.615*	2.7 ± 2.5	4.8
Equation 3	Marsh, 2012							
	Male (<i>n</i> = 29)	64.0 ± 3.5	1.7 ± 2.3	4.116*	0.787*	0.704*	2.3 ± 1.7	3.6
	Female (<i>n</i> = 16)	59.2 ± 3.4	3.6 ± 5.2	2.728*	0.008	0.006	4.8 ± 4.2	9.0
	All (<i>n</i> = 45)	62.3 ± 4.2	2.4 ± 3.7	4.352*	0.671*	0.586*	3.1 ± 3.0	5.6

CE, constant error, arithmetic mean of the difference between estimated and measured VO_{2max} ; *t*, *t* value from paired sample *t*-test; *r*, Pearson correlation coefficient; ICC, intraclass correlation coefficient; MAE, mean absolute error; and MAPE, mean absolute percentage error.

*Statistically significant ($P \leq 0.05$).

Equation 1: VO_{2max} ($mL \cdot kg^{-1} \cdot min^{-1}$) in males = $(3.2131 + 0.0076 \times PWC170) / \text{body mass}$, VO_{2max} ($mL \cdot kg^{-1} \cdot min^{-1}$) in females = $(2.4138 + 0. \times PWC170) / \text{body mass}$.

Equation 2: VO_2 ($mL \cdot kg^{-1} \cdot min^{-1}$) = $(\text{speed} (m \cdot min^{-1}) \times 0.2) + (\text{gradient} \times \text{speed} (m \cdot min^{-1}) \times 0.9) + 3.5$, where the estimated maximal speed was calculated as following steps, (1) linear regression was used based on steady-state heart rate values and running speed were obtained for each stage of the submaximal running test, (2) the linear line was then extrapolated to estimated maximal heart rate ($220 - \text{age}$) to determine the value of estimated maximal speed.

Equation 3: VO_{2max} ($mL \cdot kg^{-1} \cdot min^{-1}$) = $VO_{2\text{-stage 4}} + b (HR_{max} - HR\text{-stage 4})$, where the $VO_{2\text{-stage 4}}$ was calculated based on the steady-state HR in stage 4 of the submaximal running test as the Eq. (2), HR_{max} refers to estimated maximal heart rate ($220 - \text{age}$), and the additional coefficient *b* is calculated from $b = (VO_{2\text{-stage 4}} - VO_{2\text{-stage 3}}) / (HR\text{-stage 4} - HR\text{-stage 3})$, which is the ratio of the difference between the estimated VO_2 of last two stages of submaximal running test and corresponding change of steady-state heart rate.

2007; Akay et al., 2011; Kendall et al., 2012; De Brabandere et al., 2018). Kendall et al. (2012) found an accurate predictive equation [percentage of total error (%TE) = 5.1%, $R^2 = 0.707$, and %SEE = 4.6%] based on critical velocity and anaerobic rowing test trials. Other studies (Vehrs et al., 2007; Akay et al., 2011) also showed accurate equations for predicting VO_{2max} ($r = 0.91$, SEE = $2.54 mL \cdot kg^{-1} \cdot min^{-1}$; $r = 0.94$, SEE = $1.80 mL \cdot kg^{-1} \cdot min^{-1}$) using only the single-stage submaximal treadmill jogging test in healthy adults. However, these equations were developed for the nonathletic population and should be used with caution for well-trained athletes. The findings of this study revealed that the FFT method had a fairly accurate VO_{2max} estimation for well-trained rowers and paddlers and that the FFT method has been proven to accurately estimate VO_{2max} in nonathletic populations (college students, healthy adults, and recreational runners; Kraft and Roberts, 2017; Kraft and Dow, 2019; Passler et al., 2019). However, the rowing exercise program was not included. In the present study, the validity of the FFT method was verified in estimating VO_{2max} in well-trained rowers using a specific submaximal rowing test program.

Analysis of Comparison Between the FFT Method and Previous Predictive Equations

In the rowing program, although the results showed that the previous predictive equation (Eq. 1) by Klusiewicz et al. (2016) had a fairly accurate prediction of VO_{2max} , the FFT method showed even higher accuracy and lower error terms in all groups, as well as in both the male and female subgroups (Table 4). Interestingly, the estimated VO_{2max}

from both the FFT method and Eq. 1 showed a significant overestimation of VO_{2max} (FFT, $1.3 \pm 3.5 mL \cdot kg^{-1} \cdot min^{-1}$; Eq. 1, $3.9 \pm 4.8 mL \cdot kg^{-1} \cdot min^{-1}$). Well-trained athletes have a lower HR at the same intensity than the nonathletic population and have a relatively higher predicted VO_{2max} value based on the linear regression model, which may be the main reason. In addition, the estimation of VO_{2max} from the FFT method in the female subgroup had a nonsignificant difference and a lower MAPE than that in the male subgroup, indicating that the FFT method may provide a more accurate estimation of VO_{2max} in female athletes than that in male athletes when using the rowing program.

In the running program, Marsh (2012) created two equations for estimating VO_{2max} that were suitable for different populations. The present study used these two equations to estimate VO_{2max} in well-trained athletes in the submaximal treadmill running program and found that they all had acceptable accuracy for the overall group. The two equations had similar validity coefficients and agreement levels. However, both equations showed poor accuracy in the female subgroup, which was probably due to the fact that these equations were cross-validated in male athletes instead of in female athletes in the study by Marsh (2012). Similar to the rowing program, the FFT method demonstrated a more accurate prediction of VO_{2max} than these two equations for the overall group. Additionally, the FFT method also had an accurate VO_{2max} estimation in the female subgroup, which was better than that from the two equations. The FFT method modifies equations in the software based on the relevant background variables (e.g., activity class, training zone, and HR variability) and then

improves the accuracy of the VO_{2max} estimation, which could explain this phenomenon.

Analysis of Estimating Bias From the FFT Method

Sartor et al. (2013) have argued that many HR-based submaximal test programs are known to underestimate VO_{2max} because the workload is not high enough to promote adequate parasympathetic withdrawal and concomitant sympathetic activation. The tendency of underestimation was discovered in previous FFT-related studies (Anderson et al., 2019; Passler et al., 2019). Unlike those findings, this study found that the FFT method overestimated VO_{2max} in both the rowing and running programs. A pilot study to detect suitable workloads in submaximal programs may contribute to this phenomenon. Previous studies have indicated that individualized submaximal testing has been utilized in running (Vesterinen et al., 2017), rowing (Otter et al., 2015), and cycling (Lamberts et al., 2011), and high correlations have been found between power or speed at 90% HR_{max} and maximal endurance performance. Other studies also concluded that an optimal submaximal test program includes a proper target intensity, and different workloads for different characteristics may yield a more accurate prediction (Sartor et al., 2013). This study used workloads that may be close to maximal in some individuals, especially in the running program, which may overestimate VO_{2max} .

Practical Application of the FFT Method

Taken together, using the FFT method for VO_{2max} estimation has several practical advantages in the evaluation of aerobic capacity in well-trained athletes. First, only a wearable HR device is needed, and the HR data are recorded during the submaximal testing program available in the software; then, the estimated VO_{2max} value with acceptable accuracy would be automatically calculated. Additionally, the FFT method only requires submaximal tests, and multiple athletes can be tested simultaneously, making its use more feasible during the busy training schedule compared to the direct measurement method for VO_{2max} in the laboratory. Thus, the FFT method can be considered as a potentially convenient and cost-effective alternative to measure the maximal aerobic capacity of well-trained athletes, especially for rowing and running.

Analysis of Limitations

This study had some limitations. The first one is the lack of information regarding the underlying equation of VO_{2max} estimation for the reason that the exact equation can not be obtained from the company. Second, unlike rowers, the lack of sports event-specific testing (paddling ergometer) in paddlers may limit the applicability of the results. However, the running program was performed in this study for the reason that the achievement of VO_{2max} by treadmill running is consistent with paddling ergometer in well-trained paddlers (Augusto Rodrigues dos Santos et al., 2012). Nonetheless, further studies are needed to develop and investigate the predictive models of the FFT method based on submaximal rowing

and running programs, not only in terms of validity but also reliability.

CONCLUSION

The results of the present study indicate that the FFT method provides a considerably accurate estimation of VO_{2max} in well-trained rowers, kayakers, and canoeists, which can be considered as a potentially convenient and cost-effective alternative to measure the maximal aerobic capacity of well-trained athletes, especially for rowing and running.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**, further inquiries can be directed to the corresponding author/s.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by Ethics Committee of Zhejiang Institute of Sports Science, Hangzhou, China. Written informed consent to participate in this study was provided by the participants' legal guardian/next of kin.

AUTHOR CONTRIBUTIONS

WG, HF, and QC performed the material preparation, and data collection and analysis. WG and O-PN wrote the draft of the manuscript. WG, O-PN, and XC conducted the revision. XC supervise the whole program. All authors contributed to the conception and design of this study and approved the final manuscript.

FUNDING

This study was funded by The Wenzhou Medical University Scientific Research Foundation for Talented Scholars (grant number QTJ19033).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fphys.2021.701541/full#supplementary-material>

Supplementary Table 1 | Submaximal FFT programs for both the ROW group and RUN group.

Supplementary Table 2 | GXT programs for both the ROW group and RUN group.

REFERENCES

- Akay, M. F., Zayid, E. I. M., Aktürk, E., and George, J. D. (2011). Artificial neural network-based model for predicting VO₂max from a submaximal exercise test. *Expert Syst. Appl.* 38, 2007–2010. doi: 10.1016/j.eswa.2010.07.135
- Anderson, J. C., Chisenall, T., Tolbert, B., Ruffner, J., Whitehead, P. N., and Conners, R. T. (2019). Validating the commercially available garmin fenix 5x wrist-worn optical sensor for aerobic capacity. *Int. J. Innov. Educ. Res.* 7, 147–158. doi: 10.31686/ijer.vol7.iss1.1293
- Augusto Rodrigues dos Santos, J., Filipe Santos Sousa, R., and Patricia Amorim, T. (2012). Comparison of treadmill and kayak ergometer protocols for evaluating peak oxygen consumption. *Open Sports Sci. J.* 5, 130–133. doi: 10.2174/1875399x01205010130
- Bassett, D. R., and Howley, E. T. (1997). Maximal oxygen uptake: “classical” versus “contemporary” viewpoints. *Med. Sci. Sports Exerc.* 29, 591–603. doi: 10.1097/00005768-199705000-00002
- Beltz, N. M., Gibson, A. L., Janot, J. M., Kravitz, L., Mermier, C. M., and Dalleck, L. C. (2016). Graded exercise testing protocols for the determination of VO₂ max: historical perspectives, progress, and future considerations. *J. Sports Med.* 2016, 1–12. doi: 10.1155/2016/3968393
- Bland, M. J., and Altman, D. G. (1986). Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 327, 307–310. doi: 10.1016/S0140-6736(86)90837-8
- Brink-Elfegoun, T., Kaijser, L., Gustafsson, T., and Ekblom, B. (2007). Maximal oxygen uptake is not limited by a central nervous system governor. *J. Appl. Physiol.* 102, 781–786. doi: 10.1152/jappphysiol.00566.2006
- De Brabandere, A., Op De Beëck, T., Schütte, K. H., Meert, W., Vanwanseele, B., and Davis, J. (2018). Data fusion of body-worn accelerometers and heart rate to predict VO₂max during submaximal running. *PLoS One* 13:e0199509. doi: 10.1371/journal.pone.0199509
- Fletcher, G. F., Ades, P. A., Kligfield, P., Arena, R., Balady, G. J., Bittner, V. A., et al. (2013). Exercise standards for testing and training: a scientific statement from the American Heart Association. *Circulation* 128, 873–934. doi: 10.1161/cir.0b013e31829b5b44
- Howley, E. T., Bassett, D. R. Jr., and Welch, H. G. (1995). Criteria for maximal oxygen uptake: review and commentary. *Med. Sci. Sports Exerc.* 27, 1292–1301.
- Kendall, K. L., Fukuda, D. H., Smith, A. E., Cramer, J. T., and Stout, J. R. (2012). Predicting maximal aerobic capacity (VO₂max) from the critical velocity test in female collegiate rowers. *J. Strength Cond. Res.* 26, 733–738. doi: 10.1519/JSC.0b013e318225f3ac
- Klusiewicz, A., Borkowski, L., Sitkowski, D., Burkhard-Jagodzińska, K., Szczepańska, B., and Ladyga, M. (2016). Indirect methods of assessing maximal oxygen uptake in rowers: practical implications for evaluating physical fitness in a training cycle. *J. Hum. Kinet.* 50, 187–194. doi: 10.1515/hukin-2015-0155
- Klusiewicz, A., and Faff, J. (2003). Indirect methods of estimating maximal oxygen uptake on the rowing ergometer. *Biol. Sport* 20, 181–194.
- Kraft, G. L., and Dow, M. (2019). Validation of the garmin forerunner 920XT VO₂max estimation and the polar RS300X fitness test. *Int. J. Innov. Educ. Res.* 7, 22–28. doi: 10.31686/ijer.vol7.iss9.1658
- Kraft, G. L., and Roberts, R. A. (2017). Validation of the garmin forerunner 920XT fitness watch VO₂peak Test. *Int. J. Innov. Educ. Res.* 5, 61–67.
- Kyröläinen, H., Hackney, A. C., Salminen, R., Repola, J., Häkkinen, K., and Haimi, J. (2018). Effects of combined strength and endurance training on physical performance and biomarkers of healthy young women. *J. Strength Cond. Res.* 32, 1554–1561. doi: 10.1519/jsc.0000000000002034
- Lakomy, H. K. A., and Lakomy, J. (1993). Estimation of maximum oxygen uptake from submaximal exercise on a concept II rowing ergometer. *J. Sports Sci.* 11, 227–232. doi: 10.1080/02640419308729989
- Lamberts, R. P., Swart, J., Noakes, T. D., and Lambert, M. I. (2011). A novel submaximal cycle test to monitor fatigue and predict cycling performance. *Br. J. Sports Med.* 45, 797–804. doi: 10.1136/bjism.2009.061325
- Larsen, G. E., George, J. D., Alexander, J. L., Fellingham, G. W., Aldana, S. G., and Parcell, A. C. (2002). Prediction of maximum oxygen consumption from walking, jogging, or running. *Res. Q. Exerc. Sport* 73, 66–72. doi: 10.1080/02701367.2002.10608993
- Levine, B. D. (2008). $\dot{V}O_2$: What do we know, and what do we still need to know? *J. Physiol.* 586, 25–34. doi: 10.1113/jphysiol.2007.147629
- Marsh, C. E. (2012). Evaluation of the American College of Sports Medicine submaximal treadmill running test for predicting $\dot{V}O_2$ max. *J. Strength Cond. Res.* 26, 548–554. doi: 10.1519/JSC.0b013e318220d9a8
- Matabuena, M., Vidal, J. C., Hayes, P. R., and Huelin Trillo, F. (2018). A 6-minute sub-maximal run test to predict VO₂ max. *J. Sports Sci.* 36, 2531–2536. doi: 10.1080/02640414.2018.1468149
- Midgley, A. W., Carroll, S., Marchant, D., McNaughton, L. R., and Siegler, J. (2009). Evaluation of true maximal oxygen uptake based on a novel set of standardized criteria. *Appl. Physiol. Nutr. Metab.* 34, 115–123. doi: 10.1139/H08-146
- Midgley, A. W., McNaughton, L. R., and Carroll, S. (2007). Effect of the VO₂ time-averaging interval on the reproducibility of VO₂max in healthy athletic subjects. *Clin. Physiol. Funct. Imag.* 27, 122–125. doi: 10.1111/j.1475-097x.2007.00725.x
- Montgomery, P. G., Green, D. J., Etxebarria, N., Pyne, D. B., Saunders, P. U., and Minahan, C. L. (2009). Validation of heart rate monitor-based predictions of oxygen uptake and energy expenditure. *J. Strength Cond. Res.* 23, 1489–1495. doi: 10.1519/JSC.0b013e3181a39277
- Nelson, M. B., Kaminsky, L. A., Dickin, D. C., and Montoye, A. H. K. (2016). Validity of consumer-based physical activity monitors for specific activity types. *Med. Sci. Sports Exerc.* 48, 1619–1628. doi: 10.1249/MSS.0000000000000933
- Otter, R. T. A., Brink, M. S., Lamberts, R. P., and Lemmink, K. A. P. M. (2015). A new submaximal rowing test to predict 2,000-m rowing ergometer performance. *J. Strength Cond. Res.* 29, 2426–2433. doi: 10.1519/JSC.0000000000000902
- Passler, S., Bohrer, J., Blöching, L., and Senner, V. (2019). Validity of wrist-worn activity trackers for estimating VO₂max and energy expenditure. *Int. J. Environ. Res. Public Health* 16:3037. doi: 10.3390/ijerph16173037
- Riebe, D., Ehrman, J., Liguori, G., and Magal, M. (2018). *ACSM's Guidelines for Exercise Testing and Prescription*. Philadelphia: Wolters Kluwer.
- Saltin, B., and Strange, S. (1992). Maximal oxygen uptake: “old” and “new” arguments for a cardiovascular limitation. *Med. Sci. Sports Exerc.* 24, 30–37. doi: 10.1249/00005768-199201000-00007
- Sartor, F., Vernillo, G., De Morree, H. M., Bonomi, A. G., La Torre, A., Kubis, H. P., et al. (2013). Estimation of maximal oxygen uptake via submaximal exercise testing in sports, clinical, and home settings. *Sport. Med.* 43, 865–873. doi: 10.1007/s40279-013-0068-3
- Shephard, R. J. (2009). Is the measurement of maximal oxygen intake passe? *Br. J. Sports Med.* 43, 83–85. doi: 10.1136/bjism.2008.052506
- Tanner, R. K., and Gore, C. J. (2013). *Physiological tests for Elite Athletes*, 2nd Edn. Champaign, IL: Human Kinetics.
- Tsiaras, V., Zafeiridis, A., Dipla, K., Patras, K., Georgoulis, A., and Kellis, S. (2010). Prediction of peak oxygen uptake from a maximal treadmill test in 12- to 18-year-old active male adolescents. *Pediatr. Exerc. Sci.* 22, 624–637. doi: 10.1123/pes.22.4.624
- Vehrs, P. R., George, J. D., Fellingham, G. W., Plowman, S. A., and Dustman-Allen, K. (2007). Submaximal treadmill exercise test to predict VO₂ max in fit adults. *Meas. Phys. Educ. Exerc. Sci.* 11, 61–72. doi: 10.1080/10913670701294047
- Vesterinen, V., Nummela, A., Laine, T., Hynynen, E., Mikkola, J., and Häkkinen, K. (2017). A submaximal running test with postexercise cardiac autonomic and neuromuscular function in monitoring endurance training adaptation. *J. Strength Cond. Res.* 31, 233–243. doi: 10.1519/JSC.00000000000001458

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Gao, Nuutila, Fang, Chen and Chen. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.