



# Melatonin Improves Waterlogging Tolerance of *Malus baccata* (Linn.) Borkh. Seedlings by Maintaining Aerobic Respiration, Photosynthesis and ROS Migration

Xiaodong Zheng<sup>1</sup>, Jingzhe Zhou<sup>2</sup>, Dun-Xian Tan<sup>3</sup>, Na Wang<sup>1</sup>, Lin Wang<sup>1</sup>, Dongqian Shan<sup>1</sup> and Jin Kong<sup>1\*</sup>

<sup>1</sup> College of Horticulture, China Agricultural University, Beijing, China, <sup>2</sup> Beijing Soil and Fertilizer Work Station, Beijing, China, <sup>3</sup> Department of Cellular and Structural Biology, UT Health Science Center San Antonio, San Antonio, TX, USA

## OPEN ACCESS

### Edited by:

Wei Hu,  
Institute of Tropical Bioscience and  
Biotechnology, Chinese Academy of  
Tropical Agricultural Sciences, China

### Reviewed by:

Shi Xiao,  
Sun Yat-sen University, China  
Wei Fan,  
Yunnan Agricultural University, China

### \*Correspondence:

Jin Kong  
jinkong@cau.edu.cn

### Specialty section:

This article was submitted to  
Plant Cell Biology,  
a section of the journal  
Frontiers in Plant Science

**Received:** 26 December 2016

**Accepted:** 20 March 2017

**Published:** 05 April 2017

### Citation:

Zheng X, Zhou J, Tan D-X, Wang N,  
Wang L, Shan D and Kong J (2017)  
Melatonin Improves Waterlogging  
Tolerance of *Malus baccata* (Linn.)  
Borkh. Seedlings by Maintaining  
Aerobic Respiration, Photosynthesis  
and ROS Migration.  
*Front. Plant Sci.* 8:483.  
doi: 10.3389/fpls.2017.00483

Waterlogging, one of the notorious abiotic stressors, retards the growth of apple plants and reduces their production. Thus, it is an urgent agenda for scientists to identify the suitable remedies for this problem. In the current study, we found that melatonin significantly improved the tolerance of apple seedlings against waterlogging stress. This was indicated by the reduced chlorosis and wilting of the seedlings after melatonin applications either by leaf spray or root irrigation. The mechanisms involve in that melatonin functions to maintain aerobic respiration, preserves photosynthesis and reduces oxidative damage of the plants which are under waterlogging stress. Melatonin application also enhances the gene expression of its synthetic enzymes (MbT5H1, MbAANAT3, MbASMT9) and increases melatonin production. This is the first report of a positive feedback that exogenous melatonin application promotes the melatonin synthesis in plants. A post-transcriptional regulation apparently participated in this regulation. When exogenous melatonin meets the requirement of the plants it is found that the protein synthesis of MbASMT9 was suppressed. Taken together, the results showed that melatonin was an effective molecule to protect plant, particularly apple plant, against waterlogging stress.

**Keywords:** melatonin, waterlogging, ROS, oxidative stress, aerobic respiration, photosynthesis

## INTRODUCTION

Waterlogging, is a major agricultural constraint that limits crop growth and reduces their yield (Xu et al., 2013). It is frequently encountered during the raining seasons in many areas worldwide. The excessive waterlogging causes root damage, impairs the water uptake and, finally leads to chlorosis and wilting of the plants (Arbona et al., 2008). It was estimated that waterlogging stress resulted in nearly 40–80% of the crop yield loss in the area greater than 17 million km<sup>2</sup> (Voeselek and Sasidharan, 2013; Shabala et al., 2014).

The reactive oxygen species (ROS) is believed to play a critical role in the response of plant to waterlogging stress. At the early stage during waterlogging, the elevated ROS molecules functions as an important second messenger in signaling for response. Following the prolonged waterlogging, the increased anaerobic respiration of root and the responsive stomata closure in leaves induce a burst of excessive ROS production. If the excessive ROS is not migrated properly, it will cause plant oxidative damage and finally, it leads to roots rotting and leaves wilting (Hossain et al., 2009).

Plants have already developed a series of antioxidant mechanisms to defend themselves against oxidative stress. These include small molecule antioxidants (SMA) and antioxidant enzymes (Apel and Hirt, 2004; Mittler et al., 2004). SMA includes ascorbic acid, carotenoids, tocopherol, glutathione, polyphenol, etc. They can scavenge ROS with different chemical reactions (Smirnoff, 2000; Kuzniak and Sklodowska, 2001; Foyer and Noctor, 2005). The antioxidant enzymes are mainly those of superoxide dismutase (SOD), catalase (CAT), peroxidase (POD), etc. Both SMA and antioxidant enzymes work coordinately to keep the oxidative stress in check (Hossain et al., 2009). In addition to the SMA mentioned above, melatonin (*N*-acetyl-5-methoxytryptamine) is a potent free radical scavenger and an antioxidant (Tan et al., 1993, 2012). Melatonin was identified in plants in Dubbels et al. (1995) and Hattori et al. (1995). Since then it has been reported to exist in many plants and plant products (Ramakrishna et al., 2012; Yip et al., 2013; Shi et al., 2015b; Zhao et al., 2015a; Li C. et al., 2016; Ma et al., 2016; Xu et al., 2016). Different from other antioxidants, it is an amphiphilic molecule which makes it distribute in all cellular compartments including cytosol, membrane, mitochondria and chloroplasts (López et al., 2009; Byeon et al., 2013; Back et al., 2016). Also, melatonin as well as its metabolites can eliminate different kinds of ROS including superoxide anion ( $O_2^{\bullet-}$ ), hydrogen peroxide ( $H_2O_2$ ), hydroxyl radical ( $\bullet OH$ ), singlet oxygen ( $^1O_2$ ), peroxynitrite anion ( $ONOO^-$ ) and nitric oxide (NO) (Tan et al., 2013). Melatonin has been reported to protect plants against a variety of abiotic and biotic stresses (Zuo et al., 2014; Liang et al., 2015; Liu et al., 2015; Shi et al., 2015a; Wang et al., 2015; Zhao et al., 2015b; Li C. et al., 2016; Xu et al., 2016). The transgenic *Arabidopsis* plants, ecotopically expressing melatonin synthetic gene *MzASMT1*, had higher endogenous melatonin production and significantly lower ROS than that of their wild types. The melatonin enriched transgenic *Arabidopsis* had a greater tolerance to drought stress than the wild types (Zuo et al., 2014). The exogenous melatonin also improved tolerance of tomato plants against alkaline stress by migrating  $O_2^{\bullet-}$  and  $H_2O_2$  (Liu et al., 2015). Melatonin application to apple leaves alleviated the drought-induced inhibition of photosynthesis (Wang et al., 2012). However, the effects of the exogenous melatonin application on apple plants which are under the sustained waterlogging condition have not been reported yet.

Reduced apple yield caused by waterlogging stress is a worldwide problem to be solved. Apple is a perennial woody plant, thus, the damage from waterlogging stress not only reduces apple yield of the current year, but also suppresses the tree vigor which also leads to the yield loss in the following years (Qu et al., 1999). *Malus baccata* (Linn.) Borkh. is used as a rootstock, which frequently suffers from the waterlogging stress (Wang et al., 2013). Therefore improvement of its waterlogging resistance by melatonin application will provide a potential cultivation method for apple production.

In current study, we investigated the protective effects of melatonin on *M. baccata* seedlings, which were subjected to the waterlogging stress. Furthermore, the potential mechanisms of these protections were also explored and discussed.

## MATERIALS AND METHODS

### The Cultivation of Plant Material

Seeds of apple (*M. baccata*) were sown in sippy vermiculite. Two weeks later, the seedlings were watered with half-strength Hoagland's nutrient solution (Li M.Q. et al., 2016). When the seedlings developed to have four leaves, they were watered with complete nutrient solution. The plants were kept in green house with the temperature at a constant  $22 \pm 2^\circ C$  and a 16/8 h light/dark cycle. The light intensity was approximately  $100 \mu mol \cdot m^{-2} \cdot s^{-1}$ .

### Waterlogging Stress/Melatonin Treatment and Sample Collection

After the *M. baccata* seedlings developed to have four leaves, a total of 96 seedlings were transplanted into the glass container with sterilized matrix soil. They were divided into eight groups. Plants in group I were watered with the 200 mL normal nutrient solution. Totally 25 mL normal nutrient solution was added every 3 days as control. The waterlogging stress was conducted in the remaining seven groups by keeping the soil being covered with 300 mL the nutrient solution and added 25 mL normal nutrient solution every 3 days.

The seedlings from group II was waterlogging stressed without supply of exogenous melatonin. The waterlogging stressed seedlings from group III to VIII were treated with different concentration of melatonin by spraying or irrigation. Melatonin was dissolved in 100% ethanol at a concentration of 10 mM and stored at  $-20^\circ C$  as a stock solution. When use, melatonin was then diluted into 50, 100, and 200  $\mu M$ , respectively with deionized water. These different concentrations of melatonin were sprayed to the leaves of seedlings every other day in group III, IV, and V, respectively. Two pieces of hardboard were used to avoid sprayed melatonin dropping into the soil. In group VI, VII, and VIII, melatonin was directly supplemented to the nutrient solution at the concentrations of 200, 400, and 600  $\mu M$ , respectively. Groups I and II was also applied with equal volume of ethanol. The experiments were independently repeated three times. Photos were taken before and after 9 days of waterlogging stress/melatonin treatments.

The leaves and roots were collected from the all groups of seedlings, respectively, before and after 9 days of waterlogging stress/melatonin treatment.

### Melatonin Measurement

The leaves collected from the seedlings were immediately frozen at  $-80^\circ C$  for future melatonin detection. Around 1 g of frozen leaves of *M. baccata* was ground to a fine powder in liquid nitrogen. The powder was mixed with 10 mL methanol and ultrasonicated (80 Hz) for 35 min at  $45^\circ C$ . The sample preparation and HPLC detection of melatonin were performed as described by Zhao et al. (2013). Each experiment was independently repeated three times.

### RNA Extraction and RT-PCR Analysis

Total RNA was isolated from the leaves of seedlings of the eight groups, respectively, before and after 9 days of waterlogging

stress/melatonin treatment, using the EASY spin Plant RNA Rapid Extraction Kit (Biomed, Beijing, China). The first-strand cDNA was synthesized following the protocol of Kit (Promega, Madison, WI, USA). The cDNAs were used as template for RT-PCR. The specific primers were designed according to the sequence of melatonin synthesized enzyme genes *MbASMT9* (KJ156531), *MbAANAT3* (KJ156532), and *MbT5H1*, respectively, by Primer 5 software and checked by BLAST search in the apple genome<sup>1</sup> (*MbASMT9* Forward Primer 5'-TGATCTGCCCCATGTCGT-3', Reverse Primer 5'-CTTTGTGGCGAGGGAAAC-3'; *MbAANAT3* Forward Primer 5'-CGCTCCCTAACTACCAACCA-3', Reverse Primer 5'-ACAAATCCCTTCCCTACCAG-3'; *MbT5H1* Forward Primer 5'-ATCCGTAAGATTTGTATACTTGAGCT-3', Reverse Primer 5'-TCACCGACCAAGATAATAGCCT-3').

RT-PCRs were performed use of 20  $\mu$ L reaction mixtures containing 20 ng of first-strand cDNA, 2  $\times$  PCR Mix 5  $\mu$ L (CW BIO, Beijing, China), 0.5  $\mu$ M of each of the forward and reverse primers and appropriate amounts of ddH<sub>2</sub>O. The *Actin* gene was used as the internal standard, and the PCR program for *Actin* was as following: 94°C for 5 min; 28 cycles at 94°C for 30 s, 55°C for 30 s, and 72°C for 30 s; and 72°C for 10 min, amplified with primers (Forward Primer 5'-CAATGCCTGCCATGTATG-3', Reverse Primer 5'-CCAGCAGCTTCCATTCCAAT-3'). The PCR products were analyzed in 1% TAE-agarose gel stained by goodview. The quantification of amplified *Actin*, *MbT5H1*, *MbAANAT3*, and *MbASMT9* fragment was done by the ImageJ software<sup>2</sup>, the ratios of *MbT5H1*, *MbAANAT3*, *MbASMT9* and *Actin* were calculated.

## The Cloning of *MbASMT9* Gene and the Expression of *MbASMT9* Protein in *Escherichia coli*

The coding frame of *MbASMT9* was amplified and ligated into pMD 19-T Simple, which was then digested with BamHI/EcoRI and inserted into the pGEX-6p-1. The protein expression of *MbASMT9* was analyzed according to Li et al. (2015). The purity of the GST-MbASMT9 protein was confirmed by SDS-PAGE.

## Protein Extraction and Western Blot Analysis

The total protein was isolated from leaves of seedlings according to Wang et al. (2006). Western blot was applied with antibody of *MbASMT9* (rabbit, 1:3000). The recombinant GST-MbASMT9 was used as an antigen to raise polyclonal GST-MbASMT9 antibodies in rabbit. The preparation of *MbASMT9* antibody was carried out in accordance with relevant guidelines and regulations. The experimental protocols of the *MbASMT9* antibody preparation were reviewed and approved by Beijing Municipal Science and Technology Commission. The chemiluminescent signals were detected using an ECL detection kit (Amersham-Pharmacia, USA). The loading control of the Western blot was stained by Coomassie Brilliant Blue.

<sup>1</sup><https://www.rosaceae.org/species/malus/all>

<sup>2</sup><http://en.wikipedia.org/wiki/ImageJ>

## Detection of Antioxidant Enzyme Activities

A total of 0.3 g leaves or 0.3 g roots of *M. baccata* seedlings were ground with 8 mL chilled 50 mM phosphate buffer (pH 7.8), then they were transferred into 10 mL tubes and centrifuged at 4°C for 15 min at 10,000 g. The supernatants were diluted with phosphate buffer to 10 mL and used for enzyme activity detection. The enzyme activity of SOD was detected according to the method of Stewart and Bewley (1980). CAT activity was measured as the absorbance at 240 nm wave length according to the method of Ma et al. (2016). POD (Peroxidase) activity was measured by the changes in absorbance at 470 nm due to guaiacol oxidation according to the method of Shi et al. (2015b). Each experiment was independently repeated at least three times.

## Measurements of Enzyme Activities of Alcohol Dehydrogenase (ADH) and Succinate Dehydrogenase (SDH)

A total of 0.3 g roots for each group were collected to detect the enzyme activity of ADH and SDH before and after 9 days of waterlogging stress/melatonin treatment to identify the alterations of anaerobic and aerobic respiration. ADH activity was measured as reported by Yamashita et al. (2015). SDH activity was detected according to Landi et al. (2009). Each experiment was independently repeated at least three times.

## Detections of ROS Level and Malondialdehyde (MDA) Content

The ROS level in roots was detected according to the method modified from Zuo et al. (2014). The intact roots were collected from each group and the fresh roots were immediately used for ROS detection before and after 9 days of waterlogging stress/melatonin treatment. Simply, the entire roots were incubated with the 5-(and 6)-chloromethyl-2'-7'-dichlorodihydrofluorescein diacetate acetyl ester (CM-H<sub>2</sub>DCFDA) solution for 20 min and washed with distilled H<sub>2</sub>O to remove excess CM-H<sub>2</sub>DCFDA. The fluorescence images were obtained with a Leica stereoscope (Leica, Wetzlar, Germany) ( $\times 10$ ).

In addition to roots, the ROS level in leaves was also detected. The leaves were collected from the each group, respectively before and after 9 days of waterlogging stress/melatonin treatments and used for detection of O<sub>2</sub><sup>•-</sup> and H<sub>2</sub>O<sub>2</sub> immediately. For O<sub>2</sub><sup>•-</sup> measurement, the leaf histochemical staining was vacuum infiltrated with 0.1 mgmL<sup>-1</sup> nitroblue tetrazolium in 25 mM K-HEPES buffer (pH 7.9) for 40 min. Then the samples were kept at 25°C in dark for an additional 4 h. For the H<sub>2</sub>O<sub>2</sub> detection the leaves were vacuum infiltrated with 0.1 mg·mL<sup>-1</sup> DAB in 50 mM Tris-acetate (pH 3.8) and were incubated at 25°C in dark for 24 h. Then leaves for either O<sub>2</sub><sup>•-</sup> or H<sub>2</sub>O<sub>2</sub> detection were washed in 80% ethanol every 10 min at 80°C until the leaves lost green color completely (Gong et al., 2014). The MDA detection in the leaf samples were performed according to Zhao et al. (2013). Each experiment was independently repeated at least three times.

## Analyses of Chlorophyll (Chl) Content and Photosynthetic Rate (Pn) in Leaves of Seedlings

The extraction of Chl was conducted according to Porra et al. (1989). A total of 0.2 g leaves were homogenized in 2–3 mL 80% acetone. After filtration with four layers of gauze (1 mm × 1 mm), the homogenate was diluted with 80% acetone to 25 mL. The absorbance of the extract was measured at 645 and 663 nm. Chl concentration was calculated from the following equations: Chl a =  $12.72 \times OD_{663} - 2.59 \times OD_{645}$ ; Chl b =  $22.88 \times OD_{645} - 4.67 \times OD_{663}$ ; chl = chl a + chl b (Arnao and Hernández-Ruiz, 2009).

The photosynthetic rate was measured by LI-6400XT (LI-COR, Lincoln, NE, USA) according to the producer's protocol. The light intensity was at  $800 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ . The humidity was about 50% and the temperature was 23°C. Each experiment was independently repeated three times.


## Statistical Analysis

The data are expressed as means  $\pm$  SD. One-way ANOVA was used for the normality evaluation followed by a Tukey–Kramer multiple comparison test. The statistical significant difference was set up when  $P < 0.05$ . Statistical evaluations were carried out using SPSS software (IBM, Armonk, NY, USA).

## RESULTS

### Melatonin Applications Improved the Tolerance of *M. baccata* Seedlings against Waterlogging Stress

The results showed that waterlogging stress significantly impaired the *M. baccata* seedling growth. Those plants which were under waterlogging stress for 9 days wilted severely compared to the normal controls (Figure 1A). When melatonin was sprayed to



**FIGURE 1 |** The effects of melatonin on the *M. baccata* seedlings which were suffered from waterlogging stress. **(A)** Before and after waterlogging stress. **(B):** (a) The phenotype of *M. baccata* seedlings with melatonin leaf spray at different concentrations before and after waterlogging stress. (b) The phenotype of *M. baccata* seedlings with melatonin irrigation at different concentrations before and after waterlogging stress.

the seedlings at the concentrations of 50, 100, and 200  $\mu\text{M}$ , respectively, it was apparent that melatonin spraying improved tolerance of seedlings against waterlogging stress. The protective effects of melatonin indicated a dose-responsive manner. The seedlings treated with 200  $\mu\text{M}$  of melatonin, even under the waterlogging stress, showed a similar phenotype as the normal controls. The similar results were also observed in the seedlings those melatonin was irrigated (Figure 1B). When melatonin was irrigated to the seedlings at the concentrations of 200, 400, and 600  $\mu\text{M}$ , respectively, it also increased waterlogging resistance of seedlings as a dose-dependent manner. The irrigation with 600  $\mu\text{M}$  of melatonin, gained a similar phenotype as spraying with 200  $\mu\text{M}$  melatonin. The percentage of leaf chlorosis was also analyzed. It was clearly to see that the percentage of leaf chlorosis was significantly increased after waterlogging treatment. After melatonin was applied to the seedlings under waterlogging, the percentage of leaf chlorosis were significantly decreased (Supplementary Figure 1).


### The Effects of Waterlogging Stress and Melatonin Application on *De novo* Melatonin Synthesis in Seedlings

To find the potential effects of waterlogging stress and melatonin application on the endogenous melatonin synthesis, the melatonin level and its synthetic gene expression were detected. The results showed that waterlogging stress significantly elevated plants melatonin levels in leaves (87 vs. 21 ng/g FW of control) and in roots (73 vs. 16 ng/g FW of control) (Figure 2A).

When exogenous melatonin was applied to the stressed seedlings, their endogenous melatonin was further increased compared to the plants under waterlogging stress alone (Figure 2A). For example, the endogenous leaf melatonin level in melatonin (200  $\mu\text{M}$ ) sprayed plants was 256 ng/g FW, which was 2.9 times higher than that in leaves of waterlogged seedling alone (87 ng/g FW). The similar results were observed in roots and also in melatonin irrigated seedlings. In accordance with the elevated melatonin production, the gene expressions of melatonin synthetic enzymes including ASMT (acetylserotonin *O*-methyltransferase), AANAT (aralkylamine *N*-acetyltransferase) and T5H (tryptamine 5-hydroxylase), were slightly upregulated under the waterlogging stress and melatonin treatment compared to the controls (Figure 2B). The upregulated gene expression of *ASMT9*, which is the supposed melatonin synthetic rate-limiting enzyme, failed to result in the increase in its protein level. In contrast, its protein level was declined when compared to the controls (Figure 2C). Obviously, the post-transcriptional regulation occurred for *ASMT9*.

### The Effects of Melatonin on Antioxidant Enzymes

The activities of the antioxidant enzymes including SOD, POD, CAT were measured in four groups of *M. baccata* seedlings. These groups included the normal control group, waterlogging stress alone, waterlogging stress treated with melatonin (200  $\mu\text{M}$  spray and waterlogging stress treated with





**FIGURE 2 | Effects of waterlogging and exogenous melatonin application on the melatonin synthesis in *M. baccata* seedlings 9 days after waterlogging stress. (A)** Endogenous melatonin levels in leaves and in roots of the *M. baccata* seedlings. **(B)** The relative expression of melatonin synthetic genes (*MbASMT9*, *MbAANAT3* and *MbT5H1*) in the leaves. **(C)** The protein level of *MbASMT9* detected by Western blot analysis in the leaves. The data are means  $\pm$  SD of triplicate experiments. Asterisks (\*) and different letters indicate significant differences from the control ( $P < 0.05$ ).

melatonin (600  $\mu\text{M}$ ) irrigation. Waterlogging stress significantly suppressed all antioxidant enzymes tested in both levels and roots of the seedlings. However, melatonin treatment could recover the activities of these antioxidant enzymes suppressed by waterlogging stress in a great degree. For example, melatonin (200  $\mu\text{M}$ ) spray recovered the activity of CAT in leaves to the level comparable to the control seedlings (Figure 3).

### Effects of Melatonin on Aerobic and Anaerobic Respiration of *M. baccata* Seedlings under Waterlogging Stress

The activities of ADH and SDH indicate the anaerobic and aerobic respirations, respectively. It was found that the ADH activity (11.71 U/mg), the index of anaerobic respiration, was significantly higher in the roots of waterlogging stressed seedlings



than that of normal control (7.01 U/mg). Both melatonin spray and irrigation reduced the anaerobic respiration indicated by decreased ADH activity (Figure 4A). In contrast, waterlogging stress suppressed the aerobic respiration indicated by the significantly reduced SDH activity compared to the control seedlings. This tendency was reversed by melatonin treatment in a great degree (Figure 4B).

### The Effect of Melatonin on $O_2^{\bullet-}$ and $H_2O_2$ Production and the Oxidative Damage Induced by Waterlogging Stress


Waterlogging stress significantly elevated the  $O_2^{\bullet-}$  and  $H_2O_2$  production both in leaves and in roots of the seedlings. These were indicated by the increased fluorescent staining intensities on them. Melatonin treatment significantly reduced  $O_2^{\bullet-}$  and  $H_2O_2$  productions, no matter melatonin was sprayed to the leaves or irrigated to the roots of the seedlings (Figures 5A,C). Accordingly, waterlogging stress caused oxidative damage in both leaves and roots of the seedlings. This was indicated by the increased content of MDA. It was expected that melatonin treatment significantly reduced the MDA levels in leaves and also in roots (Figures 5B,D).

### The Effect of Melatonin on the Chlorophyll Content and Photosynthetic Rate


Waterlogging stress dramatically reduced the Chl content in the leaves of seedlings. After 9 days of waterlogging stress, the Chl content of these plants (2.4 mg/g FW) was only roughly half of the value of control seedlings (4.1 mg/g FW). The photosynthetic rate of the waterlogging stressed plants was 1.67  $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ . The value was significantly lower than that of control seedlings (2.67  $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ ). Melatonin treatments (spray or irrigation) improved both Chl content and photosynthetic rate which were suppressed by the waterlogging stress (Figures 6A,B).

## DISCUSSION

Waterlogging, caused by poor drainage, flooding, and long periods of rainfall, usually occurs in summer and autumn, which hampers apple tree growth and results in yield loss (Yu et al., 2015). Because of the global warming effect, it is predicted that this climate change will bring more rainfall yearly and thus




**FIGURE 5 | Effects of melatonin on ROS levels and oxidative damage in *M. baccata* seedlings which were under waterlogging stress. (A)** ROS level in roots. **(B)** MDA production in roots. **(C)** The levels of superoxide anion and hydrogen peroxide in leaves. **(D)** MDA production in leaves. The waterlogging stress lasted for 9 days. Scale bars in **(A)** represent 0.5 cm. Scale bars in **(C)** represent 1cm. The data are means ± SD of triplicate experiments. Different letters indicate significant differences from the control ( $P < 0.05$ ).



**FIGURE 6 | Effects of melatonin on the chlorophyll content and photosynthetic rate of the *M. baccata* seedlings before and after waterlogging stress. (A)** Chlorophyll content. **(B)** Photosynthetic rate. The data are means ± SD of triplicate experiments. Different letters indicate significant differences from the control ( $P < 0.05$ ).

the waterlogging will be frequently encountered by the crops worldwide. This will bring severe problem for these crops which are intolerant to the waterlogging stress including the apple trees. To solve this problem, it has become an urgent agenda for scientist.

In the current study, a unique molecule, melatonin, was tested with this purpose. Melatonin was reported to enhance plant tolerance against various abiotic stressors including cold, hot, drought, salinity, and chemical pollutants (Bajwa et al., 2014; Wei et al., 2014; Liang et al., 2015; Liu et al., 2015; Shi et al., 2015b;



Li X. et al., 2016; Xu et al., 2016). But the effect of melatonin on waterlogging resistance is still to be investigated.

Long lasting waterlogging caused apple seedling damage. Under waterlogging, the seedlings changed their metabolism from aerobic to anaerobic respiration (Fukao and Bailey-Serres, 2004). During this transition, a burst of ROS was generated and this resulted in oxidative damage of the seedlings (Hossain et al., 2009; Xu et al., 2013; Takahashi et al., 2015). This was indicated by the phenotype alterations of the seedlings such as chlorosis and wilting (Figure 1). However, all these changes were partially or completely prevented by melatonin application either by spraying to the leaves or irrigating to the roots. The primary mechanism of melatonin to improve the tolerance of apple seedlings against waterlogging stress may relate its potent free radical scavenging and antioxidant capacity (Tan et al., 1993, 2012). Melatonin not only directly scavenges the ROS but also upregulates the activities of a variety of antioxidant enzymes (Wang et al., 2012). This was in accordance with our observations in the current study (Figures 3, 5).

In the current study, both waterlogging stress and melatonin application significantly increased melatonin level of the plants. It is well known that various stressors induce melatonin production in plants as well as in animals (Byeon et al., 2012; Li C. et al., 2016; Xu et al., 2016). This is considered as self-defense of organisms against external insults (Tan et al., 2015). Our observation provided new evidence to support this consideration. Waterlogging induced the gene expression of the melatonin synthetases and increased melatonin

production in the apple seedlings. This positive response mechanism at the RNA level, for melatonin production in apple plants, probably also, exists in other plants. When exogenous melatonin was applied under waterlogging stress, the stress-induced expression of the melatonin synthetases still maintained their high expression. But obviously, the slight expression change between melatonin + waterlogging treatment and only waterlogging treatment can't explain the significant melatonin increase after exogenous melatonin applied under waterlogging. It seemed that melatonin absorbed from outside mainly resulted in melatonin increase. In addition, for the MbASMT9, a melatonin synthetic rate-limiting enzyme, which was located in chloroplasts (Kang et al., 2013; Zheng et al., 2017), its upregulated RNA expression did not result in elevated protein level. In contrast, its protein level reduced after melatonin application. This probably was the first report to document a post-transcriptional regulation of melatonin synthesis in plants. There are accumulated post-transcriptional regulation reported, including RNA processing, transport and degradation, translation control to fine-tune biological processes for plants in response to the environmental changes (Palatnik et al., 2003, 2007; Cowley et al., 2012; Saze et al., 2013). Obviously, herein, the post-transcriptional regulation conserves resources and energy of the plants to avoid extra melatonin production when exogenous applied melatonin meets their requirement. This observation provided valuable information as to use of exogenous melatonin to improve plant tolerance to against stressors.

There are accumulated reports described the waterlogging caused transition from aerobic to anaerobic respiration in root, which is also confirmed by us (Figure 4). Our results also uncovered the role of melatonin to maintain aerobic respiration under waterlogging stress, by efficient suppression of the ROS burst and subsequent mitochondria degradation. The chlorosis is a typical sign unavoidably happening after severe waterlogging. Due to the reason that waterlogging leads to water shortage and subsequent quick stomata closure, the high concentration of  $O_2$  can't be released out and photosynthetic electron transportation is blocked in chloroplasts. Therefore, the ion leakage from the electron transport chain would induce in over-produced  $O_2^{\bullet-}$  and  $H_2O_2$ , which destroy chlorophyll and lead to the disintegration of chloroplasts (Musgrave, 1994; Smethurst and Shabala, 2003). The protective effects of melatonin on Chl decay, photosynthetic capacity and stomata configurations have been reported previously in other stressors such as in drought and hot (Wang et al., 2012; Xu et al., 2016). Actually, chloroplasts, as one of the most suffered organelles from ROS, were proved to be the major site for melatonin production. A large amount of melatonin is needed to maintain its structure and function. Therefore absorbed and *in vivo* synthesized melatonin can function together to migrate waterlogging induced ROS and help to survive the stress. Here we reported that melatonin application preserved the Chl content and maintained the photosynthetic rate in seedlings suffered from waterlogging stress. The high content of Chl and efficient photosynthesis are required for high yield of apple production (Li et al., 2000). It is our

speculation that melatonin application in the field will increase the tolerance of apple tree and reduces apple yield loss against waterlogging stress. The speculated mechanisms are summarized in **Figure 7**.

## AUTHOR CONTRIBUTIONS

Designed the studies: JK, Undertook the experimental work: XZ, JZ, NW, LW, and DS. Contributed to figures and manuscript preparation: XZ, D-XT, and JK. All authors read and approved the final manuscript.

## REFERENCES

- Apel, K., and Hirt, H. (2004). Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* 55, 373–399. doi: 10.1146/annurev.arplant.55.031903.141701
- Arbona, V., Hossain, Z., López-Clement, M. F., Pérez-Clemente, R. M., and Gómez-Cadenas, A. (2008). Antioxidant enzymatic activity is linked to waterlogging stress tolerance in citrus. *Physiol. Plant.* 132, 452–466. doi: 10.1111/j.1399-3054.2007.01029.x
- Arnao, M. B., and Hernández-Ruiz, J. (2009). Protective effect of melatonin against chlorophyll degradation during the senescence of barley leaves. *J. Pineal Res.* 46, 58–63. doi: 10.1111/j.1600-079X.2008.00625.x
- Back, K., Tan, D. X., and Reiter, R. J. (2016). Melatonin biosynthesis in plants: multiple pathways catalyze tryptophan to melatonin in the cytoplasm or chloroplasts. *J. Pineal Res.* 61, 426–437. doi: 10.1111/jpi.12364
- Bajwa, V. S., Shukla, M. R., Sherif, S. M., Murch, S. J., and Saxena, P. K. (2014). Role of melatonin in alleviating cold stress in *Arabidopsis thaliana*. *J. Pineal Res.* 56, 238–245. doi: 10.1111/jpi.12115
- Byeon, Y., Lee, H. Y., Lee, K., Park, S., and Back, K. (2013). Cellular localization and kinetics of the rice melatonin biosynthetic enzymes SNAT and ASMT. *J. Pineal Res.* 56, 107–114. doi: 10.1111/jpi.12103
- Byeon, Y., Park, S., Kim, Y. S., Park, D. H., Lee, S., and Back, K. (2012). Light-regulated melatonin biosynthesis in rice during the senescence process in detached leaves. *J. Pineal Res.* 53, 107–111. doi: 10.1111/j.1600-079X.2012.00976.x
- Cowley, M., Wood, A. J., Böhm, S., Schulz, R., and Oakey, R. J. (2012). Epigenetic control of alternative mRNA processing at the imprinted Hec3/Nap15 locus. *Nucleic Acids Res.* 40, 8917–8926. doi: 10.1093/nar/gks654
- Dubbels, R., Reiter, R. J., Klenke, E., Goebel, A., Schnakenberg, E., Ehlers, C., et al. (1995). Melatonin in edible plants identified by radioimmunoassay and by high performance liquid chromatography-mass spectrometry. *J. Pineal Res.* 18, 28–31. doi: 10.1111/j.1600-079X.1995.tb00136.x
- Foyer, C. H., and Noctor, G. (2005). Redox homeostasis and antioxidant signaling: a metabolic interface between stress perception and physiological responses. *Plant Cell* 17, 1866–1875. doi: 10.1105/tpc.105.033589
- Fukao, T., and Bailey-Serres, J. (2004). Plant responses to hypoxia—is survival a balancing act? *Trends Plant Sci.* 9, 449–456. doi: 10.1016/j.tplants.2004.07.005
- Gong, B., Li, X., VandenLangenberg, K. M., Wen, D., Sun, S., Wei, M., et al. (2014). Overexpression of S-adenosyl-L-methionine synthetase increased tomato tolerance to alkali stress through polyamine metabolism. *Plant Biotechnol. J.* 12, 694–708. doi: 10.1111/pbi.12173
- Hattori, A., Migitaka, H., Iigo, M., Itoh, M., Yamamoto, K., Ohtani-Kaneko, R., et al. (1995). Identification of melatonin in plants and its effects on plasma melatonin levels and binding to melatonin receptors in vertebrates. *Biochem. Mol. Biol. Int.* 35, 627–634.
- Hossain, Z., López-Clement, M. F., Arbona, V., Pérez-Clemente, R. M., and Gómez-Cadenas, A. (2009). Modulation of the antioxidant system in citrus under waterlogging and subsequent drainage. *J. Plant Physiol.* 166, 1391–1404. doi: 10.1016/j.jplph.2009.02.012
- Kang, K., Lee, K., Park, S., Byeon, Y., and Back, K. (2013). Molecular cloning of rice serotonin N-acetyltransferase, the penultimate gene in plant melatonin biosynthesis. *J. Pineal Res.* 55, 7–13. doi: 10.1111/jpi.12011
- Kuzniak, E., and Sklodowska, M. (2001). Ascorbate, glutathione and related enzymes in chloroplasts of tomato leaves infected by *Botrytis cinerea*. *Plant Sci.* 160, 723–731. doi: 10.1016/S0168-9452(00)00457-X
- Landi, L., Capocasa, F., Costantini, E., and Mezzetti, B. (2009). *ROLC* strawberry plant adaptability, productivity, and tolerance to soil-borne disease and mycorrhizal interactions. *Transgenic Res.* 18, 933–942. doi: 10.1007/s11248-009-9279-7
- Li, C., Liang, B., Chang, C., Wei, Z., Zhou, S., and Ma, F. (2016). Exogenous melatonin improved potassium content in *Malus* under different stress conditions. *J. Pineal Res.* 61, 218–229. doi: 10.1111/jpi.12342
- Li, J., Liu, J., Wang, G., Cha, J. Y., Li, G., Chen, S., et al. (2015). A chaperone function of NO CATALASE ACTIVITY1 is required to maintain catalase activity and for multiple stress responses in Arabidopsis. *Plant Cell* 27, 908–925. doi: 10.1105/tpc.114.135095
- Li, M. Q., Hasan, M. K., Li, C. X., Ahammed, G. J., Xia, X. J., and Shi, K. (2016). Melatonin mediates selenium-induced tolerance to cadmium stress in tomato plants. *J. Pineal Res.* 61, 291–302. doi: 10.1111/jpi.12346
- Li, W., Lu, Q., Hao, N., Zhang, Q., Ge, Q., and Kuang, T. (2000). The high photosynthesis-efficiency characteristics of high-yield varieties in soybean. *Acta Biophys. Sin.* 16, 421–426.
- Li, X., Tan, D. X., Jiang, D., and Liu, F. (2016). Melatonin enhances cold tolerance in drought-primed wild-type and abscisic acid-deficient mutant barley. *J. Pineal Res.* 61, 328–339. doi: 10.1111/jpi.12350
- Liang, C., Zheng, G., Li, W., Wang, Y., Hu, B., Wang, H., et al. (2015). Melatonin delays leaf senescence and enhances salt stress tolerance in rice. *J. Pineal Res.* 59, 91–101. doi: 10.1111/jpi.12243
- Liu, N., Jin, Z. Y., Wang, S. S., Gong, B., Wen, D., Wang, X. F., et al. (2015). Sodic alkaline stress mitigation with exogenous melatonin involves reactive oxygen metabolism and ion homeostasis in tomato. *Sci. Hort.* 181, 18–25. doi: 10.1016/j.scienta.2014.10.049
- López, A., Garcia, J. A., Escames, G., Venegas, C., Ortiz, F., and López, L. C. (2009). Melatonin protects the mitochondria from oxidative damage reducing oxygen consumption, membrane potential, and superoxide anion production. *J. Pineal Res.* 46, 188–198. doi: 10.1111/j.1600-079X.2008.00647.x
- Ma, Q., Zhang, T., Zhang, P., and Wang, Z. Y. (2016). Melatonin attenuates postharvest physiological deterioration of cassava storage roots. *J. Pineal Res.* 60, 424–434. doi: 10.1111/jpi.12325
- Mittler, R., Vanderauwera, S., Gollery, M., and Van, B. F. (2004). Reactive oxygen gene network of plants. *Trends Plant Sci.* 9, 490–498. doi: 10.1016/j.tplants.2004.08.009
- Musgrave, M. E. (1994). Waterlogging effects on yield and photosynthesis in eight winter wheat cultivars. *Crop Sci.* 34, 1314–1318. doi: 10.2135/cropsci1994.0011183X003400050032x
- Palatnik, J. F., Allen, E., Wu, X., Schommer, C., Schwab, R., Carrington, J. C., et al. (2003). Control of leaf morphogenesis by microRNAs. *Nature* 425, 257–263. doi: 10.1038/nature01958
- Palatnik, J. F., Wollmann, H., Schommer, C., Schwab, R., Boisbouvier, J., Rodriguez, R., et al. (2007). Sequence and expression differences underlie functional specialization of *Arabidopsis* microRNAs miR159 and miR319. *Dev. Cell* 13, 115–125. doi: 10.1016/j.devcel.2007.04.012
- Porra, R. J., Thompson, W. A., and Kriedmann, P. E. (1989). Determination of accurate extinction coefficients and simultaneous equations for assaying

## ACKNOWLEDGMENTS

This research was supported by Chinese Universities Scientific Fund (NCET13-0555) and the NSFC (National Natural Science Research Foundation of China, Project No: 31372034).

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fpls.2017.00483/full#supplementary-material>

- chlorophylls a and b extracted with four different solvents: verification of the concentration of chlorophyll standards by atomic absorption spectroscopy. *Biochim. Biophys. Acta* 975, 384–394. doi: 10.1016/S0005-2728(89)80347-0
- Qu, G., Li, X., Zhao, F., Wang, H. X., and Shu, H. R. (1999). Effect of water stress on microstructure of apple leaves and newborn roots. *Acta Hortic. Sin.* 26, 147–151.
- Ramakrishna, A., Giridhar, P., Sankar, K. U., and Ravishankar, G. A. (2012). Melatonin and serotonin profiles in beans of *Coffea* species. *J. Pineal Res.* 52, 470–476. doi: 10.1111/j.1600-079X.2011.00964.x
- Saze, H., Kitayama, J., Takashima, K., Miura, S., Harukawa, Y., Ito, T., et al. (2013). Mechanism for full-length RNA processing of *Arabidopsis* genes containing intragenic heterochromatin. *Nat. Commun.* 4, 2301. doi: 10.1038/ncomms3301
- Shabala, S., Shabala, L., Barcelo, J., and Poschenrieder, C. (2014). Membrane transporters mediating root signalling and adaptive responses to oxygen deprivation and soil flooding. *Plant Cell Environ.* 37, 2216–2233. doi: 10.1111/pce.12339
- Shi, H., Chen, Y., Tan, D. X., Reiter, R. J., Chan, Z., and He, C. (2015a). Melatonin induces nitric oxide and the potential mechanisms relate to innate immunity against bacterial pathogen infection in *Arabidopsis*. *J. Pineal Res.* 59, 102–108. doi: 10.1111/jpi.12244
- Shi, H., Wang, X., Tan, D. X., Reiter, R. J., and Chan, Z. (2015b). Comparative physiological and proteomic analyses reveal the actions of melatonin in the reduction of oxidative stress in Bermuda grass (*Cynodon dactylon* (L.) Pers.). *J. Pineal Res.* 59, 120–131. doi: 10.1111/jpi.12246
- Smethurst, C. F., and Shabala, S. (2003). Screening methods for waterlogging tolerance in lucerne: comparative analysis of waterlogging effects on chlorophyll fluorescence, photosynthesis, biomass and chlorophyll content. *Funct. Plant Biol.* 30, 335–343. doi: 10.1071/FP02192
- Smirnoff, N. (2000). Ascorbic acid: metabolism and functions of a multi-faceted molecule. *Curr. Opin. Plant Biol.* 3, 229–235. doi: 10.1016/S1369-5266(00)00069-8
- Stewart, R. R., and Bewley, J. D. (1980). Lipid peroxidation associated with accelerated aging of soybean axes. *Plant Physiol.* 65, 245–248. doi: 10.1104/pp.65.2.245
- Takahashi, H., Yamauchi, T., Rajhi, I., Nishizawa, N. K., and Nakazono, M. (2015). Transcript profiles in cortical cells of maize primary root during ethylene-induced lysigenous aerenchyma formation under aerobic conditions. *Ann. Bot.* 115, 879–894. doi: 10.1093/aob/mcv018
- Tan, D. X., Chen, L. D., Poeggeler, B., Manchester, L. C., and Reiter, R. J. (1993). Melatonin: a potent, endogenous hydroxyl radical scavenger. *Endocr. J.* 1, 57–60.
- Tan, D. X., Hardeland, R., Manchester, L. C., Korkmaz, A., Ma, S., Rosales-Corral, S., et al. (2012). Functional roles of melatonin in plants, and perspectives in nutritional and agricultural science. *J. Exp. Bot.* 63, 577–597. doi: 10.1093/jxb/err256
- Tan, D. X., Manchester, L. C., Esteban-Zubero, E., Zhou, Z., and Reiter, R. J. (2015). Melatonin as a potent and inducible endogenous antioxidant: synthesis and metabolism. *Molecules* 20, 18886–18906. doi: 10.3390/molecules201018886
- Tan, D. X., Manchester, L. C., Liu, X., Rosales-Corral, S. A., Acuna-Castroviejo, D., and Reiter, R. J. (2013). Mitochondria and chloroplasts as the original sites of melatonin synthesis: a hypothesis related to melatonin's primary function and evolution in eukaryotes. *J. Pineal Res.* 54, 127–138. doi: 10.1111/jpi.12026
- Voesenek, L., and Sasidharan, R. (2013). Ethylene-and oxygen signalling-drive plant survival during flooding. *Plant Biol.* 15, 426–435. doi: 10.1111/plb.12014
- Wang, L., Wang, Z., and Li, X. (2013). Optimization of ultrasonic-assisted extraction of phenolic antioxidants from *Malus baccata* (Linn.) Borkh. using response surface methodology. *J. Sep. Sci.* 36, 1652–1658. doi: 10.1002/jssc.201300062
- Wang, P., Sun, X., Wang, N., Tan, D. X., and Ma, F. (2015). Melatonin enhances the occurrence of autophagy induced by oxidative stress in *Arabidopsis* seedlings. *J. Pineal Res.* 58, 479–489. doi: 10.1111/jpi.12233
- Wang, P., Yin, L. H., Liang, D., Li, C., Ma, F. W., and Yue, Z. Y. (2012). Delayed senescence of apple leaves by exogenous melatonin treatment: toward regulating the ascorbate–glutathione cycle. *J. Pineal Res.* 53, 11–20. doi: 10.1111/j.1600-079X.2011.00966.x
- Wang, W., Vignani, R., Scali, M., and Cresti, M. (2006). A universal and rapid protocol for protein extraction from recalcitrant plant tissues for proteomic analysis. *Electrophoresis* 27, 2782–2786. doi: 10.1002/elps.200500722
- Wei, W., Li, Q. T., Chu, Y. N., Reiter, R. J., Yu, X. M., Zhu, D. H., et al. (2014). Melatonin enhances plant growth and abiotic stress tolerance in soybean plants. *J. Exp. Bot.* 66, 695–707. doi: 10.1093/jxb/eru392
- Xu, Q. T., Yang, L., Zhou, Z. Q., Mei, F. Z., Qu, L. H., and Zhou, G. S. (2013). Process of aerenchyma formation and reactive oxygen species induced by waterlogging in wheat seminal roots. *Planta* 238, 969–982. doi: 10.1007/s00425-013-1947-4
- Xu, W., Cai, S. Y., Zhang, Y., Wang, Y., Ahammed, G. J., Xia, X. J., et al. (2016). Melatonin enhances thermotolerance by promoting cellular protein protection in tomato plants. *J. Pineal Res.* 61, 457–469. doi: 10.1111/jpi.12359
- Yamashita, H., Goto, M., Matsui-Yuasa, I., and Kojima-Yuasa, A. (2015). Ecklonia cava polyphenol has a protective effect against ethanol-induced liver injury in a cyclic AMP-dependent manner. *Mar. Drugs* 13, 3877–3891. doi: 10.3390/md13063877
- Yip, H. K., Chang, Y. C., Wallace, C. G., Chang, L. T., Tsai, T. H., Chen, Y. L., et al. (2013). Melatonin treatment improve adipose-derived mesenchymal stem cell therapy for acute lung ischemia–reperfusion injury. *J. Pineal Res.* 54, 207–221. doi: 10.1111/jpi.12020
- Yu, F., Han, X. S., Geng, C. J., Zhao, Y. X., Zhang, Z. X., and Qiu, F. Z. (2015). Comparative proteomic analysis revealing the complex network associated with waterlogging stress in maize (*Zea mays* L.) seedling root cells. *Proteomics* 15, 135–147. doi: 10.1002/pmic.201400156
- Zhao, H., Su, T., Huo, L., Wei, H., Jiang, Y., and Xu, L. (2015a). Unveiling the mechanism of melatonin impacts on maize seedling growth: sugar metabolism as a case. *J. Pineal Res.* 59, 255–266. doi: 10.1111/jpi.12258
- Zhao, H., Xu, L., Su, T., Jiang, Y., Hu, L., and Ma, F. (2015b). Melatonin regulates carbohydrate metabolism and defenses against *Pseudomonas syringae* pv. *tomato* DC3000 infection in *Arabidopsis thaliana*. *J. Pineal Res.* 59, 109–119. doi: 10.1111/jpi.12245
- Zhao, Y., Tan, D. X., Lei, Q., Chen, H., Wang, L., Li, Q. T., et al. (2013). Melatonin and its potential biological functions in the fruits of sweet cherry. *J. Pineal Res.* 55, 79–88.
- Zheng, X., Tan, D. X., Allan, A. C., Zuo, B., Zhao, Y., Reiter, R. J., et al. (2017). Chloroplastic biosynthesis of melatonin and its involvement in protection of plants from salt stress. *Sci. Rep.* 7:41236. doi: 10.1038/srep41236
- Zuo, B. X., Zheng, X. D., He, P. L., Wang, L., Lei, Q., Feng, C., et al. (2014). Overexpression of *MzASMT* improves melatonin production and enhances drought tolerance in transgenic *Arabidopsis thaliana* plants. *J. Pineal Res.* 57, 408–417. doi: 10.1111/jpi.12180

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Zheng, Zhou, Tan, Wang, Wang, Shan and Kong. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.