



Corrigendum: Earlywood and Latewood Stable Carbon and Oxygen Isotope Variations in Two Pine Species in Southwestern China during the Recent Decades

OPEN ACCESS

Edited and reviewed by:

Jian-Guo Huang, University of Chinese Academy of Sciences, China

*Correspondence:

Ze-Xin Fan fanzexin@xtbg.org.cn Achim Bräuning achim.braeuning@fau.de

Specialty section:

This article was submitted to Functional Plant Ecology, a section of the journal Frontiers in Plant Science

> Received: 14 May 2017 Accepted: 16 May 2017 Published: 31 May 2017

Citation:

Fu P-L, Grießinger J, Gebrekirstos A, Fan Z-X and Bräuning A (2017) Corrigendum: Earlywood and Latewood Stable Carbon and Oxygen Isotope Variations in Two Pine Species in Southwestern China during the Recent Decades. Front. Plant Sci. 8:923. doi: 10.3389/fpls.2017.00923 Pei-Li Fu¹, Jussi Grießinger², Aster Gebrekirstos^{2,3}, Ze-Xin Fan^{1*} and Achim Bräuning^{2*}

¹ Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Menglun, China, ² Institute of Geography, University of Erlangen-Nürnberg, Erlangen, Germany, ³ World Agroforestry Centre, Nairobi, Kenya

Keywords: stable carbon isotope, stable oxygen isotope, intrinsic water use efficiency, subtropical pine species, Asian summer monsoon, intra-annual resolution

A corrigendum on

Earlywood and Latewood Stable Carbon and Oxygen Isotope Variations in Two Pine Species in Southwestern China during the Recent Decades

by Fu, P.-L., Grießinger, J., Gebrekirstos, A., Fan, Z.-X., and Bräuning, A. (2017). Front. Plant Sci. 7:2050. doi: 10.3389/fpls.2016.02050

In the original article, there was a mistake in **Figure 4** as published. The unit of the y-axis in **Figures 4A,B** should be μ mol mol⁻¹, not μ mol s⁻² s⁻¹. The corrected **Figure 4** appears below. The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Fu, Grießinger, Gebrekirstos, Fan and Bräuning. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 4 | Intrinsic water use efficiency (iWUE) for earlywood (iWUE_{EW}, open circle) and latewood of secondary forest pine *P. kesiya* (PIKE) (iWUE_{LW}, closed circle) (**A**), and for earlywood (iWUE_{EW}, open triangle) and latewood of natural forest pine *P. armandii* (PIAR) (iWUE_{LW}, closed triangle) (**B**), intercellular CO₂ concentration (c_i) of earlywood (ci_{EW}, open circle) and latewood of secondary forest pine *P. kesiya* (ci_{LW}, closed circle) (**C**), and for earlywood (ci_{EW}, open triangle) and latewood of natural forest pine Pinus armandii (ci_{LW}, closed triangle) (**D**). Mean values and standard deviations of iWUE and c_i series are indicated inside the panels.