



The Calcium-Dependent Protein Kinase CPK33 Mediates Strigolactone-Induced Stomatal Closure in *Arabidopsis thaliana*

Xuening Wang¹, Shuo Lv¹, Xiangyu Han¹, Xiongjuan Guan¹, Xiong Shi¹, Jingke Kang¹, Luosha Zhang¹, Bing Cao¹, Chen Li², Wei Zhang³, Guodong Wang^{1*} and Yonghong Zhang^{2,4*}

¹ National Engineering Laboratory for Resource Development of Endangered Crude Drugs in Northwest China, Key Laboratory of Medicinal Resources and Natural Pharmaceutical Chemistry, Ministry of Education, College of Life Sciences, Shaanxi Normal University, Xi'an, China, ² Laboratory of Medicinal Plant, Institute of Basic Medical Sciences, School of Basic Medicine, Biomedical Research Institute, Hubei Key Laboratory of Wudang Local Chinese Medicine Research, Hubei University of Medicine, Shiyan, China, ³ Key Laboratory of Plant Development and Environment Adapting Biology, Ministry of Education, School of Life Science, Shandong University, Qingdao, China, ⁴ Hubei Key Laboratory of Embryonic Stem Cell Research, Hubei University of Medicine, Shiyan, China

OPEN ACCESS

Edited by:

Kazimierz Trebacz,
Maria Curie-Skłodowska University,
Poland

Reviewed by:

Fabien Jammes,
Pomona College, United States
Danilo M. Daloso,
Universidade Federal do Ceará, Brazil

*Correspondence:

Guodong Wang
guodong_wang@snnu.edu.cn
Yonghong Zhang
zhangyonghong52@gmail.com

Specialty section:

This article was submitted to
Plant Physiology,
a section of the journal
Frontiers in Plant Science

Received: 27 August 2019

Accepted: 19 November 2019

Published: 17 December 2019

Citation:

Wang X, Lv S, Han X, Guan X, Shi X,
Kang J, Zhang L, Cao B, Li C,
Zhang W, Wang G and Zhang Y (2019)
The Calcium-Dependent Protein
Kinase CPK33 Mediates
Strigolactone-Induced Stomatal
Closure in *Arabidopsis thaliana*.
Front. Plant Sci. 10:1630.
doi: 10.3389/fpls.2019.01630

Strigolactones (SLs) are known to mediate plant acclimation to environmental stress. We recently reported that SLs acted as prominent regulators in promotion of stomatal closure. However, the detailed mechanism by which SLs induce stomatal closure requires further investigation. Here we studied the essential role of the calcium (Ca²⁺) signal mediating by the calcium-dependent protein kinase (CPK) in SL-induced stomatal closure. SL-induced stomatal closure was strongly inhibited by a Ca²⁺ chelator and Ca²⁺ channel blockers, indicating that Ca²⁺ functions in SL promotion of stomatal closure. Through examining a collection of *cpk* mutants, we identified CPK33, potentially acting as a Ca²⁺ transducer, which is implicated in guard cell SL signaling. SL- and Ca²⁺-induced stomatal closure were impaired in *cpk33* mutants. CPK33 kinase activity is essential for SL induction of stomatal closure as SL-induced stomatal closure is blocked in the dead kinase mutant of CPK33. The *cpk33* mutant is impaired in H₂O₂-induced stomatal closure, but not in SL-mediated H₂O₂ production. Our study thus uncovers an important player CPK33 which functions as an essential Ca²⁺ signals mediator in *Arabidopsis* guard cell SL signaling.

Keywords: Ca²⁺, calcium-dependent protein kinase, guard cell, stomatal closure, strigolactones

INTRODUCTION

Plants are sessile organisms that confront with a wide range of biotic and abiotic stress conditions during their life cycle. As a strategy to cope with environmental stress, plants utilize stomatal pores, each consisting of a pair of guard cells, that open and close to modulate gas exchange for photosynthesis, transpirational water loss, and stomatal immunity, thereby allowing plants to respond properly to diverse environmental stress (Hetherington and Woodward, 2003; Ruszala et al., 2011; Blatt et al., 2017; Melotto et al., 2017). Guard cells have developed sophisticated mechanisms which enable plants to appropriately control of stomatal apertures in mediating

response of environmental stimuli such as light, drought, and external calcium ion (Ca^{2+}) (Kim et al., 2010; Murata et al., 2015). Particularly, phytohormones, including abscisic acid (ABA), ethylene, brassinosteroids, strigolactones (SLs), salicylic acid (SA), jasmonic acid (JA), and small signaling peptides have been found to play pivotal roles through their coordination with various key guard cell signaling components to modulate stomatal apertures in response to fluctuating environmental stress (Kim et al., 2010; Daszkowska-Golec and Szarejko, 2013; Munemasa et al., 2015; Murata et al., 2015; Cardinale et al., 2018; Mostofa et al., 2018; Qu et al., 2019; Zhang et al., 2019).

Other than their notable roles in shoot branching (Al-Babili and Bouwmeester, 2015), SLs have been found to be implicated in many plant developmental processes such as primary root development (Kapulnik et al., 2011; Ruyter-Spira et al., 2011), adventitious root formation (Rasmussen et al., 2012; Sun et al., 2015), secondary growth (Agusti et al., 2011), photomorphogenesis (Shen et al., 2007; Shen et al., 2012), flower development (Snowden et al., 2005; Kohlen et al., 2012; Liu et al., 2013), and hypocotyl elongation (Tsuchiya et al., 2010; Jia et al., 2014). Notably, accumulating data indicated that SLs are also involved in mediating plant responses to environmental stress, rendering plants to defend against abiotic stress as well as against specific bacterial and fungal species (Marzec, 2016; Cardinale et al., 2018; Mostofa et al., 2018). Specifically, SL-deficient and SL-signaling mutants exhibited drought hypersensitivity, whereas SLs application rescued drought-sensitive phenotypes of SL-deficient mutants and strengthened drought tolerance of wild-type (WT) plants (Bu et al., 2014; Ha et al., 2014; Liu et al., 2015; Li et al., 2017; Zhang et al., 2018). We recently revealed that SLs could induce stomatal closure through enhancing hydrogen peroxide (H_2O_2) and nitric oxide production in an ABA-independent manner, possibly preventing water loss and pathogen invasion and thereby resulting in plant acclimation to environmental stress (Lv et al., 2018; Zhang et al., 2018). However, the detail molecular mechanism, especially the intracellular events that are initiated by SLs in guard cells, remains largely unclear. To this end, the potential downstream component(s) that transduces guard cell SL signaling is thus required to be determined.

It has long been known that calcium functions as a secondary messenger in stomatal closure (Blatt, 2000; Bowler and Fluhr, 2000; Kim et al., 2010; Murata et al., 2015; Ray, 2017). For instance, through H_2O_2 activation of Ca^{2+} -permeable cation channels, ABA triggers an increment of cytosolic Ca^{2+} [$(\text{Ca}^{2+})_{\text{cyt}}$] that includes Ca^{2+} influx elevation from extracellular spaces and Ca^{2+} release from intracellular stores (Pei et al., 2000). In addition, JA-induced stomatal closure is mediated by cytosolic Ca^{2+} since JA signaling in guard cells is inhibited by Ca^{2+} channel blockers (Suhita et al., 2003; Suhita et al., 2004). Likewise, Ca^{2+} signaling is found to be implicated in SA induction of stomatal closure in a mode of action similar to studies of ABA- and JA-mediated stomatal closure (Prodhon et al., 2018). Eventually, the resultant guard cell cytosolic Ca^{2+} elevation promotes stomatal closure by stimulation of SLOW

ANION CHANNEL-ASSOCIATED 1 (SLAC1) anion channels and/or the GATED OUTWARDLY-RECTIFYING K^+ (GORK) channel (Kim et al., 2010; Murata et al., 2015; Roux and Leonhardt, 2018).

Calcium-dependent protein kinases (CPKs) function as Ca^{2+} signal transducers involving in various biological processes including Ca^{2+} -dependent guard cell signaling (Boudsocq and Sheen, 2013; Schulz et al., 2013; Simeunovic et al., 2016). Mechanically, CPKs are known to activate SLAC1 and GORK channels to induce stomatal closure (Kim et al., 2010; Boudsocq and Sheen, 2013; Schulz et al., 2013; Simeunovic et al., 2016; Corratgé-Faillie et al., 2017). To date, a number of CPKs have been identified to be implicated in guard cell signaling to mediate stomatal movement. Several CPKs, including CPK3, CPK4, CPK6, CPK8, CPK9, CPK10, CPK11, and CPK33, are involved in ABA-mediated stomatal closure through distinct modes of action (Boudsocq and Sheen, 2013; Schulz et al., 2013; Li C.L et al., 2016; Chen et al., 2019). For instance, CPK3 and CPK6 activated ABA-induced stomatal closure and slow-type (S-type) anion channel activity (Mori et al., 2006). Disruption of *CPK6* impaired JA-mediated stomatal closure and S-type anion channels activation, implying that CPK6 acted as a positive regulator in guard cell MeJA signaling (Munemasa et al., 2011). CPK3 and CPK6 functioned additively in SA-induced stomatal closure and SA activation of S-type anion channels (Prodhon et al., 2018). CPK33 suppressed ABA-induced stomatal closure and S-type anion channel activity (Li C.L et al., 2016; Chen et al., 2019), whereas CPK33 stimulated GORK activity to promote stomatal closure (Corratgé-Faillie et al., 2017). It was reported that CPK10, possibly association with HEAT SHOCK PROTEIN 20-LIKE PROTEIN 1 (HSP1), functioned in ABA- and Ca^{2+} -mediated stomatal closure in response to drought stress (Zou et al., 2010). Altogether, these findings underscore the importance of CPKs in the modulation of stomatal closure. CPKs and Ca^{2+} -independent kinases [e.g. SnRK2-type protein kinase OPEN STOMATA1 (OST1)] have long been recognized to be involved in activating ion channels and stimulating stomatal closure (Geiger et al., 2009; Lee et al., 2009; Geiger et al., 2010; Geiger et al., 2011; Brandt et al., 2012; Brandt et al., 2015).

In this study, the importance of Ca^{2+} in SL-induced stomatal closure was firstly determined by performing pharmacological studies, which indicate that Ca^{2+} acts as a prominent mediator functions in SL induction of stomatal closure. Through examining a collection of *cpk* mutants, we identified CPK33 (and possibly CPK10) which is implicated in guard cell SL signaling. SL activation of stomatal closure, as well as Ca^{2+} -induced stomatal closure, was greatly impaired in *cpk33* mutants. Additionally, CPK33 kinase activity is essential for SL induction of stomatal closure. We further found that H_2O_2 -induced stomatal closure was moderately impaired in *cpk33* mutants whereas SL-mediated H_2O_2 production was maintained in *cpk33* mutants. Taken together, our study uncovers an important player CPK33 that functions as an important mediator in SL signaling in *Arabidopsis* guard cells.

MATERIALS AND METHODS

Plant Growth Conditions and Mutants Isolation

The *Arabidopsis* ecotype Columbia-0 (Col-0) was used as wild-type (WT) plants in this study. The following mutants have been described previously: *max1-1*, *max2-1*, *max2-2*, *max3-9*, and *max4-1* (Stirnberg et al., 2002; Umehara et al., 2008), *d14-5* (Yao et al., 2016), *cpk4-1* and *cpk11-2* (Zhu et al., 2007), *cpk8* (Zou et al., 2015), *cpk10* (Zou et al., 2010), *cpk23* (Ma and Wu, 2007), *cpk33-1*, *cpk33-2*, *35S::CPK33* and *35S::CPK33^{K102R}* (Li C.L et al., 2016), *cpk3 cpk5 cpk6 cpk11* (Guzel Deger et al., 2015), and *cpk5 cpk6 cpk11 cpk23* (Wang et al., 2018).

To obtain the double mutant *cpk10 cpk33*, the *cpk10* mutant and *cpk33* mutants (*cpk33-1* and *cpk33-2*) were crossed. Homozygous double mutants were determined by PCRs using a combination of a gene-specific primer and a T-DNA border primer. The primers used were listed in **Table S1**.

Seeds were sterilized and sown for germination in 1/2 Murashige and Skoog (MS) medium supplemented with 0.8% (w/v) agar and 1% sucrose (w/v). Seedlings were transplanted into pots and subsequently kept in a growth chamber with a 16h/8h (light/darkness) regime at 21°C. Fully expanded rosette leaves detached from 4- to 6-week-old healthy plants were harvested for immediate use.

Stomatal Aperture Bioassay

Stomatal apertures were measured as described previously (Lv et al., 2018; Zhang et al., 2019). In brief, epidermal strips of fully expanded leaves were incubated in the MES-KCl buffer (10 mM MES-KOH/50 mM KCl, pH 6.15) to promote stomatal opening following treatments as described in each experiment. Finally, stomatal apertures on the abaxial epidermis were measured and presented as mean \pm SE of three replicates.

Chemicals

The preparation of synthetic SL analog GR24 (Chiralix Nijmegen, the Netherlands) and 2',7'-dichlorofluorescein diacetate (H₂DCF-DA; Biotium Hayward, USA) were performed as described previously (Lv et al., 2018; Zhang et al., 2019). The ethylene glycol-bis(β -aminoethyl ether)-N,N,N',N'-tetra acetic acid (EGTA; Sigma, USA), lanthanum chloride (LaCl₃; Solarbio, China), aluminum chloride (AlCl₃, Solarbio, China), trifluoperazine (TFP), and trifluoperazine dihydrochloride (TFP; Santa Cruz, USA) were dissolved appropriately according to the supplier information and prepared stock solutions for further use. Other chemicals used in this study, including H₂O₂, 2-(N-morpholino) ethanesulphonic acid (MES), and calcium chloride (CaCl₂), were purchased from Sigma-Aldrich. All chemicals used are of the highest analytical grade.

H₂O₂ Content Detection in Guard Cells

H₂DCF-DA was utilized to determine H₂O₂ content in stomata. The measurement of H₂O₂ content was performed according to Lv et al. (2018) and Zhang et al. (2019). Briefly, epidermal strips with open stomata were incubated in the Tris-KCl solution containing 50 μ M H₂DCF-DA for 10 min of darkness at 25°C

in the absence or presence of GR24. Epidermal strips were washed three times with Tris-KCl buffer to remove excess dye under darkness. The fluorescence in stomata was visualized using a TCS SP2 confocal laser scanning microscope (Leica Lasertechnik GmbH, Germany). The fluorescence intensity representing the endogenous H₂O₂ content was determined using ImageJ software. Fluorescence intensities were normalized to those of controls. The data for fluorescence intensities represent the mean \pm SE of three replicates.


Statistical Analyses

Statistical analyses were performed using a one-way ANOVA to discriminate significant differences followed by the least significant difference test.

RESULTS

Ca²⁺ Is Required for SL-Induced Stomatal Closure in *Arabidopsis*

Calcium is known to act as a key secondary messenger in mediating stomatal closure (Blatt, 2000; Bowler and Fluhr, 2000; Pei et al., 2000; Kim et al., 2010; Murata et al., 2015; Ray, 2017). We therefore took a pharmacological approach to determine whether Ca²⁺ is required for SL-triggered stomatal closure. To this end, we examined the effects of ethylene glycol-bis(β -aminoethyl ether)-N,N,N',N'-tetra acetic acid (EGTA; a Ca²⁺ chelator), and LaCl₃ and AlCl₃ (Ca²⁺ channel blockers) on SL-induced stomatal closure (Schwartz, 1985; Zhao et al., 2007; Li Y et al., 2016). As expected, the synthetic SL analog GR24 could significantly induce stomatal closure (**Figure 1**; Lv et al., 2018). However, SL-induced stomatal closure was inhibited by EGTA, LaCl₃, and AlCl₃, respectively (**Figure 1**). These results



indicate that Ca^{2+} and Ca^{2+} channels are involved in SL-induced stomatal closure.

It was reported that exogenous Ca^{2+} could promote stomatal closure (Blatt, 2000; Ray, 2017). Since we have shown that Ca^{2+} plays a crucial role in SL-induced stomatal closure, we further examined the stomatal response of SL-related mutants in response to Ca^{2+} . We found that addition of Ca^{2+} (CaCl_2) induced a similar stomatal closure in SL-related mutants as in WT plants, whereas SL-signaling mutants *max2* and *d14* were insensitive to GR24 as reported previously (Figure S1; Lv et al., 2018). This result indicates that Ca^{2+} might act as a signaling molecule downstream of SL-signaling and SL-biosynthetic genes.

SL- and Ca^{2+} -Induced Stomatal Closure Were Impaired in the *Cpk33* Mutant

Given that Ca^{2+} is essential for SL-induced stomatal closure (Figure 1), we sought to identify the specific Ca^{2+} sensor(s) that is involved in guard-cell SL signaling. CPKs, as Ca^{2+} sensors, are important in the regulation of ABA-mediated and Ca^{2+} -mediated stomatal closure (Kim et al., 2010; Boudsocq and Sheen, 2013; Schulz et al., 2013; Simeunovic et al., 2016). To determine whether any CPK is required for SL signaling in guard cells, we firstly examined SL-induced stomatal closure in the absence and presence of CPK inhibitors TFP and staurosporine (ST) (Li et al., 1998). TFP and ST significantly inhibited GR24-induced stomatal closure (Figure S2), suggesting that CPK(s) is truly indispensable for SL-induced stomatal closure.

Furthermore, to identify CPK(s) that might be involved in SL signaling in guard cells, the effect of GR24 on stomatal aperture was examined in a collection of *cpk* mutants, including *cpk4*, *cpk8*, *cpk10*, *cpk11*, *cpk23*, *cpk33-1*, *cpk33-2*, *cpk3 cpk5 cpk6 cpk11*, and *cpk5 cpk6 cpk11 cpk23* (Figure 2). GR24 significantly induced stomatal closure in most of examined *cpk* mutants. Conversely, SL-induced stomatal closure is greatly impaired in *cpk33-1* and *cpk33-2* mutants, and marginally impaired in *cpk10* mutants (Figure 2). To overcome the potential redundant function

between CPK10 and CPK33, we generated double mutants *cpk10 cpk33-1* and *cpk10 cpk33-2*. The double mutants *cpk10 cpk33* exhibited insensitivity to SL-induced stomatal closure to a similar extent to that of the *cpk33* mutants (Figure 2), suggesting that CPK10 exerted slight effect on SL-induced stomatal closure. Nevertheless, we found that CPK33 is predominantly involved in guard cell SL signaling, and we thus concentrated on CPK33 in subsequent studies.

Because SL promotion of stomatal closure depends on Ca^{2+} , we hypothesized that disruption of CPK33 would impair Ca^{2+} signaling transduction. To test this hypothesis, we investigated the stomatal response following exogenous Ca^{2+} application in *cpk33-1* and *cpk33-2* mutants. Exogenous Ca^{2+} stimulated stomatal closure in WT plants, but not in *cpk33-1* and *cpk33-2* mutants (Figure 3). Taken together, our results indicate that CPK33 acts as an important Ca^{2+} sensor that is involved in guard-cell SL signaling.

CPK33 Kinase Activity Is Essential for SL-Induced Stomatal Closure

Previous studies have shown that the kinase activity of CPK33 is essential for SLAC1 channels and GORK channels activities and ABA-induced stomatal movement (Li C.L et al., 2016; Corratgé-Faillie et al., 2017). To investigate whether the *in vivo* kinase activity of CPK33 is also required for SL-induced stomatal closure, we performed stomatal bioassay analysis upon GR24 treatment using *cpk33-1* and *cpk33-2* mutants, two independent *cpk33-1* complementation lines (35::CPK33 *cpk33* #1 and #2), and two independent lines expressing a kinase-inactive CPK33^{K102R} construct in *cpk33-1* (35::CPK33^{K102R} *cpk33* #1 and #2). Two 35::CPK33 *cpk33* lines exhibited GR24-sensitive phenotype similar to that of WT plants, while we found that 35::CPK33^{K102R} *cpk33* plants, as well as *cpk33-1* and *cpk33-2* mutants, failed to close stomata in response to GR24 (Figure 4). Thus, this result suggests that CPK33 kinase activity is essential for stomatal closure induced by SLs.




FIGURE 2 | Strigolactone-stimulated stomatal closure is mediated by CPK33. Leaf epidermal peels of WT plants and a collection of *cpk* mutants were exposed to the MES-KCl buffer in the absence and presence of 1 μM GR24. Stomatal apertures were measured and presented as means \pm SE of three independent experiments. Bars with different letters represent statistically significant differences.




FIGURE 3 | Ca²⁺-induced stomatal closure is significantly impaired in *cpk33* mutants. Leaves of WT plants, *cpk33-1*, and *cpk33-2* mutants with open stomata were exposed to the MES-KCl buffer containing different concentrations of CaCl₂. Stomatal apertures were subsequently measured and presented as means ± SE of three independent experiments. Bars with different letters represent statistically significant differences.




FIGURE 4 | CPK33 kinase activity is essential for strigolactone-induced stomatal closure. Leaves of WT, *cpk33-1*, *cpk33-2*, 35S::CPK33 *cpk33* #1, 35S::CPK33 *cpk33* #2, 35S::CPK33^{K102R} *cpk33* #1, and 35S::CPK33^{K102R} *cpk33* #2 plants with open stomata were exposed to the MES-KCl buffer in the absence or presence of 1 µM GR24. Stomatal apertures were measured and presented as means ± SE of three independent experiments. Bars with different letters represent statistically significant differences.

Effects of the *cpk33* Mutation on SL-Induced H₂O₂ Production and H₂O₂-Mediated Stomatal Closure in Guard Cells

It was reported previously that SLs stimulate H₂O₂ accumulation, and the resultant H₂O₂ acts as an early signal component in the induction of stomatal closure triggered by SLs (Lv et al., 2018). To investigate the genetic relationship between H₂O₂ and CPK33 in guard cell SL signaling, we examined the SL effect on H₂O₂

production in *cpk33* mutants using the H₂O₂ fluorescent probe H₂DCF-DA. In line with previous data, GR24 stimulated H₂O₂ production in WT plants. Similarly, GR24 also stimulated H₂O₂ production in *cpk33* mutants (Figure 5A), suggesting that CPK33 disruption did not affect SL-induced H₂O₂ production.

To further investigate the position of H₂O₂ production and CPK33 in guard cell SL signaling pathway, we next examined the stomatal aperture upon exogenous H₂O₂ in *cpk33* mutants. Exogenous application of H₂O₂ significantly stimulated stomatal closure in WT plants, but H₂O₂-induced stomatal closure is partially impaired in *cpk33* mutants in comparison with WT plants (Figure 5B). Altogether, our results indicate that CPK33 possibly functions downstream of H₂O₂ production in guard-cell SL signaling. It appears that, as a paradigm, H₂O₂ activates CPK(s) that function as signal mediator(s) in guard cells. However, it remains to be explored whether and how CPK33 is activated by SL-induced H₂O₂

DISCUSSION

It has been shown that SLs were implicated in various developmental processes of plants and mediate their responses to environmental stress (Al-Babili and Bouwmeester, 2015; Marzec, 2016; Cardinale et al., 2018; Mostofa et al., 2018). Recently we found that SLs function as common regulators to induce stomatal closure (Lv et al., 2018; Zhang et al., 2018). We further elucidated that SL-induced stomatal closure is accomplished through enhancing production of H₂O₂ and nitric oxide, eventually promoting plant resilience to environmental stress (Lv et al., 2018; Zhang et al., 2018). However, no significant effect on stomatal aperture was observed when spraying GR24 onto intact plants, although SL-related mutants were genetically confirmed to display higher stomatal conductance (Kalliola et al., 2019). The discrepancy in SL-induced stomatal closure may be due to the different materials used, intact plants and epidermal strips, which possibly results in different efficacy of GR24 such as permeability problem. It has been reported that Ca²⁺ and its sensors CPKs are crucial for stomatal closing (Blatt, 2000; Bowler and Fluhr, 2000; Pei et al., 2000; Kim et al., 2010; Murata et al., 2015; Ray, 2017). OST1, a Ca²⁺-independent kinase, was found to be dispensable for SL-induced stomatal closure (Lv et al., 2018). Based on these observations, we thus hypothesize that Ca²⁺ as well as its transducers CPK(s) is implicated in SL-triggered stomatal closure.

In this study we found that Ca²⁺ chelator EGTA, and Ca²⁺ channel blockers LaCl₃ and AlCl₃ significantly suppressed SL-induced stomatal closure (Figure 1), suggesting the requirement of extracellular free Ca²⁺ and Ca²⁺ channels in the modulation of SL-induced stomatal closure. Our pharmacological and further genetic analyses suggest that SL requires cytosolic Ca²⁺ signals to promote stomatal closure (Figures 1 and 3). It is possible that SLs stimulate either Ca²⁺ sensitivity priming or cytosolic Ca²⁺ elevation to perturb Ca²⁺ signals, thereby resulting in stomatal closure. It is thus demanded to detect the alteration of Ca²⁺ signals through Ca²⁺ detection fluorescence dyes and/or different Ca²⁺ biosensors in the future analyses.

It has long been known that CPKs, acting as Ca²⁺ sensors, are important mediators of Ca²⁺-dependent stomatal closure and ion





FIGURE 5 | The role of CPK33 in strigolactone-induced H_2O_2 production and H_2O_2 -mediated stomatal closure in guard cells. **(A)** Effects of GR24 on H_2O_2 production in guard cells of WT plants, *cpk33-1* and *cpk33-2* mutants. Fluorescence intensities were normalized to those of controls that were taken as 100% for the indicated experiments. **(B)** Epidermal strips of WT plants with open stomata were exposed to the MES-KCl buffer containing different concentrations of H_2O_2 . The stomatal apertures were measured and presented as means \pm SE of three independent experiments. Bars with different letters represent statistically significant differences.

channel activation (Kim et al., 2010; Boudsocq and Sheen, 2013; Schulz et al., 2013; Simeunovic et al., 2016). We identified a potential Ca^{2+} transducer CPK33 acting as an intermediate component downstream of H_2O_2 in guard cell SL signaling pathway. SL activation of stomatal closure, as well as Ca^{2+} -induced stomatal closure, were impaired in *cpk33* mutants (Figures 2 and 3). Interestingly, CPK10 disruption slightly impaired SL-induced stomatal closure (Figure 2). The double mutant *cpk10 cpk33* exhibited insensitivity to SL-induced stomatal closure to a similar extent to that of the *cpk33* mutant (Figure 2), suggesting a prime role of CPK33 and a differential contribution of CPK10 and CPK33 in SL-induced stomatal closure. It was found previously that CPK10 was involved in plant responses to drought stress *via* modulation of ABA- and Ca^{2+} -mediated stomatal closure (Zou et al., 2010). Considering the complexity of crosstalk of distinct signaling pathways, it is plausible that CPK10 might indirectly impact on the SL activation of stomatal closure. Alternatively, given that there are 34 CPKs in *Arabidopsis* (Boudsocq and Sheen, 2013), a yet unidentified CPK, rather than

CPK33, may function redundant with CPK10 in modulation of SL-induced stomatal closure. Nevertheless, the role of CPK10 (possible with another redundant CPK) in SL promotion of stomatal closure needs to be investigated.

Intriguingly, guard cell outward potassium channel GORK is specifically stimulated by CPK33 to active outward potassium ion currents, showing that, unlike its negative regulation of anion channels (Li C.L et al., 2016), CPK33 positively modulates the GORK channel activity to promote stomatal closure (Corratgé-Faillie et al., 2017). Consequently, *cpk33* mutants were delayed in stomatal closure under normal conditions (Corratgé-Faillie et al., 2017). In this regard, it is possible that SLs could exploit CPK33-activated GORK channels to promote stomatal closure. It thus remains to be investigated whether SLs could stimulate CPK33-activated GORK channel activity. CPK33 was reported as a negative regulator in ABA-modulated stomatal closure (Li C.L et al., 2016). SL activation of stomatal closure, however, were inhibited in *cpk33* mutants (Figure 2), indicating that CPK33 positively functions in SL-induced stomatal closure. Taken together, our results and



previous reports suggest that the CPK-dependent Ca²⁺ recognition conveying by CPK33 could be essential for SL signaling, as well as ABA signaling in guard cells. Nevertheless, the role of opposite effect of CPK33 on anion and potassium channels, as well as the discrepant role of CPK33 in guard cell ABA and SL signaling, is required to be further studied.

Integrated previous studies with our present genetic and physiological analyses, we propose that CPK33, as a Ca²⁺ transducer, acts downstream of H₂O₂ and Ca²⁺ (Figure 6). In guard cells, following perception by D14 and MAX2, SLs stimulate the production of H₂O₂ that possibly activates CPK33, which likely modulates anion and potassium channels to promote stomatal closure (Li C.L et al., 2016; Corratgé-Faillie et al., 2017; Figure 6). Indeed, it has been reported that CPK33 positively stimulates the GORK channels while negatively regulates SLAC1 channels to promote stomatal closure (Li C.L et al., 2016; Corratgé-Faillie

et al., 2017). However, it remains to be examined whether and how CPK33 is activated by SL-induced H₂O₂. It has been reported that H₂O₂-stimulated cytosolic Ca²⁺ elevation is crucial for stomatal closure (Pei et al., 2000). CPK33 was confirmed to be a typical Ca²⁺-dependent kinase (Corratgé-Faillie et al., 2017), indicating that CPK33 activity requires cytosolic Ca²⁺ increase. Therefore, it will be intriguing to investigate whether SLs and/or SL-induced H₂O₂ are able to promote cytosolic Ca²⁺ which presumably is sensed by CPK33 in SL-mediated guard cell signaling. The finding that the stomata of *cpk33* mutants remain considerable response to H₂O₂ indicates that other factor(s) might also required for SL-triggered stomatal closure. In the meanwhile, our results emphasizes the essential role of H₂O₂ serving as a hub in the complicated hormone-mediated stomatal closure. Altogether, our study reinforces the understanding of the molecular mechanism by which SLs induce stomatal closure and provides new insights to improve stress acclimatization of plants.

DATA AVAILABILITY STATEMENT

The datasets generated for this study are available on request to the corresponding author.

AUTHOR CONTRIBUTIONS

YZ and GW designed the experiments. XW, SL, XH, XG, XS, JK, ZL, and YZ performed the experiments. BC assisted in microscope. WZ contributed materials. YZ, CL, and GW wrote the manuscript. All authors reviewed the manuscript.

FUNDING

The study is supported by the National Natural Science Foundation of China (31701294 and 31801210), the Fundamental Research Funds for Central Universities (GK201702016), the SNNU Laboratory Technology Research Funds (SYJS201739), the Start-up Foundation of Hubei University of Medicine (2016QDJZR14), and the Natural Science Foundation of Hubei Provincial Department of Education (Q20182104).

ACKNOWLEDGMENTS

We thank Profs Wei-Hua Wu, Da-Peng Zhang, Rob Roelfsema, Daoxin Xie, Cun Wang, and Shinjiro Yamaguchi for sharing genetic materials.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2019.01630/full#supplementary-material>

REFERENCES

- Agusti, J., Herold, S., Schwarz, M., Sanchez, P., Ljung, K., Dun, E. A., et al. (2011). Strigolactone signaling is required for auxin-dependent stimulation of secondary growth in plants. *Proc. Natl. Acad. Sci. U.S.A.* 108, 20242–20247. doi: 10.1073/pnas.1111902108
- Al-Babili, S., and Bouwmeester, H. J. (2015). Strigolactones, a novel carotenoid-derived plant hormone. *Annu. Rev. Plant Biol.* 66, 161–186. doi: 10.1146/annurev-arplant-043014-114759
- Blatt, M. R., Brodribb, T. J., and Torii, K. U. (2017). Small pores with a big impact. *Plant Physiol.* 174, 467–469. doi: 10.1104/pp.17.00642
- Blatt, M. R. (2000). Ca^{2+} signalling and control of guard-cell volume in stomatal movements. *Curr. Opin. Plant Biol.* 3, 196–204. doi: 10.1016/s1369-5266(00)80065-5
- Boudsocq, M., and Sheen, J. (2013). CDPKs in immune and stress signaling. *Trends Plant Sci.* 18, 30–40. doi: 10.1016/j.tplants.2012.08.008
- Bowler, C., and Fluhr, R. (2000). The role of calcium and activated oxygens as signals for controlling cross-tolerance. *Trends Plant Sci.* 5, 241–246. doi: 10.1016/s1360-1385(00)01628-9
- Brandt, B., Brodsky, D. E., Xue, S., Negi, J., Iba, K., Kangasjärvi, J., et al. (2012). Reconstitution of abscisic acid activation of SLAC1 anion channel by CPK6 and OST1 kinases and branched ABI1 PP2C phosphatase action. *Proc. Natl. Acad. Sci. U.S.A.* 109, 10593–10598. doi: 10.1073/pnas.1116590109
- Brandt, B., Munemasa, S., Wang, C., Nguyen, D., Yong, T., Yang, P. G., et al. (2015). Calcium specificity signaling mechanisms in abscisic acid signal transduction in *Arabidopsis* guard cells. *eLife* 4, e03599. doi: 10.7554/eLife.03599
- Bu, Q., Lv, T., Shen, H., Luong, P., Wang, J., Wang, Z., et al. (2014). Regulation of drought tolerance by the F-box protein MAX2 in *Arabidopsis*. *Plant Physiol.* 164, 424–439. doi: 10.1104/pp.113.226837
- Cardinale, F., Krukowski, P. K., Schubert, A., and Visentin, I. (2018). Strigolactones: mediators of osmotic stress responses with a potential for agrochemical manipulation of crop resilience. *J. Exp. Bot.* 69, 2291–2303. doi: 10.1093/jxb/erx494
- Chen, D. H., Liu, H. P., and Li, C. L. (2019). Calcium-dependent protein kinase CPK9 negatively functions in stomatal abscisic acid signaling by regulating ion channel activity in *Arabidopsis*. *Plant Mol. Biol.* 99, 113–122. doi: 10.1007/s11103-018-0805-y
- Corratgé-Faillie, C., Ronzier, E., Sanchez, F., Prado, K., Kim, J. H., Lanciano, S., et al. (2017). The *Arabidopsis* guard cell outward potassium channel GORK is regulated by CPK33. *FEBS Lett.* 591, 1982–1992. doi: 10.1002/1873-3468.12687
- Daszkowska-Golec, A., and Szarejko, I. (2013). Open or close the gate-stomata action under the control of phytohormones in drought stress conditions. *Front. Plant Sci.* 4, 138–138. doi: 10.3389/fpls.2013.00138
- Geiger, D., Scherzer, S., Mumm, P., Stange, A., Marten, I., Bauer, H., et al. (2009). Activity of guard cell anion channel SLAC1 is controlled by drought stress signaling kinase-phosphatase pair. *Proc. Natl. Acad. Sci. U.S.A.* 106, 21425–21430. doi: 10.1073/pnas.0912021106
- Geiger, D., Scherzer, S., Mumm, P., Marten, I., Ache, P., Matschi, S., et al. (2010). Guard cell anion channel SLAC1 is regulated by CDPK protein kinases with distinct Ca^{2+} affinities. *Proc. Natl. Acad. Sci. U.S.A.* 107, 8023–8028. doi: 10.1073/pnas.0912030107
- Geiger, D., Maierhofer, T., Al-Rasheid, K. A., Scherzer, S., Mumm, P., Liese, A., et al. (2011). Stomatal closure by fast abscisic acid signaling is mediated by the guard cell anion channel SLAH3 and the receptor RCAR1. *Sci. Signal.* 4, ra32. doi: 10.1126/scisignal.2001346
- Guzel Deger, A., Scherzer, S., Nuhkat, M., Kedzierska, J., Kollist, H., Brosché, M., et al. (2015). Guard cell SLAC1-type anion channels mediate flagellin-induced stomatal closure. *New Phytol.* 208, 162–173. doi: 10.1111/nph.13435
- Ha, C. V., Leyva-González, M. A., Osakabe, Y., Tran, U. T., Nishiyama, R., Watanabe, Y., et al. (2014). Positive regulatory role of strigolactone in plant responses to drought and salt stress. *Proc. Natl. Acad. Sci. U.S.A.* 111, 851–856. doi: 10.1073/pnas.1322135111
- Hetherington, A. M., and Woodward, F. I. (2003). The role of stomata in sensing and driving environmental change. *Nature* 424, 901–908. doi: 10.1038/nature01843
- Jia, K. P., Luo, Q., He, S. B., Lu, X. D., and Yang, H. Q. (2014). Strigolactone-regulated hypocotyl elongation is dependent on cryptochrome and phytochrome signaling pathways in *Arabidopsis*. *Mol. Plant* 7, 528–540. doi: 10.1093/mp/sst093
- Kalliola, M., Jakobson, L., Davidsson, P., Pennanen, V., Waszczak, C., Yarmolinsky, D., et al. (2019). The role of strigolactones in regulation of stomatal conductance and plant-pathogen interactions in *Arabidopsis thaliana*. *BioRxiv*. doi: 10.1101/573873
- Kapulnik, Y., Delaux, P. M., Resnick, N., Mayzlish-Gati, E., Winer, S., Bhattacharya, C., et al. (2011). Strigolactones affect lateral root formation and root-hair elongation in *Arabidopsis*. *Planta* 233, 209–216. doi: 10.1007/s00425-010-1310-y
- Kim, T. H., Böhmer, M., Hu, H., Nishimura, N., and Schroeder, J. I. (2010). Guard cell signal transduction network: advances in understanding abscisic acid, CO_2 and Ca^{2+} signaling. *Annu. Rev. Plant Biol.* 61, 561–591. doi: 10.1146/annurev-arplant-042809-112226
- Kohlen, W., Charnikhova, T., Lammers, M., Pollina, T., Tóth, P., Haider, I., et al. (2012). The tomato *CAROTENOID CLEAVAGE DIOXYGENASE (SICCD8)* regulates rhizosphere signaling, plant architecture and affects reproductive development through strigolactone biosynthesis. *New Phytol.* 196, 535–547. doi: 10.1111/j.1469-8137.2012.04265.x
- Lee, S. C., Lan, W. Z., Buchanan, B. B., and Luan, S. (2009). A protein kinase-phosphatase pair interacts with an ion channel to regulate ABA signaling in plant guard cells. *Proc. Natl. Acad. Sci. U.S.A.* 106, 21419–21424. doi: 10.1073/pnas.0910601106
- Li, J., Lee, Y. R., and Assmann, S. M. (1998). Guard cells possess a Calcium-dependent protein kinase that phosphorylates the KAT1 potassium channel. *Plant Physiol.* 116, 785–795. doi: 10.1104/pp.116.2.785
- Li, C. L., Wang, M., Wu, X. M., Chen, D. H., Lv, H. J., Shen, J. L., et al. (2016). TH11, a thiamine thiazole synthase, interacts with Ca^{2+} -dependent protein kinase CPK33 and modulates the S-type anion channels and stomatal closure in *Arabidopsis*. *Plant Physiol.* 170, 1090–1104. doi: 10.1104/pp.15.01649
- Li, Y., Xu, S., Gao, J., Pan, S., and Wang, G. (2016). Glucose- and mannose-induced stomatal closure is mediated by ROS production, Ca^{2+} and water channel in *Vicia faba*. *Physiol. Plant* 156, 252–261. doi: 10.1111/ppl.12353
- Li, W., Nguyen, K. H., Chu, H. D., Ha, C. V., Watanabe, Y., Osakabe, Y., et al. (2017). The karrikin receptor KAI2 promotes drought resistance in *Arabidopsis thaliana*. *PLoS Genet.* 13, e1007076. doi: 10.1371/journal.pgen.1007076
- Liu, J., Novero, M., Charnikhova, T., Ferrandino, A., Schubert, A., Ruyter-Spira, C., et al. (2013). *CAROTENOID CLEAVAGE DIOXYGENASE 7* modulates plant growth, reproduction, senescence, and determinate nodulation in the model legume *Lotus japonicus*. *J. Exp. Bot.* 64, 1967–1981. doi: 10.1093/jxb/ert056
- Liu, J., He, H., Vitali, M., Visentin, I., Charnikhova, T., Haider, I., et al. (2015). Osmotic stress represses strigolactone biosynthesis in *Lotus japonicus* roots: exploring the interaction between strigolactones and ABA under abiotic stress. *Planta* 241, 1435–1451. doi: 10.1007/s00425-015-2266-8
- Lv, S., Zhang, Y., Li, C., Liu, Z., Yang, N., Pan, L., et al. (2018). Strigolactone-triggered stomatal closure requires hydrogen peroxide synthesis and nitric oxide production in an abscisic acid-independent manner. *New Phytol.* 217, 290–304. doi: 10.1111/nph.14813
- Ma, S. Y., and Wu, W. H. (2007). AtCPK23 functions in *Arabidopsis* responses to drought and salt stresses. *Plant Mol. Biol.* 65, 511–518. doi: 10.1007/s11103-007-9187-2
- Marzec, M. (2016). Strigolactones as part of the plant defence system. *Trends Plant Sci.* 21, 900–903. doi: 10.1016/j.tplants.2016.08.010
- Melotto, M., Zhang, L., Oblessuc, P. R., and He, S. Y. (2017). Stomatal defense a decade later. *Plant Physiol.* 174, 561–571. doi: 10.1104/pp.16.01853
- Mori, I. C., Murata, Y., Yang, Y., Munemasa, S., Wang, Y. F., Andreoli, S., et al. (2006). CDPKs CPK6 and CPK3 function in ABA regulation of guard cell S-type anion- and Ca^{2+} -permeable channel and stomatal closure. *PLoS Biol.* 4, e327. doi: 10.1371/journal.pbio.0040327
- Mostafa, M. G., Li, W., Nguyen, K. H., Fujita, M., and Tran, L. P. (2018). Strigolactones in plant adaptation to abiotic stresses: an emerging avenue of plant research. *Plant Cell Environ.* 41, 2227–2243. doi: 10.1111/pce.13364
- Munemasa, S., Hossain, M. A., Nakamura, Y., Mori, I. C., and Murata, Y. (2011). The *Arabidopsis* calcium-dependent protein kinase, CPK6, functions as a positive regulator of methyl jasmonate signaling in guard cells. *Plant Physiol.* 155, 553–561. doi: 10.1104/pp.110.162750
- Munemasa, S., Hauser, F., Park, J., Waadt, R., Brandt, B., and Schroeder, J. I. (2015). Mechanisms of abscisic acid-mediated control of stomatal aperture. *Curr. Opin. Plant Biol.* 28, 154–162. doi: 10.1016/j.pbi.2015.10.010

- Murata, Y., Mori, I. C., and Munemasa, S. (2015). Diverse stomatal signaling and the signal integration mechanism. *Annu. Rev. Plant Biol.* 66, 369–392. doi: 10.1146/annurev-arplant-043014-114707
- Pei, Z. M., Murata, Y., Benning, G., Thomine, S., Klüsener, B., Allen, G. J., et al. (2000). Calcium channels activated by hydrogen peroxide mediate abscisic acid signalling in guard cells. *Nature* 406, 731–734. doi: 10.1038/35021067
- Prodhan, M. Y., Munemasa, S., Nahar, M. N., Nakamura, Y., and Murata, Y. (2018). Guard cell salicylic acid signaling is integrated into abscisic acid signaling via the Ca^{2+} /CPK-dependent pathway. *Plant Physiol.* 178, 441–450. doi: 10.1104/pp.18.00321
- Qu, X., Cao, B., Kang, J., Wang, X., Han, X., Jiang, W., et al. (2019). Fine-tuning stomatal movement through small signaling peptides. *Front. Plant Sci.* 10, 69. doi: 10.3389/fpls.2019.00069
- Rasmussen, A., Mason, M. G., Cuyper, C. D., Brewer, P. B., Herold, S., Agusti, J., et al. (2012). Strigolactones suppress adventitious rooting in *Arabidopsis* and pea. *Plant Physiol.* 158, 1976–1987. doi: 10.1104/pp.111.187104
- Ray, S. D. (2017). “ Ca^{2+} , The miracle molecule in plant hormone signaling during abiotic stress,” in *Mechanism of Plant Hormone Signaling under Stress*. Eds. G. K. Pandey (New Jersey, NJ: Wiley Blackwell), 45–90.
- Roux, B., and Leonhardt, N. (2018). Chapter Seven - The regulation of Ion channels and transporters in the guard cell. *Adv. Bot. Res.* 87, 171–214. doi: 10.1016/bs.abr.2018.09.013
- Ruszala, E. M., Beerling, D. J., Franks, P. J., Chater, C., Casson, S. A., Gray, J. E., et al. (2011). Land plants acquired active stomatal control early in their evolutionary history. *Curr. Biol.* 21, 1025–1029. doi: 10.1016/j.cub.2011.04.044
- Ruyter-Spira, C., Kohlen, W., Charnikhova, T., van Zeijl, A., van Bezouwen, L., de Ruijter, N., et al. (2011). Physiological effects of the synthetic strigolactone analog GR24 on root system architecture in *Arabidopsis*: another belowground role for strigolactones? *Plant Physiol.* 155, 721–734. doi: 10.1104/pp.110.166645
- Schulz, P., Herde, M., and Romeis, T. (2013). Calcium-dependent protein kinases: hubs in plant stress signaling and development. *Plant Physiol.* 163, 523–530. doi: 10.1104/pp.113.222539
- Schwartz, A. (1985). Role of Ca^{2+} and EGTA on Stomatal Movements in *Commelina communis* L. *Plant Physiol.* 79, 1003–1005. doi: 10.1104/pp.79.41003
- Shen, H., Luong, P., and Huq, E. (2007). The F-box protein MAX2 functions as a positive regulator of photomorphogenesis in *Arabidopsis*. *Plant Physiol.* 145, 1471–1483. doi: 10.1104/pp.107.107227
- Shen, H., Zhu, L., Bu, Q. Y., and Huq, E. (2012). MAX2 affects multiple hormones to promote photomorphogenesis. *Mol. Plant* 5, 750–762. doi: 10.1093/mp/sss029
- Simeunovic, A., Mair, A., Wurzing, B., and Teige, M. (2016). Know where your clients are: subcellular localization and targets of calcium-dependent protein kinases. *J. Exp. Bot.* 67, 3855–3872. doi: 10.1093/jxb/erw157
- Snowden, K. C., Simkin, A. J., Janssen, B. J., Templeton, K. R., Loucas, H. M., Simons, J. L., et al. (2005). The *Decreased apical dominance1/Petunia hybrida* CAROTENOID CLEAVAGE DIOXYGENASE8 gene affects branch production and plays a role in leaf senescence, root growth, and flower development. *Plant Cell* 17, 746–759. doi: 10.1105/tpc.104.027714
- Stirnberg, P., van De Sande, K., and Leyser, H. M. (2002). MAX1 and MAX2 control shoot lateral branching in *Arabidopsis*. *Development* 129, 1131–1141.
- Suhita, D., Kolla, V. A., Vavasseur, A., and Raghavendra, A. S. (2003). Different signaling pathways involved during the suppression of stomatal opening by methyl jasmonate or abscisic acid. *Plant Sci.* 164, 481–488. doi: 10.1016/s0168-9452(02)00432-6
- Suhita, D., Raghavendra, A. S., Kwak, J. M., and Vavasseur, A. (2004). Cytoplasmic alkalization precedes reactive oxygen species production during methyl jasmonate- and abscisic acid-induced stomatal closure. *Plant Physiol.* 134, 1536–1545. doi: 10.1104/pp.103.032250
- Sun, H., Tao, J., Hou, M., Huang, S., Chen, S., Liang, Z., et al. (2015). A strigolactone signal is required for adventitious root formation in rice. *Ann. Bot.* 115, 1155–1162. doi: 10.1093/aob/mcv052
- Tsuchiya, Y., Vidaurre, D., Toh, S., Hanada, A., Nambara, E., Kamiya, Y., et al. (2010). A small-molecule screen identifies new functions for the plant hormone strigolactone. *Nat. Chem. Biol.* 6, 741–749. doi: 10.1038/nchembio.435
- Umehara, M., Hanada, A., Yoshida, S., Akiyama, K., Arite, T., Takeda-Kamiya, N., et al. (2008). Inhibition of shoot branching by new terpenoid plant hormones. *Nature* 455, 195–200. doi: 10.1038/nature07272
- Wang, C., Zhang, J., Wu, J., Brodsky, D. E., and Schroeder, J. I. (2018). Cytosolic malate and oxaloacetate activate S-type anion channels in *Arabidopsis* guard cells. *New Phytol.* 220, 178–186. doi: 10.1111/nph.15292
- Yao, R., Ming, Z., Yan, L., Li, S., Wang, F., Ma, S., et al. (2016). DWARF14 is a non-canonical hormone receptor for strigolactone. *Nature* 536, 469–473. doi: 10.1038/nature19073
- Zhang, Y., Lv, S., and Wang, G. (2018). Strigolactones are common regulators in induction of stomatal closure in planta. *Plant Signal. Behav.* 13, 1–12. doi: 10.1080/15592324.2018.1444322
- Zhang, L., Shi, X., Zhang, Y., Wang, J., Yang, J., Ishida, T., et al. (2019). CLE9 peptide-induced stomatal closure is mediated by abscisic acid, hydrogen peroxide, and nitric oxide in *Arabidopsis thaliana*. *Plant Cell Environ.* 42, 1033–1044. doi: 10.1111/pce.13475
- Zhao, M. G., Tian, Q. Y., and Zhang, W. H. (2007). Ethylene activates a plasma membrane Ca^{2+} -permeable channel in tobacco suspension cells. *New Phytol.* 174, 507–515. doi: 10.2307/4640974
- Zhu, S. Y., Yu, X. C., Wang, X. J., Zhao, R., Li, Y., Fan, R. C., et al. (2007). Two calcium-dependent protein kinases, CPK4 and CPK11, regulate abscisic acid signal transduction in *Arabidopsis*. *Plant Cell* 19, 3019–3036. doi: 10.1105/tpc.107.050666
- Zou, J. J., Wei, F. J., Wang, C., Wu, J. J., Ratnasekera, D., Liu, W. X., et al. (2010). *Arabidopsis* calcium-dependent protein kinase CPK10 functions in abscisic acid- and Ca^{2+} -mediated stomatal regulation in response to drought stress. *Plant Physiol.* 154, 1232–1243. doi: 10.1104/pp.110.157545
- Zou, J. J., Li, X. D., Ratnasekera, D., Wang, C., Liu, W. X., Song, L. F., et al. (2015). *Arabidopsis* CALCIUM-DEPENDENT PROTEIN KINASE8 and CATALASE3 function in abscisic acid-mediated signaling and H_2O_2 homeostasis in stomatal guard cells under drought stress. *Plant Cell* 27, 1445–1460. doi: 10.1105/tpc.15.00144

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Wang, Lv, Han, Guan, Shi, Kang, Zhang, Cao, Li, Zhang, Wang and Zhang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.