Check for updates

OPEN ACCESS

EDITED BY Nadine Töpfer, University of Cologne, Germany

REVIEWED BY

Anna Maria Mastrangelo, Council for Agricultural and Economics Research (CREA), Italy Matthias Salomon, University of Adelaide, Australia Patrick Schäfer, University of Giessen, Germany

*CORRESPONDENCE Gwendolin Wehner Gwendolin.wehner@juliuskuehn.de

SPECIALTY SECTION

This article was submitted to Plant Systems and Synthetic Biology, a section of the journal Frontiers in Plant Science

RECEIVED 13 October 2022 ACCEPTED 12 December 2022 PUBLISHED 12 January 2023

CITATION

Matros A, Schikora A, Ordon F and Wehner G (2023) QTL for induced resistance against leaf rust in barley. *Front. Plant Sci.* 13:1069087. doi: 10.3389/fpls.2022.1069087

COPYRIGHT

© 2023 Matros, Schikora, Ordon and Wehner. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

QTL for induced resistance against leaf rust in barley

Andrea Matros¹, Adam Schikora², Frank Ordon^{1,2} and Gwendolin Wehner^{1*}

¹Julius Kühn Institute (JKI), Federal Research Centre for Cultivated Plants, Institute for Resistance Research and Stress Tolerance, Quedlinburg, Germany, ²Julius Kühn Institute (JKI), Federal Research Centre for Cultivated Plants, Institute for Epidemiology and Pathogen Diagnostics, Braunschweig, Germany

Leaf rust caused by Puccinia hordei is one of the major diseases of barley (Hordeum vulgare L.) leading to yield losses up to 60%. Even though, resistance genes Rph1 to Rph28 are known, most of these are already overcome. In this context, priming may promote enhanced resistance to P. hordei. Several bacterial communities such as the soil bacterium *Ensifer* (syn. *Sinorhizobium*) meliloti are reported to induce resistance by priming. During quorum sensing in populations of gram negative bacteria, they produce N-acyl homoserinelactones (AHL), which induce resistance in plants in a species- and genotypespecific manner. Therefore, the present study aims to detect genotypic differences in the response of barley to AHL, followed by the identification of genomic regions involved in priming efficiency of barley. A diverse set of 198 spring barley accessions was treated with a repaired E. meliloti natural mutant strain expR+ch producing a substantial amount of AHL and a transformed E. meliloti strain carrying the lactonase gene attM from Agrobacterium tumefaciens. For P. hordei resistance the diseased leaf area and the infection type were scored 12 dpi (days post-inoculation), and the corresponding relative infection and priming efficiency were calculated. Results revealed significant effects (p<0.001) of the bacterial treatment indicating a positive effect of priming on resistance to P. hordei. In a genome-wide association study (GWAS), based on the observed phenotypic differences and 493,846 filtered SNPs derived from the Illumina 9k iSelect chip, genotyping by sequencing (GBS), and exome capture data, 11 quantitative trait loci (QTL) were identified with a hot spot on the short arm of the barley chromosome 6H, associated to improved resistance to P. hordei after priming with E. meliloti expR+ch. Genes in these QTL regions represent promising candidates for future research on the mechanisms of plant-microbe interactions.

KEYWORDS

biotic stress resistance, Ensifer meliloti, AHL priming, Puccinia hordei, barley, leaf rust, GWAS, QTL

1 Introduction

Leaf rust caused by the biotrophic fungus Puccinia hordei is a serious disease of barley, leading to yield losses from 15-25% (Whelan et al., 1997) up to 62% (Park et al., 2015). Visual symptoms vary from small chlorotic lesions to large orangebrown pustules of about 0.5 mm in size, often surrounded by green leaf areas (Clifford, 1985). A major strategy in controlling leaf rust epidemics is breeding of resistant barley cultivars. While several resistance genes (Rph1 to Rph28, and Rph_{MBR1012}) have been identified in recent years (Singh et al., 2015; Fazlikhani et al., 2019; Chen et al., 2021), most of these are already overcome. Breakdown of resistance by mutations in effector genes of the pathogen frequently lead to the occurrence of new virulent races (Park, 2003), and thus a rapid decrease of the number of effective Rph genes (Kavanagh et al., 2017). Other effective management practices include the application of chemical fungicides, which can cause negative environmental (Ballabio et al., 2018) and public health effects (Perlin et al., 2017) if applied incorrectly. In this regard, decision-based management strategies may help to reduce their environmental impact considerably in the coming years (Lázaro et al., 2021). However, to promote more sustainable agricultural systems establishing alternative strategies to manage leaf rust epidemics is of prime importance, besides the ongoing genetic mapping of resistance genes.

Novel approaches for sustainable and economical crop production under ever-changing climate conditions include microbiome-facilitated crop management strategies (Liu et al., 2020; Trivedi et al., 2021). Beneficial plant-associated microbes include predominantly rhizobacteria and fungi as well as viruses, actinomycetes, cyanobacteria, archaea (Mauch-Mani et al., 2017; Mahapatra et al., 2020). The barley-microbiome was reported to largely vary between the different plant organs, developmental stages, origin, as well as between genotypes (Yang et al., 2017; Alegria Terrazas et al., 2020; Bziuk et al., 2021). Reported positive effects of such interactions in barley include reduced virulence of plant pathogenic fungi (Wehner et al., 2019; Karlsson et al., 2021), higher tolerance to abiotic stress (Caddell et al., 2019; Yang et al., 2020), as well as plant growth promotion and nutrient uptake under adverse conditions (Trivedi et al., 2020; Ribaudo et al., 2020).

The activation of induced defense mechanisms by various stimuli, such as from pathogens, beneficial microbes, arthropods, as well as chemicals and abiotic cues, is generally regarded as priming (Mauch-Mani et al., 2017; Desmedt et al., 2021). Upon priming, plants respond stronger and faster to a biotic or abiotic stress factor, resulting in resistance and securing yield. The use of priming for enhancing resistance has a long tradition, and the phenomenon called "sensitization" or "preformed defense" was investigated and used since the 1930s (Chester, 1933; Starkey, 1958). In this respect, it has to be noted, that microbial priming-induced plant responses vary between species and depend on the composition of the soil-

microbiome as well as on the genotype and vice versa (Jack et al., 2019; Morella et al., 2020; Alegria Terrazas et al., 2020; Si et al., 2021).

Various organic compounds, for instance salicylic acid (SA), benzothiadiazole, β -aminobutyric acid, and azelaic acid are known to induce priming (Conrath et al., 2002; Jung et al., 2009; Mauch-Mani et al., 2017). In addition, many recent reports are highlighting bacterial quorum sensing (QS) molecules as priming-inducers in plants. Related to plant-pathogen interactions, the best-studied QS molecules are *N*-acyl homoserine lactones (AHL), which are produced by numerous gram negative bacteria to communicate and to monitor the density of populations (Kaplan and Greenberg, 1985; Fuqua et al., 1994). Plants generally respond to short-chained AHL with modifications in growth, while long-chained AHL induce AHL-priming for enhanced resistance against pathogen infection (Shrestha et al., 2020).

In tomato plants, root colonization with AHL-producing *Serratia liquefaciens* MG1 and *Pseudomonas putida* IsoF lead to increased systemic resistance against the fungal leaf pathogen *Alternaria alternate*, likely in an SA- and ethylene-dependent manner (Schuhegger et al., 2006). The effect has been related to the four AHLs N-(3-oxo-hexanoyl)-L-homoserine lactone (3-oxo-C6-HSL), 3-oxo-C8-HSL, 3-oxo-C10-HSL, and 3-oxo-C12-HSL produced by *P. putida* IsoF (Gantner et al., 2006), and to the two AHLs butanoyl-homoserine lactone (C4-HSL) and 3-oxo-C6-HSL produced by *S. liquefaciens* MG1. Plants inoculated with the AHL-negative mutant strain *S. liquefaciens* MG44 exhibited a susceptible phenotype (Schuhegger et al., 2006).

Another rhizobacterium with known AHL-priming capacity is Ensifer meliloti, a naturally occurring gram negative soil bacterium (Teplitski et al., 2003; Fagorzi et al., 2020). By studying AHL-priming with E. meliloti as a model in the laboratory, different authors tested resistance against various pathogens, for instance Pseudomonas syringae in Arabidopsis thaliana (Zarkani et al., 2013; Shrestha et al., 2020), Blumeria graminis (Shrestha et al., 2019), Puccinia hordei in barley (Hernández-Reyes et al., 2014; Wehner et al., 2019; Shrestha and Schikora, 2020) and the pest Pratylenchus penetrans in soybean (Adss et al., 2021). Notably, barley genotypes showed different sensitivity to AHL-priming when treated with E. meliloti, indicated by diverse resistance responses of seven barley accessions to P. hordei infection (Wehner et al., 2019) and of eight barley accessions to B. graminis infection (Shrestha et al., 2019). These results suggest that AHL-priming is highly influenced by the genetic background of the respective accession, and strongly propose the utilization of priming-efficiency as a future breeding goal. However, the genetic background underlying the variability of AHL-priming-induced resistance in plants has not yet been investigated.

In order to map genetic loci underlying differences in response to AHL-priming, genome wide association studies (GWAS) are a powerful tool since high throughput marker technologies like the 9k iSelect chip (Comadran et al., 2012), genotyping by sequencing (Poland and Rife, 2012) or exome capture (EC) sequencing (Mascher et al., 2013) are available in barley. While family-based linkage analysis searches for associations within populations developed from bi-parental crosses, association mapping utilizes historic patterns of recombination that have occurred within a sample of individuals. Association mapping is based on the principle that over multiple generations of recombination, correlations only with markers tightly linked to the trait of interest will be present. As a predominantly inbreeding species, cultivated barley is an attractive target for association mapping as its genome contains extensive blocks of chromatin in linkage disequilibrium (Rostoks et al., 2006), providing a well-defined haplotype structure from which markertrait associations can be identified (Cockram et al., 2008).

In this study, we aim to investigate the genetic background of resistance priming in barley. Thus, the objective is identifying barley varieties with an enhanced potential to respond to *E. meliloti* treatment with an increased resistance to a *P. hordei* infection by screening a set of barley genotypes with a diverse genetic background. A set of 198 spring barley accessions was challenged with either the 3-oxo-C14-HSL accumulating *E. meliloti* strain *expR+ch* or with the control *E. meliloti* strain *attM*, not being able to accumulate AHL (Zarkani et al., 2013). Responses after *P. hordei* infection were monitored and relative infection values were used as a target for GWAS. Identified primable accessions and QTL are discussed regarding the molecular mechanisms of AHL-priming and breeding applications.

2 Material and methods

2.1 Plant material, experimental design, and treatment

In order to determine the effects of priming in relation to an infection with *P. hordei* causing leaf rust, we made use of the

potentially most diverse GWAS panel for barley comprising 200 spring barley accessions and landraces of worldwide origin from the spring barley IPK-SB224 panel, including the "Genobar" panel, (Haseneyer et al., 2010; Pasam et al., 2012). Our study panel consisted of 111 two-rowed and 83 six-rowed accessions from the Genobar panel and additional six barley lines, namely Barke, Morex, Steptoe, Golden Promise, Großklappige, and Roland, making it a total of 200 spring barley accessions of which 198 were genotyped (Table S1 and Table S2, Figure S1).

This panel was investigated in greenhouse pot experiments for three growing seasons (2017, 2018 and 2019) in a split plot design with three pots per accession. The experimental workflow is illustrated in Figure 1. After two days of germination on wet filter paper, seedlings were transferred to potting mix (Fruhstorfer Typ T) with three plants per pot (7x7x6 cm filled with 0.2 L of substrate) and cultivated in the greenhouse at night/ day temperatures of 18°C/22°C with additional lighting for 10 h. For watering of pots to full water holding capacity, tap water was used, and the substrate was kept at similar moisture for all accessions and treatments during the course of the experiment. Two days after planting (dap) all accessions were treated either with the repaired *E. meliloti* natural mutant strain expR+ch(Rm2011 expR+ch) or with the transformed E. meliloti strain attM (pBBR2-attM). ExpR+ch is able to accumulate a large amount of the AHL 3-oxo-C14-HSL whereas attM was used as a control, carrying a lactonase gene from Agrobacterium tumefaciens and is therefore not able to accumulate AHL. Both bacteria strains were grown in tryptone yeast (TY) extract medium until the OD_{600nm} of 0.6 to 0.8. Bacterial cultures were centrifuged at $2,500 \times g$ for 10 min and suspended in 10 mM MgCl₂ (Zarkani et al., 2013). The bacteria solutions were applied three times in total. In fact, the substrate was treated with



3.5 ml of bacteria solution (OD_{600nm} of 0.1, corresponding to 10^8 CFU/ml) each, at two, eight and 14 dap (Wehner et al., 2019) using a multi-dispenser pipette equipped with a 50 ml tip vessel. Inoculation of plants with the *P. hordei* virulent strain I-80 was performed at the three-leaf stage at 16 dap. Therefore, trays containing 24 pots each were placed in sealable containers. The leaf rust spores (25 mg) were mixed with white clay (1:3) and applied equally to the plants in the containers, using a manual powder spray bottle. Immediately after the inoculation, the containers were covered with foil to keep a moist atmosphere and the plants were stored for 24 hours without light. After removing the trays from the containers, plants were further cultivated in the greenhouse as stated above.

2.2 Phenotyping for relative infection and priming efficiency

The scoring was carried out 12 dpi (Figure 1) by estimating the diseased leaf area in percentage [%] (Moll et al., 2000) and the infection type (Levine and Cherewick, 1952). Thereof, the relative infection as well as priming efficiency were calculated according to Wehner et al. (2019).

Relative infection = $0.2 \times ln (P. hordei \%) + ln (P. hordei scores)$

Priming efficiency

= relative infection (attM) – relative infection (expR + ch)

Figures showing relative infection results were generated with the data analysis software JMP version 14 (https://www.jmp.com/en_gb/software/data-analysis-software.html). The barley accessions were clustered in primable and non-primable genotypes, by the difference of the relative infection between the priming treatments (= priming efficiency), applying a threshold for priming efficiency of >0.2 for primable ones and a significance level of p<0.05. Best linear unbiased estimates (BLUEs) of relative infection and priming efficiency for the set of accessions under investigation are provided in Table S2.

The descriptive statistics of the phenotypic data were carried out in the software "RStudio" version 3.4.2 (https://www.rstudio.com/ products/rstudio/) using the R-packages "pastecs" and "agricolae". The minimum (Min), the maximum (Max), mean, standard deviation (SD), coefficient of variation (CV, standard deviation divided by mean), least significant difference (LSD), and repeatability for relative infection with *P. hordei* for the 198 accessions and for three years experiments were calculated for *attM* as well as for *expR+ch* (Table 1). Repeatability of the experiments in the greenhouse was calculated based on the variance components of the accessions and the variance associated with the accession by year interaction implemented in the R-package "Ime4". Based on these data, an analysis of variances (ANOVA) was carried out over the total number of accessions in order to check for any significant effects for the accessions, the priming treatment and their interaction (Table 2). Thus, a linear mixed effects (lme) model by means of the R-package "nlme" was applied, using priming, accessions, and the interaction of both as fixed effects and the plant per pot, replication, and year as random effects.

2.3 Genotyping and GWAS

For 198 accessions single nucleotide polymorphism marker data was gained (Table S1) derived from the Illumina 9k iSelect chip (SGS Trait genetics) following Comadran et al. (2012). In addition, for 175 accessions genotyping by sequencing (GBS) and for 155 accessions Exome Capture analyses (Mascher et al., 2013) were created. Thus, 5,764 iSelect markers, 17,653 GBS markers and 470,429 exome capture (EC) markers were used after filtering for<12.5% heterozygous SNPs, >5% minor allele frequency and<10% missing values. Genetic positions for all markers were extracted from the barley reference genome based on MorexV3 (Mascher et al., 2021) by means of the Barleymap software module "Locate by position" (https://floresta.eead.csic. es/barleymap/locate/), which was published by the international barley sequencing consortium (Cantalapiedra et al., 2015).

SNP markers from the iSelect Chip were used for calculation of the population structure in STRUCTURE (Pritchard et al., 2000). In this program, ten independent runs of Monte Carlo Markov chain with burn-in period of 500,000 were calculated to obtain the optimum k-value and the corresponding q-matrix. Two subpopulations clustered by row-type (Figure S1) were identified according to the method of Evanno et al. (2005). Furthermore, linkage disequilibrium (LD) was calculated according to Hill and Weir (1988) using the GenABEL package in R.

A genome-wide association study (GWAS) was carried out in order to detect genetic loci that underlie the difference in the response

TABLE 1 Descriptive statistics of relative infection with *P. hordei* for the 198 accessions treated with either the *attM* or *expR+ch* bacterial strain.

Priming	attM	expR+ch
Min	1.110	1.060
Max	1.690	1.440
Mean	1.490	1.310
SD	0.081	0.099
CV	0.054	0.075
LSD	0.004	0.010
r ²	0.636	0.314

Minimum (Min), maximum (Max), mean, standard deviation (SD), coefficient of variation (CV, standard deviation divided by mean), least significant difference (LSD), and repeatability (r^2) are listed.

TABLE 2 Analysis of variances (ANOVA) of relative infection with *P. hordei* for the 198 accessions.

Effect	F-value	P-value
Priming (αi)	306.06	< 0.001
Accession (βj)	8.89	< 0.001
(αβ)ij	3.33	< 0.001

to the bacterial priming. The total data set of 493,846 SNP markers, phenotypic data by BLUEs, and the q-matrix was used for the GWAS. The GAPIT tool in R was used in which a compressed mixed linear model (cMLM) was applied using q-matrix and kinship (calculated by Van Raden within the package) as random factors (Lipka et al., 2012) after a model selection step including cMLM with kinship as cofactor, a generalized linear model (GLM) and cMLM with qmatrix and kinship as cofactors (Figure S2). In this comparison, closest alignment to the line of expected probabilities (black) was shown for cMLM with q-matrix and kinship as cofactors (green, subjacent the black line in the graph), which was therefore selected for GWAS. Furthermore, SNP permutation (1,000 permutations) was applied to assess the empirical significance of SNPs and compression level of 1.9 was used as suggested in the GAPIT package. Significant marker trait associations (MTA) were identified using a threshold of the false discovery rate (FDR) at p<0.05 which in this case represents a likelihood of odds (LOD) score of 4. Significant MTAs were further clustered in corresponding putative QTL by the LD of 21,000 bp. LD was calculated using the R package "GenABEL". Physical positions together with marker types and the phenotypic variances explained (PVE) are visualized using MapChart program (Voorrips, 2002).

2.4 Annotations regarding the Morex reference genome

Annotation of genes at the identified marker positions was obtained from information provided in the Barleymap software module "Locate by position" containing annotations from the Morex genome (Mascher et al., 2017) and updated to the MorexV3 genome released in 2021 (Mascher et al., 2021). Annotation of several iSelect markers was obtained from sequence alignments using the BLAST tool of the UniProt reference database (https://www.uniprot.org/blast/). Gene transcript expression data were obtained from the Expression Atlas database for barley (https://www.ebi.ac.uk/gxa/home#).

3 Results

3.1 The priming effect varies between the various accessions

An established priming system using *E. meliloti* expR+ch as primary trigger was applied (Wehner et al., 2019). The low

standard deviation and the associated high repeatability shows that the priming method used, was reproducible.

The priming effect of the 198 accessions analyzed is represented in the average relative infection with P. hordei (Figure 2). Results clearly show that the expR+ch bacterial strain treated group of plants was less infected in comparison to the control group treated with the attM bacterial strain (Figure 2 and Table 1). However, a large variation of the accessions regarding the relative infection was observed and allowed for the differentiation in primable and non-primable accessions based on priming efficiency. Significant treatment effects with p<0.05 were detected for 87 accessions meaning that 43.5% of the accessions in our study can be defined as primable with AHL (Figure 2, Table S2). Concordantly, the large diversity for priming efficiency in the panel is also shown by the high coefficients of variation (CV) for both treatments (Table 1). Furthermore, the effects can be assumed stable over the three year's experiments, with a calculated repeatability of 63.6% for attm control treatments, while repeatability was low with 31.4% for the expR+ch primed plants. In addition, ANOVA results revealed significant effects (p<0.001) of the priming treatment indicating a positive effect of priming on resistance to P. hordei infection (Table 2). Also, significant (p<0.001) effects between the accessions and for accessions times priming interactions were observed. Thus, from the results a significant priming effect can be concluded for AHL-induced P. hordei resistance, which is accession-specific.

The ten accessions with the highest priming efficiency are all of non-European origin, and comprise eight six-rowed and two two-rowed accessions (Table S2): BCC129 (MAR, 6-row), HOR11403 (IND, 6-row), BCC421 (CHN. 6-row), BCC551 (IND, 6-row), BCC538 (IND, 6-row), BCC93 (IRQ, 2-row), Morex (USA, 6-row), BCC868 (MEX, 6-row), BCC903 (CAN, 2-row), and BCC881 (CAN, 6-row). Among the ten accessions with the lowest priming efficiency are two of European origin, and four accessions are of six-rowed and six of two-rowed ear



Relative infection with *P. hordei* (BLUES) in the 198 accessions treated with either the *attM* (red) or *expR+ch* (blue) bacterial strain and ordered by priming efficiency. Significant treatment effects with p<0.05 were detected for 87 accessions.

type: HOR8160 (TUR, 2-row), BCC192 (SYR, 2-row), BCC899 (CHL, 2-row), HOR11373 (ISR, 2-row), BCC1385 (POL, 2-row), BCC927 (PER, 6-row), BCC445 (CHN, 6-row), BCC844 (COL, 6-row), BCC1433 (DEU, 2-row), and BCC807 (URY, 6-row). However, apart from a trend to non-European origin and six-rowed ear type being correlated with higher priming efficiency no clear relation to genetically determined phenotypic traits could be made.

3.2 GWAS identified markers for relative infection

For the three-year data on relative infection of primed (expR +ch) and control plants (attM), 129 significant marker-trait associations (MTAs) were identified by GWAS (Table S3, Figure S3). Notably, primarily exome capture (EC) markers were associated, highlighting the predominance of this marker type in our dataset. Identified MTAs were clustered into QTL using the critical LD decay value of 21,000 bp and MTA not in LD were considered as an independent QTL. Overall, 70 QTL were identified (Table S3). The identified QTL were unevenly distributed across the barley chromosomes with one QTL detected on chromosome 1H, five on chromosome 2H, three on chromosome 3H, one on chromosome 4H, seven on chromosome 5H. A hot spot of 42 QTL was detected on chromosome 6H, and 11 QTL were detected on chromosome 7H (Figure 3). Detected QTL showed a positive effect on relative infection with significant LOD values ranging between 4.01 and 6.21 (Table S3). The lowest LOD value was found for MTA 33 in

QTL 21 on chromosome 6H and the highest LOD value was found for MTA 102, 103 and 104 in QTL 65 on chromosome 7H. The respective QTL explain up to 13.5% of the phenotypic variance (PVE) as specified in Figure 3 and listed in Table S3, indicating the potential to increase resistance to *P. hordei* in spring barley.

We have not detected QTL overlapping for relative infection of primed (expR+ch) and control plants (attM) in our study. Of all identified significant MTA, 112 MTA clustered in 59 QTL, which were associated with the relative infection under attmtreatment. Therein, the percentage of PVE by each MTA varied from 9.0 to 13.5% (Figure 3). In addition, 17 MTA clustered in 11 QTL, which are associated with the relative infection under expR+ch priming (Table 3). These 11 QTL were distributed over the barley chromosomes 2H (one QTL), 3H (one QTL), 5H (three QTL), 6H (four QTL), and 7H (two QTL). Variation of PVE by each MTA varied between 8.3 to 10.1% (Figure 3).

Out of the 70 detected QTL, annotation of marker positions within 23 QTL could not be obtained. This relates to 34 of the 129 identified MTA (Table S3). From the annotated genes, 13 are related to proteins with unknown functions (unknown protein or undescribed protein). Promising candidates associated with increased resistance to *P. hordei* treated with the *attm* control strain included i) several transporters: Magnesium transporter NIPA2 (HORVU1Hr1G077020) in QTL one, Mechanosensitive ion channel protein 2 (HORVU5Hr1G095750) in QTL 17, LETM1 and EF-hand domain-containing protein 1 (HORVU6Hr1G055750) in QTL 28, ii) transcription factors: WRKY DNA-binding protein 2 (HORVU5Hr1G058640) in QTL 58,



TABLE 3 In a genome-wide association study (GWAS), based on the observed phenotypic differences and 493,846 filtered SNPs derived from the Illumina 9k iSelect (iSelect) chip, genotyping by sequencing (GBS), and exome capture data (EC), 11 quantitative trait loci (QTL) associated to improved resistance to *P. hordei* after priming with *E. meliloti expR+ch*, were identified with a peak on the short arm of the barley chromosome (Chr.) 6H.

QTL	SNP	Marker type	Chr.	Position in Mb	LOD	Annotation (Barleymap, ^a UniProt BlastX)	Gene Name	Differential Expression (up-regu- lated)	Location	GO	InterPro
5	chr2H_44543906	GBS	2H	44,543906	4,13	WRKY transcription factor 4	HORVU2Hr1G017720	Infection, wound	Nucleus	DNA-binding, transcription factor activity	WRKY domain
7	chr3H_15470342EC	EC	3H	15,470342	4,84	Xylanase inhibitor	HORVU3Hr1G006440	Infection, wound		Aspartic-type endopeptidase activity, proteolysis	Xylanase inhibitor, Aspartic peptidase A1
	chr3H_15470343EC	EC	3H	15,470343	4,84	Xylanase inhibitor	HORVU3Hr1G006440	Infection, wound		Aspartic-type endopeptidase activity, proteolysis	Xylanase inhibitor, Aspartic peptidase A1
11	chr5H_5584131EC	EC	5H	5,584131	4,30	Laccase	HORVU5Hr1G001990			Lignin catabolic process	Reverse transcriptase-like
	chr5H_5584216EC	EC	5H	5,584216	4,35	Laccase	HORVU5Hr1G001990			Lignin catabolic process	Reverse transcriptase-like
13	SCRI_RS_205235	iSelect	5H	491,233648	4,25	ATP-dependent zinc metalloprotease FtsH 2 ^a	HORVU5Hr1G063340	Infection, wound	Integral membrane	ATPase activity, ATP binding	ATPase, nucleoside triphosphate hydrolase
14	BOPA1_8215-496	iSelect	5H	498,937856	4,73	PDZ domain- containing protein ^a	HORVU5Hr1G065220		Nucleus, cytoplasm	Proteasome regulatory particle assembly, protein binding	26S Proteasome non- ATPase regulatory subunit 9
18	chr6H_16152203EC	EC	6H	16,152203	4,23	Pentatricopeptide repeat-containing protein	HORVU6Hr1G008870	Drought		Nucleic acid binding, protein binding	Zinc finger C2H2 superfamily
21	SCRI_RS_182367	iSelect	6H	44,243728	4,01	Esterase/lipase/ thioesterase family protein	HORVU6Hr1G018050		ER membrane	Transferase activity, transferring acyl groups other than amino-acyl groups	Alpha/Beta hydrolase, Diacylglycerol acyltransferase
	chr6H_44247682	GBS	6H	44,247682	4,44	Esterase/lipase/ thioesterase family protein	HORVU6Hr1G018050		ER membrane	Transferase activity, transferring acyl groups other than amino-acyl groups	Alpha/Beta hydrolase, Diacylglycerol acyltransferase
	chr6H_44247749	GBS	6H	44,247749	4,44	Esterase/lipase/ thioesterase family protein	HORVU6Hr1G018050		ER membrane	Transferase activity, transferring acyl groups	Alpha/Beta hydrolase, Diacylglycerol acyltransferase
											(Continued)

TABLE 3	Continued
---------	-----------

QTL	SNP	Marker type	Chr.	Position in Mb	LOD	Annotation (Barleymap, ^a UniProt BlastX)	Gene Name	Differential Expression (up-regu- lated)	Location	GO	InterPro
										other than amino-acyl groups	
	chr6H_44247752	GBS	6H	44,247752	4,44	Esterase/lipase/ thioesterase family protein	HORVU6Hr1G018050		ER membrane	Transferase activity, transferring acyl groups other than amino-acyl groups	Alpha/Beta hydrolase, Diacylglycerol acyltransferase
	chr6H_44247776	GBS	6H	44,247776	4,44	Esterase/lipase/ thioesterase family protein	HORVU6Hr1G018050		ER membrane	Transferase activity, transferring acyl groups other than amino-acyl groups	Alpha/Beta hydrolase, Diacylglycerol acyltransferase
22	SCRI_RS_121633	iSelect	6Н	47,62853	4,33	Protein FLUORESCENT IN BLUE LIGHT	HORVU6Hr1G018610		Chloroplast membrane	Chlorophyll/tetrapyrrole biosynthetic process, protein binding	Protein FLUORESCENT IN BLUE LIGHT, Tetratricopeptide repeat- containing
23	chr6H_50004267	GBS	6H	50,004267	4,11						
60	chr7H_523570079EC	EC	7H	523,570079	4,25	Undescribed protein	HORVU7Hr1G086760				
61	chr7H_535632749EC	EC	7H	535,632749	4,17						
Positio	Positions of the detected SNP marker were mapped against the barley reference genome and identified QTL and detected SNP marker in these regions are listed together with the annotation and available gene transcript expression information.										

and iii) several proteins involved in nucleic acid and protein processing (Table S3). Unfortunately, the three markers with the highest LOD value of 6.21 (MTA 102, 103 and 104) in QTL 65 on chromosome 7H could not be annotated, yet.

The most interesting QTL associated with increased resistance to P. hordei primed with E. meliloti expR+ch was identified on the short arm of barley chromosome 6H (Figure 3, Table 3). The five marker clustering into QTL 21 are located in the gene HORVU6Hr1G018050 being annotated as esterase/ lipase/thioesterase family protein. In addition, the two markers clustering into QTL 7 on chromosome 3H, are located in the gene HORVU3Hr1G006440 which is a xylanase inhibitor. This gene has been reported to show increased transcript accumulation under infection and wounding (Table 3). The former is also the QTL containing the markers with the highest PVE among the QTL associated with increased resistance to P. hordei after AHL-priming, with 10.1% (Figure 3). Similarly, the two markers in QTL 5 and QTL 13 are located in genes, which show increased transcript expression under infection and wounding: namely HORVU2Hr1G017720, the WRKY transcription factor 4, and HORVU5Hr1G063340, the ATP-dependent zinc metalloprotease FtsH 2, respectively. Also, HORVU5Hr1G001990, a laccase involved in lignin catabolism, may be involved in increased resistance to P. hordei following AHL-priming. The two markers clustering into QTL 11 on chromosome 5H were located in this gene (Table 3).

4 Discussion

Within this three-year trial, a positive effect of priming with the gram negative soil bacterium E. meliloti strain expR+ch in relation to resistance to P. hordei was demonstrated for spring barley. The effect on relative infection and thus priming efficiency varied among the set of 198 accessions, of which 87 accessions showed significant (p<0.05) treatment effects and therefore have been considered highly primable (Figure 2). Nevertheless, in practical applications the priming effect might be influenced by environmental conditions, such as soil type and the presence of other microorganisms. To our knowledge, the presented study is the first investigation on the effects of the priming inducer E. meliloti strain *expR+ch* on cereal resistance against fungal pathogens including a larger set of genetically differentiating accessions. While several recent reports are suggesting that the microbiome varies between the different plant organs, developmental stages, origin, as well as between genotypes in wheat (Gdanetz and Trail, 2017; Kuźniar et al., 2020; Žiarovská et al., 2020) and barley (Yang et al., 2017; Alegria Terrazas et al., 2020; Bziuk et al., 2021), studies on genetic variation of priming efficiency are rare. In a previous

experiment, we observed significant differences between seven spring barley accessions, indicating genotypic differences in the response to priming by E. meliloti strain expR+ch towards P. hordei resistance (Wehner et al., 2019). Similarly, using a set of eight genetically diverse spring barley accessions, Shrestha et al. (2019) have demonstrated differences in their ability to be primed by E. meliloti strain expR+ch with respect to resistance against B. graminis f. sp. hordei. Other root associated microorganisms, for which genotype dependent responses have been shown, include arbuscular mycorrhiza fungi (AMF), reviewed in Parke and Kaeppler (2000) and Sawers et al. (2010). Varying abilities to respond to AMF were reported for wheat (Hetrick et al., 1993), maize (Kaeppler et al., 2000; An et al., 2010), and onion (Powell et al., 1982; Tawaraya et al., 2001; Galván et al., 2011). Consistently, investigation of a diverse set of 94 bread wheat genotypes (Triticum aestivum) regarding root colonization by AMF identified significant genotypic differences (p<0.001) for mycorrhizal colonization and detected 30 significant genetic markers associated with root colonization (Lehnert et al., 2017). Within the same set of genotypes, drought stress tolerance was significantly increased in the presence of AMF colonization compared to drought stress tolerance in the absence of AMF colonization (Lehnert et al., 2018).

In our results, we observed a trend to non-European origin and six-rowed ear type being correlated with higher priming efficiency (Table S2). However, no clear relation to genetically determined phenotypic traits could be made. The model cultivar Morex was among the ten accessions with highest priming efficiency. This is concordant with our previous finding, where Morex was identified as a well primable accession (Wehner et al., 2019). In this study, the accessions BCC768 and HOR7985 were also assigned as wellprimable, a result which was confirmed in the recent study. While the resistance gene Rph8 was mapped in the Morex reference genome on chromosome 5H at 12.7 Mb (Mascher et al., 2017), we observed high values for relative infection under attM control treatment in this accession (Table S2). This can be explained by breakdown of resistance gene Rph8 by the P. hordei I-80 isolate (Niks et al., 2000), which was used in the present study. Notably, some QTL identified during the control treatment were in the genomic region of known resistance genes, for example Rph19 (Park and Karakousis, 2002) on the barley chromosome 7H (see Figure 3). Together, results suggest a high genetic determination of priming responses and thus highlight the importance of screening large diversity sets for detecting quantitative trait loci or candidate genes involved in priming.

In our study, of the 129 MTA identified, 17 MTA, which cluster in 11 QTL, were associated to improved resistance to *P. hordei* after priming with the oxo-C14-HSL-producing *E. meliloti* expR+ch, with a hot spot on the short arm of barley chromosome 6H (Figure 3, Table 3, and Table S3). Except two markers, all

markers were assigned to positions in the genome within annotated barley coding genes. On chromosome 6H, the marker chr6H_16152203EC (QTL 18, LOD 4.23) is localized in the gene HORVU6Hr1G008870 encoding for a pentatricopeptide repeatcontaining protein. This gene product belongs to the class of zinc finger proteins involved in DNA binding, which may have regulatory function, and which was shown to be induced by drought stress in barley in a previous study (Guo et al., 2009). Another significant marker on chromosome 6H (SCRI_RS_121633, QTL 22, LOD 4.33) is localized in the gene HORVU6Hr1G018610 encoding for the protein FLUORESCENT IN BLUE LIGHT. In A. thaliana, this gene product was reported as a negative regulator of chlorophyll biosynthesis acting in a light dependent manner (Hou et al., 2019). How both gene products mentioned above are involved in the process of induced systemic resistance by AHL priming remains speculation. However, limitation of photosynthetic activity has been assumed one possible strategy of plants to increase pathogen resistance. For instance, the accumulation of oxylipin in distal tissues during AHL-priming in Arabidopsis promoted stomatal closure. The closed stomata enhanced plant resistance to bacterial pathogens (Schenk and Schikora, 2015). In this line, a very promising QTL (21) on chromosome 6H comprises five significant markers from two different marker types (SCRI_RS_182367, chr6H_44247682, chr6H_44247749, chr6H_44247752, chr6H_44247776), all localized in the same gene HORVU6Hr1G018050, which encodes for a esterase/lipase/thioesterase family protein. In barley, this protein was not characterized functionally yet, but a BLAST search with the protein sequence in the NCBI database (https://blast.ncbi.nlm.nih.gov/Blast.cgi) revealed highest similarity to the chloroplastic phytyl ester synthase 1 (PES1) from A. thaliana (At1g54570). It exhibits phytyl ester synthesis and diacylglycerol acyltransferase activities with broad substrate specificities, and can employ acyl-CoAs, acyl carrier proteins, and galactolipids as acyl donors. PES1 was shown to be involved in fatty acid phytyl ester synthesis in chloroplasts, a process required for the maintenance of the photosynthetic membrane integrity during abiotic stress and senescence (Lippold et al., 2012). Accordingly, deposition of free phytol and free fatty acids in the form of phytyl esters in chloroplasts was reported for etiolated barley seedlings after 6 to 8 h irradiation (Liljenberg, 1977), and in primary leaves of barley during methyl jasmonate induced leaf senescence (Springer et al., 2015). Notably, HORVU6Hr1G018050 was also found in a QTL region on chromosome 6H in the interval between 37-76 Mb associated with resistance against net blotch causing Pyrenophora teres f. teres in a barley diversity set (Novakazi et al., 2019). On chromosome 5H, the marker BOPA1_8215-496 (QTL 14, LOD 4,73) is localized in the gene HORVU5Hr1G065220 encoding for a PDZ domain-containing protein, which is likely involved in the regulation of protein translation via the proteasome with unknown

contribution to pathogen resistance. Interestingly, the marker SCRI_RS_205235 (QTL 13, LOD 4,25) is localized in the ATPdependent zinc metalloprotease FtsH coding gene 2HORVU5Hr1G063340. FtsH is the major thylakoid membrane protease required for photosynthetic pathways in plants (Kato and Sakamoto, 2018). This gene was recently shown to be relatively highly expressed in all grain organs suggesting its crucial role in the accumulation of the micronutrient metals (i.e., Cu, Fe, Mn, and Zn) in barley grains (Thabet et al., 2022). It may be speculated, that FtsH represents another component of the mechanisms activated to limit photosynthetic activity to increase pathogen resistance. However, despite the upregulation observed during infection and wounding (Table 3), the involvement of FtsH in improved resistance against P. hordei after AHL-priming needs further molecular validation. Two other markers on chromosome 5H (chr5H_5584131EC, chr5H_5584216EC) cluster in QTL 11 with LOD 4.3 and 4.35, respectively. Both are located in the same gene HORVU5Hr1G001990, which encodes for a laccase. Laccases constitute a multi-gene family of multi-copper glycoproteins and have diverse and overlapping physiological functions in plants, including involvement in redox metabolism, responses to wound healing, defense against pathogens or insects, synthesis of lignin and suberin, and cross-linking of cell wall components; cf. Janusz et al. (2020) and Tomková et al. (2012). While several laccases have been annotated in the barley genome on all chromosomes recently (https://plants.ensembl.org/Hordeum_vulgare/Search/Results? page=3;q=laccase;species=Hordeum_vulgare;collection=all;site= ensembl), only HvLac1 on chromosome 4H has been characterized in detail so far with unclear molecular function (Tomková et al., 2012). However, we suggest that the laccase, identified in our study is likely involved in cell wall fortification representing one of the processes leading to increased resistance to P. hordei by AHL priming. Consistently (Shrestha et al., 2019) showed that the principal mechanism of AHL-induced priming in barley seems similar to the mechanism in Arabidopsis, i.e., it is associated with the activation of MAPKs, enhanced expression of various defenserelated genes and of genes involved in the remodeling of cell wall structure. Additionally, laccases were also identified among a group of genes that were associated with Fusarium graminearum resistance in barley (Huang et al., 2016). In the same line, the two markers identified on chromosome 3H (chr3H_15470342EC, chr3H_15470343EC, QTL7, LOD 4.84), which are both localized in the gene (HORVU3Hr1G006440) annotated as a xylanase inhibitor, represent interesting candidates with potential function in the fortification and remodeling of cell wall structure. This gene was also found to be upregulated after infection and wounding in barley (Table 3). Most cereals contain arabinoxylan as a structural component in their cell walls (Henry, 1985). As xylanases have a general role in the depolymerization of plant and fungal cell walls (Hrmova et al., 1997), the xylanase inhibitor identified in our study

could help to reduce or overcome the effect of plant and fungal related xylanases during infection with P. hordei and thus promote resistance. A marker with high potential in translation regulatory processes was identified on chromosome H2 (chr2H_44543906, LOD 4.13) localized in the gene HORVU2Hr1G017720, which is annotated as WRKY transcription factor 4. The WRKY transcription factor gene family is one of the largest, and it is known to be involved in a wide range of plant developmental and physiological processes. Although, more than 60 unique barley genes have been annotated containing the WRKY domain, WRKY proteins of barley are not yet fully annotated and most of them are not functionally characterized (Liu et al. (2014), http://www. barleygfdb.com/second/family.php?xu=77). The data of Dey et al. (2014) suggest that bacteria-induced systemic immunity in barley is associated with the local and/or systemic induction of transcript accumulation of Ethylene Responsive Factor (ERF)-like transcription factors (HvERF-like, HvERF44411) as well as HvWRKY22 and HvWRKY38/1. Similarly, Gao et al. (2018) proposed Nonexpressor of Pathogenesis-Related Genes1 (NPR1) homologs and WRKY transcription factors as the master regulators of systemic acquired resistance in wheat and barley. Furthermore, they showed that transient expression of HvWRKY6, HvWRKY40, and HvWRKY70, in wheat leaves by Agrobacteriummediated infiltration enhanced the resistance to Puccinia triticina. In another study, HvWRKY10, HvWRKY19, and HvWRKY28 were shown to positively regulate the response of barley to B. graminis infection (Meng and Wise, 2012). On the other hand, the wound and pathogen-inducible HvWRKY1 and HvWRKY2 are known as negative defense regulators repressing the activity of the powdery mildew-induced promoter of HvGER4c, a germin-like defense-related protein (Liu et al., 2014). However, WRKY transcription factor 4 (HORVU2Hr1G017720) was not functionally characterized yet, despite the induced expression of HORVU2Hr1G017720 after fungal infection and wounding (Table 3, http://www.barleygfdb.com/second/PRJNA728113.php? xu=77). It also needs to be considered that WRKY functions may be genotype specific, such as shown for WRKY transcription factors in barley cultivars infected with Fusarium culmorum (Uluhan et al., 2019).

In conclusion, our results provided clear evidence for the improved resistance of barley against *P. hordei* infection by AHL-mediated priming. Priming responses were genotype specific and most significant genetic markers for improved resistance to *P. hordei* after priming identified in our study were related to limiting photosynthetic activity, cell wall fortification, and regulation of transcription and translation. Our findings are important, as they open doors to study the mechanisms and the interactions between the plant genetic background and AHL-priming as well as supporting novel breeding approaches for priming efficient accessions and thus sustainable crop protection after validation.

Data availability statement

The original contributions presented in the study are included in the article/Supplementary Material. Further inquiries can be directed to the corresponding author.

Author contributions

The initial research idea was developed by FO and GW. The study conception and experimental design was established by AS and GW. Material preparation and data collection was performed by GW, and data analysis was performed by GW and AM. The first draft of the manuscript was written by GW, amended by AM, and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Funding

The authors wish to acknowledge financial support by the German Federal Ministry of Education and Research (Bundesministerium für Bildung und Forschung, BMBF) for the project PrimedPlant (031B0196C and 031B0196B).

Acknowledgments

We would like to thank Christine Hoppe, Nicole Refert, Ilona Renneberg, and the greenhouse team of the Julius Kühn Institute (JKI) Federal Research Institute for Cultivated Plants for excellent technical support.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: https://www.frontiersin.org/articles/10.3389/ fpls.2022.1069087/full#supplementary-material

SUPPLEMENTARY FIGURE 1

Composition of the test panel of 200 spring barley accessions. Our study panel consisted of 111 two-rowed and 83 six-rowed accessions from the Genobar panel (Haseneyer et al., 2010; Pasam et al., 2012) comprising

References

Adss, S., Liu, B., Beerhues, L., Hahn, V., Heuer, H., and Elhady, A. (2021). Priming soybean cv. primus leads to successful systemic defense against the rootlesion nematode, *Pratylenchus penetrans. Front. In Plant Sci.* 12, 651943. doi: 10.3389/fpls.2021.651943

Alegria Terrazas, R., Balbirnie-Cumming, K., Morris, J., Hedley, P. E., Russell, J., Paterson, E., et al. (2020). A footprint of plant eco-geographic adaptation on the composition of the barley rhizosphere bacterial microbiota. *Sci. Rep.* 10, 1–13. doi: 10.1101/2020.02.11.944140

An, G.-H., Kobayashi, S., Enoki, H., Sonobe, K., Muraki, M., Karasawa, T., et al. (2010). How does arbuscular mycorrhizal colonization vary with host plant genotype? an example based on maize (*Zea mays*) germplasms. *Plant And Soil* 327, 441-453. doi: 10.1007/s11104-009-0073-3

Ballabio, C., Panagos, P., Lugato, E., Huang, J.-H., Orgiazzi, A., Jones, A., et al. (2018). Copper distribution in European topsoils: An assessment based on Lucas soil survey. *Sci. Of Total Environ.* 636, 282–298. doi: 10.1016/j.scitotenv.2018.04.268

Bziuk, N., Maccario, L., Straube, B., Wehner, G., Sørensen, S. J., Schikora, A., et al. (2021). The treasure inside barley seeds: Microbial diversity and plant beneficial bacteria. *Environ. Microbiome.* 16, 1–21. doi: 10.1186/s40793-021-00389-8

Caddell, D. F., Deng, S., and Coleman-Derr, D. (2019). "Role of the plant root microbiome in abiotic stress tolerance," in *Seed endophytes* (Springer).

Cantalapiedra, C. P., Boudiar, R., Casas, A. M., Igartua, E., and Contreras-Moreira, B. (2015). Barleymap: Physical and genetic mapping of nucleotide sequences and annotation of surrounding loci in barley. *Mol. Breed.* 35, 1–11. doi: 10.1007/s11032-015-0253-1

Chen, C., Jost, M., Clark, B., Martin, M., Matny, O., Steffenson, B. J., et al. (2021). Bed domain-containing nlr from wild barley confers resistance to leaf rust. *Plant Biotechnol. J.* 19, 1206–1215. doi: 10.1111/pbi.13542

Chester, K. S. (1933). The problem of acquired physiological immunity in plants. *Q. Rev. Of Biol.* 8, 275–324. doi: 10.1086/394440

Clifford, B. (1985). "Barley leaf rust," in *Diseases, distribution, epidemiology, and control* (Elsevier).

Cockram, J., White, J., Leigh, F. J., Lea, V. J., Chiapparino, E., Laurie, D. A., et al. (2008). Association mapping of partitioning loci in barley. *BMC Genet.* 9, 1–14. doi: 10.1186/1471-2156-9-16

Comadran, J., Kilian, B., Russell, J., Ramsay, L., Stein, N., Ganal, M., et al. (2012). Natural variation in a homolog of antirrhinum centroradialis contributed to spring growth habit and environmental adaptation in cultivated barley. *Nat. Genet.* 44, 1388. doi: 10.1038/ng.2447

Conrath, U., Pieterse, C. M., and Mauch-Mani, B. (2002). Priming in plantpathogen interactions. *Trends In Plant Sci.* 7, 210–216. doi: 10.1016/S1360-1385 (02)02244-6

Desmedt, W., Vanholme, B., and Kyndt, T. (2021). Plant defense priming in the field: A review. *Recent Highlights In Discovery And Optimization Of Crop Prot. Products.* 5, 87–124. doi: 10.1016/B978-0-12-821035-2.00045-0

Dey, S., Wenig, M., Langen, G., Sharma, S., Kugler, K. G., Knappe, C., et al. (2014). Bacteria-triggered systemic immunity in barley is associated with wrky and ethylene responsive factors but not with salicylic acid. *Plant Physiol.* 166, 2133–2151. doi: 10.1104/pp.114.249276

Evanno, G., Regnaut, S., and Goudet, J. (2005). Detecting the number of clusters of individuals using the software structure: A simulation study. *Mol. Ecol.* 14, 2611–2620. doi: 10.1111/j.1365-294X.2005.02553.x

barley accessions and landraces of worldwide origin and including a test set of six common barley lines. * The UN 3-letter country code was used.

SUPPLEMENTARY FIGURE 2

QQ-Plot for model selection of a compressed mixed linear model (cMLM, **A**) with kinship as cofactor, cMLM with q-matrix and kinship as cofactors **(B)** and a generalized linear model (GLM, **C**).

SUPPLEMENTARY FIGURE 3

Manhattan plots for relative infection of *P. hordei* for 200 accessions primed with *E. meliloti* strain attM (red) or expR+ch (blue) against physical positions (Mb) of the marker trait associations on the seven barley chromosomes.

Fagorzi, C., Ilie, A., Decorosi, F., Cangioli, L., Viti, C., Mengoni, A., et al. (2020). Symbiotic and nonsymbiotic members of the genus *Ensifer* (Syn. *Sinorhizobium*) are separated into two clades based on comparative genomics and high-throughput phenotyping. *Genome Biol. And Evol.* 12, 2521–2534. doi: 10.1093/gbe/evaa221

Fazlikhani, L., Keilwagen, J., Kopahnke, D., Deising, H., Ordon, F., and Perovic, D. (2019). High resolution mapping of *Rph* mbr 1012 conferring resistance to *Puccinia hordei* in barley (*Hordeum vulgare* l.). *Front. In Plant Sci.* 10, 640. doi: 10.3389/fpls.2019.00640

Fuqua, W. C., Winans, S. C., and Greenberg, E. P. (1994). Quorum sensing in bacteria: The luxr-luxi family of cell density-responsive transcriptional regulators. *J. Of Bacteriology* 176, 269–275. doi: 10.1128/jb.176.2.269-275.1994

Galván, G. A., Kuyper, T. W., Burger, K., Keizer, L., Hoekstra, R. F., Kik, C., et al. (2011). Genetic analysis of the interaction between allium species and arbuscular mycorrhizal fungi. *Theor. And Appl. Genet.* 122, 947–960. doi: 10.1007/s00122-010-1501-8

Gantner, S., Schmid, M., Dürr, C., Schuhegger, R., Steidle, A., Hutzler, P., et al. (2006). *In situ* quantitation of the spatial scale of calling distances and population density-independent n-acylhomoserine lactone-mediated communication by rhizobacteria colonized on plant roots. *FEMS Microbiol. Ecol.* 56, 188–194. doi: 10.1111/j.1574-6941.2005.00037.x

Gao, J., Bi, W., Li, H., Wu, J., Yu, X., Liu, D., et al. (2018). Wrky transcription factors associated with Npr1-mediated acquired resistance in barley are potential resources to improve wheat resistance to *Puccinia triticina*. *Front. In Plant Sci.*, 1486. doi: 10.3389/fpls.2018.01486

Gdanetz, K., and Trail, F. (2017). The wheat microbiome under four management strategies, and potential for endophytes in disease protection. *Phytobiomes* 1, 158–168. doi: 10.1094/PBIOMES-05-17-0023-R

Guo, P., Baum, M., Grando, S., Ceccarelli, S., Bai, G., Li, R., et al. (2009). Differentially expressed genes between drought-tolerant and drought-sensitive barley genotypes in response to drought stress during the reproductive stage. *J. Of Exp. Bot.* 60, 3531–3544. doi: 10.1093/jxb/erp194

Haseneyer, G., Stracke, S., Paul, C., Einfeldt, C., Broda, A., Piepho, H. P., et al. (2010). Population structure and phenotypic variation of a spring barley world collection set up for association studies. *Plant Breed.* 129, 271–279. doi: 10.1111/j.1439-0523.2009.01725.x

Henry, R. J. (1985). A comparison of the non-starch carbohydrates in cereal grains. J. Of Sci. Of Food And Agric. 36, 1243–1253. doi: 10.1002/jsfa.2740361207

Hernández-Reyes, C., Schenk, S. T., Neumann, C., Kogel, K. H., and Schikora, A. (2014). N-Acyl-Homoserine lactones-producing bacteria protect plants against plant and human pathogens. *Microbial Biotechnol.* 7, 580–588. doi: 10.1111/1751-7915.12177

Hetrick, B., Wilson, G., and Cox, T. (1993). Mycorrhizal dependence of modern wheat cultivars and ancestors: A synthesis. *Can. J. Of Bot.* 71, 512–518. doi: 10.1139/b93-056

Hill, W., and Weir, B. (1988). Variances and covariances of squared linkage disequilibria in finite populations. *Theor. Population Biol.* 33, 54–78. doi: 10.1016/0040-5809(88)90004-4

Hou, Z., Yang, Y., Hedtke, B., and Grimm, B. (2019). Fluorescence in blue light (Flu) is involved in inactivation and localization of glutamyl-trna reductase during light exposure. *Plant J.* 97, 517–529. doi: 10.1111/tpj.14138

Hrmova, M., Banik, M., Harvey, A. J., Garrett, T. P., Varghese, J. N., Høj, P. B., et al. (1997). Polysaccharide hydrolases in germinated barley and their role in the depolymerization of plant and fungal cell walls. *Int. J. Of Biol. Macromolecules* 21, 67–72. doi: 10.1016/S0141-8130(97)00043-3

Huang, Y., Li, L., Smith, K. P., and Muehlbauer, G. J. (2016). Differential transcriptomic responses to *Fusarium graminearum* infection in two barley quantitative trait loci associated with fusarium head blight resistance. *BMC Genomics* 17, 1–16. doi: 10.1186/s12864-016-2716-0

Jack, C. N., Wozniak, K. J., Porter, S. S., and Friesen, M. L. (2019). Rhizobia protect their legume hosts against soil-borne microbial antagonists in a host-Genotype-Dependent manner. *Rhizosphere* 9, 47–55. doi: 10.1016/j.rhisph.2018.11.005

Janusz, G., Pawlik, A., Świderska-Burek, U., Polak, J., Sulej, J., Jarosz-Wilkołazka, A., et al. (2020). Laccase properties, physiological functions, and evolution. *Int. J. Of Mol. Sci.* 21, 966. doi: 10.3390/ijms21030966

Jung, H. W., Tschaplinski, T. J., Wang, L., Glazebrook, J., and Greenberg, J. T. (2009). Priming in systemic plant immunity. *Science* 324, 89–91. doi: 10.1126/science.1170025

Kaeppler, S. M., Parke, J. L., Mueller, S. M., Senior, L., Stuber, C., and Tracy, W. F. (2000). Variation among maize inbred lines and detection of quantitative trait loci for growth At low phosphorus and responsiveness to arbuscular mycorrhizal fungi. *Crop Sci.* 40, 358–364. doi: 10.2135/cropsci2000.402358x

Kaplan, H. B., and Greenberg, E. (1985). Diffusion of autoinducer is involved in regulation of the vibrio fischeri luminescence system. *J. Of Bacteriology* 163, 1210–1214. doi: 10.1128/jb.163.3.1210-1214.1985

Karlsson, I., Persson, P., and Friberg, H. (2021). Fusarium head blight from a microbiome perspective. *Front. In Microbiol.* 12, 628373. doi: 10.3389/fmicb.2021.628373

Kato, Y., and Sakamoto, W. (2018). Ftsh protease in the thylakoid membrane: Physiological functions and the regulation of protease activity. *Front. In Plant Sci.* 9, 855. doi: 10.3389/fpls.2018.00855

Kavanagh, P. J., Singh, D., Bansal, U. K., and Park, R. F. (2017). Inheritance and characterization of the new and rare gene *Rph25* conferring seedling resistance in *Hordeum vulgare* against *Puccinia hordei*. *Plant Breed*. 136, 908–912. doi: 10.1111/ pbr.12535

Kuźniar, A., Włodarczyk, K., Grządziel, J., Goraj, W., Gałązka, A., and Wolińska, A. (2020). Culture-independent analysis of an endophytic core microbiome in two species of wheat: *Triticum aestivum* l. (Cv.'Hondia') and the first report of microbiota in *Triticum spelta* l. (Cv.'Rokosz'). *Systematic And Appl. Microbiol.* 43, 126025. doi: 10.1016/j.syapm.2019.126025

Lázaro, E., Makowski, D., and Vicent, A. (2021). Decision support systems halve fungicide use compared to calendar-based strategies without increasing disease risk. *Commun. Earth Environ.* 2, 1–10. doi: 10.1038/s43247-021-00291-8

Lehnert, H., Serfling, A., Enders, M., Friedt, W., and Ordon, F. (2017). Genetics of mycorrhizal symbiosis in winter wheat (*Triticum aestivum*). New Phytol. 215, 779–791. doi: 10.1111/nph.14595

Lehnert, H., Serfling, A., Friedt, W., and Ordon, F. (2018). Genome-wide association studies reveal genomic regions associated with the response of wheat (Triticum aestivum l.) to mycorrhizae under drought stress conditions. *Front. In Plant Sci.* 9, 1728. doi: 10.3389/fpls.2018.01728

Levine, M. N., and Cherewick, W. J. (1952). *Studies on dwarf leaf rust of barley* (Us Department Of Agriculture).

Liljenberg, C. (1977). The occurrence of phytylpyrophosphate and acyl esters of phytol in irradiated dark-grown barley seedlings and their possible role in biosynthesis of chlorophyll. *Physiologia Plantarum* 39, 101–105. doi: 10.1111/j.1399-3054.1977.tb04017.x

Lipka, A. E., Tian, F., Wang, Q., Peiffer, J., Li, M., Bradbury, P. J., et al. (2012). Gapit: Genome association and prediction integrated tool. *Bioinformatics* 28, 2397–2399. doi: 10.1093/bioinformatics/bts444

Lippold, F., Vom Dorp, K., Abraham, M., Hölzl, G., Wewer, V., et al. (2012). Fatty acid phytyl ester synthesis in chloroplasts of arabidopsis. *Plant Cell* 24, 2001–2014. doi: 10.1105/tpc.112.095588

Liu, H., Brettell, L. E., Qiu, Z., and Singh, B. K. (2020). Microbiome-mediated stress resistance in plants. *Trends In Plant Sci.* 25, 733-743. doi: 10.1016/j.tplants.2020.03.014

Liu, D., Leib, K., Zhao, P., Kogel, K.-H., and Langen, G. (2014). Phylogenetic analysis of barley wrky proteins and characterization of Hvwrky1 and-2 as repressors of the pathogen-inducible gene Hvger4c. *Mol. Genet. And Genomics* 289, 1331–1345. doi: 10.1007/s00438-014-0893-6

Mahapatra, S., Rayanoothala, P., Solanki, M. K., and Das, S. (2020). Wheat microbiome: Present status and future perspective. *Phytobiomes: Curr. Insights And Future Vistas*, 191–223. doi: 10.1007/978-981-15-3151-4_8

Mascher, M., Gundlach, H., Himmelbach, A., Beier, S., Twardziok, S. O., Wicker, T., et al. (2017). A chromosome conformation capture ordered sequence of the barley genome. *Nature* 544, 427–433. doi: 10.1038/nature22043

Mascher, M., Richmond, T. A., Gerhardt, D. J., Himmelbach, A., Clissold, L., Sampath, D., et al. (2013). Barley whole exome capture: A tool for genomic research in the genus hordeum and beyond. *Plant J.* 76, 494–505. doi: 10.1111/tpj.12294 Mascher, M., Wicker, T., Jenkins, J., Plott, C., Lux, T., Koh, C. S., et al. (2021). Long-read sequence assembly: A technical evaluation in barley. *Plant Cell* 33, 1888–1906. doi: 10.1093/plcell/koab077

Mauch-Mani, B., Baccelli, I., Luna, E., and Flors, V. (2017). Defense priming: An adaptive part of induced resistance. *Annu. Rev. Of Plant Biol.* 68, 485–512. doi: 10.1146/annurev-arplant-042916-041132

Meng, Y., and Wise, R. P. (2012). Hvwrky10, Hvwrky19, and Hvwrky28 regulate mla-triggered immunity and basal defense to barley powdery mildew. *Mol. Plant-Microbe Interact.* 25, 1492–1505. doi: 10.1094/MPMI-04-12-0082-R

Moll, E., Flath, K., and Piepho, H. (2000). Die prüfung Von pflanzen auf ihre widerstandsfähigkeit gegen schadorganismen in der biologischen bundesanstalt teil 3: Methodische anleitung zur bewertung der partiellen resistenz Von getreidesortimenten und die sas-applikation resi mitteilungen aus der biologischen bundesanstalt für land-und forstwirtschaft. *Heft* 374, 128.

Morella, N. M., Weng, F. C.-H., Joubert, P. M., Metcalf, C. J. E., Lindow, S., and Koskella, B. (2020). Successive passaging of a plant-associated microbiome reveals robust habitat and host genotype-dependent selection. *Proc. Of Natl. Acad. Of Sci.* 117, 1148–1159. doi: 10.1073/pnas.1908600116

Niks, R., Walther, U., Jaiser, H., Martinez, F., and Rubiales, D. (2000). Resistance against barley leaf rust (*Puccinia hordei*) in West-European spring barley germplasm. *Agronomie* 20, 769–782. doi: 10.1051/agro:2000174

Novakazi, F., Afanasenko, O., Anisimova, A., Platz, G. J., Snowdon, R., Kovaleva, O., et al. (2019). Genetic analysis of a worldwide barley collection for resistance to net form of net blotch disease (*Pyrenophora teres f. teres*). *Theor. And Appl. Genet.* 132, 2633–2650. doi: 10.1007/s00122-019-03378-1

Park, R. (2003). Pathogenic specialization and pathotype distribution of puccinia hordei in australia 1992 To 2001. *Plant Dis.* 87, 1311–1316. doi: 10.1094/PDIS.2003.87.11.1311

Parke, J. L., and Kaeppler, S. W. (2000). "Effects of genetic differences among crop species and cultivars upon the arbuscular mycorrhizal symbiosis," in *Arbuscular mycorrhizas: Physiology and function* (Springer).

Park, R. F., Golegaonkar, P. G., Derevnina, L., Sandhu, K. S., Karaoglu, H., Elmansour, H. M., et al. (2015). Leaf rust of cultivated barley: Pathology and control. *Annu. Rev. Phytopathol.* 53, 565–589. doi: 10.1146/annurev-phyto-080614-120324

Park, R. F., and Karakousis, A. (2002). Characterization and mapping of gene *Rph19* conferring resistance to *Puccinia hordei* in the cultivar 'Reka 1'and several Australian barleys. *Plant Breed.* 121, 232–236. doi: 10.1046/j.1439-0523.2002.00717.x

Pasam, R. K., Sharma, R., Malosetti, M., Van Eeuwijk, F. A., Haseneyer, G., Kilian, B., et al. (2012). Genome-wide association studies for agronomical traits in a world wide spring barley collection. *BMC Plant Biol.* 12, 1–22. doi: 10.1186/1471-2229-12-16

Perlin, D. S., Rautemaa-Richardson, R., and Alastruey-Izquierdo, A. (2017). The global problem of antifungal resistance: Prevalence, mechanisms, and management. *Lancet Infect. Dis.* 17, E383–E392. doi: 10.1016/S1473-3099(17) 30316-X

Poland, J. A., and Rife, T. W. (2012). Genotyping-By-Sequencing for plant breeding and genetics. *Plant Genome*. 5, 92–102. doi: 10.3835/plantgenome2012.05.0005

Powell, C. L., Clark, G., and Verberne, N. (1982). Growth response of four onion cultivars to several isolates of va mycorrhizal fungi. *New Z. J. Of Agric. Res.* 25, 465–470. doi: 10.1080/00288233.1982.10417914

Pritchard, J. K., Stephens, M., and Donnelly, P. (2000). Inference of population structure using multilocus genotype data. *Genetics* 155, 945–959. doi: 10.1093/genetics/155.2.945

Ribaudo, C., Zaballa, J. I., and Golluscio, R. (2020). Effect of the phosphorussolubilizing bacterium *Enterobacter ludwigii* on barley growth promotion. *Am. Acad. Sci. Res. J. For Engineering Technology And Sci.* 63, 144–157.

Rostoks, N., Ramsay, L., Mackenzie, K., Cardle, L., Bhat, P. R., Roose, M. L., et al. (2006). Recent history of artificial outcrossing facilitates whole-genome association mapping in elite inbred crop varieties. *Proc. Of Natl. Acad. Of Sci.* 103, 18656–18661. doi: 10.1073/pnas.0606133103

Sawers, R. J., Gebreselassie, M. N., Janos, D. P., and Paszkowski, U. (2010). Characterizing variation in mycorrhiza effect among diverse plant varieties. *Theor. And Appl. Genet.* 120, 1029–1039. doi: 10.1007/s00122-009-1231-y

Schenk, S. T., and Schikora, A. (2015). Ahl-priming functions Via oxylipin and salicylic acid. Front. In Plant Sci. 5, 784. doi: 10.3389/fpls.2014.00784

Schuhegger, R., Ihring, A., Gantner, S., Bahnweg, G., Knappe, C., Vogg, G., et al. (2006). Induction of systemic resistance in tomato by n-Acyl-L-Homoserine lactone-producing rhizosphere bacteria. *Plant Cell Environ.* 29, 909–918. doi: 10.1111/j.1365-3040.2005.01471.x

Shrestha, A., Elhady, A., Adss, S., Wehner, G., Böttcher, C., Heuer, H., et al. (2019). Genetic differences in barley govern the responsiveness to n-acyl

homoserine lactone. Phytobiomes J. 3, 191-202. doi: 10.1094/PBIOMES-03-19-0015-R

Shrestha, A., Grimm, M., Ojiro, I., Krumwiede, J., and Schikora, A. (2020). Impact of quorum sensing molecules on plant growth and immune system. *Front. In Microbiol.* 11, 1545. doi: 10.3389/fmicb.2020.01545

Shrestha, A., and Schikora, A. (2020). Ahl-priming for enhanced resistance as a tool in sustainable agriculture. *FEMS Microbiol. Ecol.* 96, Fiaa226. doi: 10.1093/femsec/fiaa226

Si, J., Froussart, E., Viaene, T., Vázquez-Castellanos, J. F., Hamonts, K., Tang, L., et al. (2021). Interactions between soil compositions and the wheat root microbiome under drought stress: From an *In silico* to *In planta* perspective. *Comput. And Struct. Biotechnol. J.* 19, 4235–4247. doi: 10.1016/ j.csbj.2021.07.027

Singh, D., Dracatos, P., Derevnina, L., Zhou, M., and Park, R. F. (2015). *Rph23*: A new designated additive adult plant resistance gene to leaf rust in barley on chromosome 7h. *Plant Breed*. 134, 62–69. doi: 10.1111/pbr.12229

Springer, A., Acker, G., Bartsch, S., Bauerschmitt, H., Reinbothe, S., and Reinbothe, C. (2015). Differences in gene expression between natural and artificially induced leaf senescence in barley. *J. Of Plant Physiol.* 176, 180–191. doi: 10.1016/j.jplph.2015.01.004

Starkey, R. L. (1958). Interrelations between microorganisms and plant roots in the rhizosphere. *Bacteriological Rev.* 22, 154–172. doi: 10.1128/br.22.3.154-172.1958

Tawaraya, K., Tokairin, K., and Wagatsuma, T. (2001). Dependence of Allium fistulosum cultivars on the arbuscular mycorrhizal fungus, *Glomus fasciculatum*. Appl. Soil Ecol. 17, 119–124. doi: 10.1016/S0929-1393(01)00126-3

Teplitski, M., Eberhard, A., Gronquist, M. R., Gao, M., Robinson, J. B., and Bauer, W. D. (2003). Chemical identification of n-acyl homoserine lactone quorum-sensing signals produced by *Sinorhizobium meliloti* strains in defined medium. *Arch. Of Microbiol.* 180, 494–497. doi: 10.1007/s00203-003-0612-x

Thabet, S. G., Alomari, D. Z., Brinch-Pedersen, H., and Alqudah, A. M. (2022). Genetic analysis toward more nutritious barley grains for a food secure world. *Botanical Stud.* 63, 1–11. doi: 10.1186/s40529-022-00334-z

Tomková, L., Kučera, L., Vaculová, K., and Milotová, J. (2012). Characterization and mapping of a putative laccase-like multicopper oxidase gene in the barley (*Hordeum vulgare* l.). *Plant Sci.* 183, 77–85. doi: 10.1016/j.plantsci.2011.11.003

Trivedi, P., Leach, J. E., Tringe, S. G., Sa, T., and Singh, B. K. (2020). Plantmicrobiome interactions: From community assembly to plant health. *Nat. Rev. Microbiol.* 18, 607–621. doi: 10.1038/s41579-020-0412-1

Trivedi, P., Mattupalli, C., Eversole, K., and Leach, J. E. (2021). Enabling sustainable agriculture through understanding and enhancement of microbiomes. *New Phytol.* 230, 2129–2147. doi: 10.1111/nph.17319

Uluhan, E., Keleş, E. N., and Tufan, F. (2019). Analysis of wrky transcription factors in barley cultivars infected with *Fusarium culmorum*. Int. J. Of Life Sci. And Biotechnol. 2, 165–174. doi: 10.38001/ijlsb.588730

Voorrips, R. (2002). Mapchart: Software for the graphical presentation of linkage maps and qtls. J. Of Heredity 93, 77–78. doi: 10.1093/jhered/93.1.77

Wehner, G., Kopahnke, D., Richter, K., Kecke, S., Schikora, A., and Ordon, F. (2019). Priming is a suitable strategy to enhance resistance towards leaf rust in barley. *Phytobiomes J.* 3, 46–51. doi: 10.1094/PBIOMES-09-18-0041-R

Whelan, H., Gaunt, R., and Scott, W. (1997). The effect of leaf rust (*Puccinia hordei*) on yield response in barley (*Hordeum vulgare* l.) crops with different yield potentials. *Plant Pathol.* 46, 397–406. doi: 10.1046/j.1365-3059.1997.d01-23.x

Yang, L., Danzberger, J., Schöler, A., Schröder, P., Schloter, M., and Radl, V. (2017). Dominant groups of potentially active bacteria shared by barley seeds become less abundant in root associated microbiome. *Front. In Plant Sci.* 8, 1005. doi: 10.3389/fpls.2017.01005

Yang, L., Schröder, P., Vestergaard, G., Schloter, M., and Radl, V. (2020). Response of barley plants to drought might be associated with the recruiting of soilborne endophytes. *Microorganisms* 8, 1414. doi: 10.3390/microorganisms8091414

Zarkani, A. A., Stein, E., Röhrich, C. R., Schikora, M., Evguenieva-Hackenberg, E., Degenkolb, T., et al. (2013). Homoserine lactones influence the reaction of plants to rhizobia. *Int. J. Of Mol. Sci.* 14, 17122–17146. doi: 10.3390/ijms140817122

Žiarovská, J., Medo, J., Kyseľ, M., Zamiešková, L., and Kačániová, M. (2020). Endophytic bacterial microbiome diversity in early developmental stage plant tissues of wheat varieties. *Plants* 9, 266. doi: 10.3390/plants9020266