



Corrigendum: To Be or Not to Be Expressed: The First Evidence of a Nucleolar Dominance Tissue-Specificity in *Brachypodium hybridum*

Natalia Borowska-Zuchowska^{1*}, Ewa Robaszkiewicz¹, Serhii Mykhailyk¹, Joanna Wartini¹, Artur Pinski¹, Ales Kovarik² and Robert Hasterok¹

¹ Plant Cytogenetics and Molecular Biology Group, Institute of Biology, Biotechnology and Environmental Protection, Faculty of Natural Sciences, University of Silesia in Katowice, Katowice, Poland, ² Department of Molecular Epigenetics, Institute of Biophysics, Academy of Sciences of the Czech Republic, Brno, Czechia

OPEN ACCESS

Approved by:
Frontiers Editorial Office,
Frontiers Media SA, Switzerland

***Correspondence:**
Natalia Borowska-Zuchowska
natalia.borowska@us.edu.pl

Specialty section:
This article was submitted to
Plant Cell Biology,
a section of the journal
Frontiers in Plant Science

Received: 11 April 2022
Accepted: 12 April 2022
Published: 28 April 2022

Citation:
Borowska-Zuchowska N,
Robaszkiewicz E, Mykhailyk S,
Wartini J, Pinski A, Kovarik A and
Hasterok R (2022) Corrigendum: To
Be or Not to Be Expressed: The First
Evidence of a Nucleolar Dominance
Tissue-Specificity in *Brachypodium
hybridum*.
Front. Plant Sci. 13:917940.
doi: 10.3389/fpls.2022.917940

Keywords: nucleolar dominance, 35S rDNA, secondary constriction, *Brachypodium*, allopolyploidy, 3D-FISH, rRNA gene expression

A Corrigendum on

To Be or Not to Be Expressed: The First Evidence of a Nucleolar Dominance Tissue-Specificity in *Brachypodium hybridum*

by Borowska-Zuchowska, N., Robaszkiewicz, E., Mykhailyk, S., Wartini, J., Pinski, A., Kovarik, A., and Hasterok, R. (2021). *Front. Plant Sci.* 12:768347. doi: 10.3389/fpls.2021.768347

In the original article, there was a mistake in **Figure 3B** as published. The gel lines were marked in an incorrect order. The corrected **Figure 3B** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Borowska-Zuchowska, Robaszkiewicz, Mykhailyk, Wartini, Pinski, Kovarik and Hasterok. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 3 | The 35S rDNA expression analysis in the different tissues of *B. hybridum* (genotypes 3-7-2 and ABR113) and *B. stacei* (genotype ABR114; control) and *B. distachyon* (genotype Bd21; control) using the RT-PCR CAPS method. **(A)** The *MluI* restriction profile of the *B. distachyon*-like and *B. stacei*-like ITS1 PCR products. The expected sizes of the bands after *MluI* digestion are presented. **(B)** The *MluI* restriction profiles of ITS1 amplification products that were obtained from the leaves cDNAs of *B. distachyon* and *B. stacei*. **(C)** The *MluI* restriction profiles of ITS1 amplification products that were obtained from the primary roots, leaves, adventitious roots, and immature spikes cDNAs. Adv. Roots, adventitious roots.