Check for updates

OPEN ACCESS

EDITED BY Ignacio García-González, University of Santiago de Compostela, Spain

REVIEWED BY Furong Niu, University of Arizona, United States Junzhou Zhang, Lanzhou University, China

*CORRESPONDENCE Keyan Fang kfang@fjnu.edu.cn Feifei Zhou zhouff1987@126.com

[†]These authors have contributed equally to this work

RECEIVED 30 January 2025 ACCEPTED 30 May 2025 PUBLISHED 25 June 2025

CITATION

Wang C, Zheng Z, Gao J, Zhou F, Rossi S and Fang K (2025) Linking growth dynamics and intra-annual density fluctuations to late-summer precipitation in humid subtropical China. *Front. Plant Sci.* 16:1568882. doi: 10.3389/fpls.2025.1568882

COPYRIGHT

© 2025 Wang, Zheng, Gao, Zhou, Rossi and Fang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Linking growth dynamics and intra-annual density fluctuations to late-summer precipitation in humid subtropical China

Chunsong Wang^{1,2†}, Zhuangpeng Zheng^{3†}, Jiani Gao^{1,2}, Feifei Zhou^{1,2*}, Sergio Rossi⁴ and Keyan Fang^{1,2*}

¹Institute of Geography, Fujian Normal University, Fuzhou, China, ²Key Laboratory of Humid Subtropical Eco-Geographical Process (Ministry of Education), College of Geographical Sciences, Fujian Normal University, Fuzhou, China, ³School of Tourismand Historical Culture, Zhaoqing University, Zhaoqing, China, ⁴Département des Sciences Fondamentales, Université du Québec à Chicoutimi, Chicoutimi, QC, Canada

Global warming has intensified extreme rainfall events and prolonged droughts, significantly impacting tree growth and wood formation. This study investigates the effects of late-summer precipitation variability on the intra-annual growth dynamics of Cunninghamia lanceolata and Cryptomeria fortunei in humid subtropical China. Microcores were collected from 12 trees at 7-10 days intervals between March and December from 2021 to 2023 in the Gushan Mountains. Typically, high temperatures and rainfall deficits in July induce cambial dormancy, while subsequent rainfall in August and September reactivates growth, resulting in a bimodal growth pattern. However, in 2022, an unprecedented drought (August-October rainfall 77% below average) shortened the growing season, causing an early cessation of growth and a rare unimodal growth pattern. In contrast, persistent rainfall in 2023 accelerated cell enlargement to 7 μ m d⁻¹ and significantly increased latewood intra-annual density fluctuations (L-IADFs). Notably, despite abundant late-summer rainfall in 2021, L-IADFs did not form, indicating a nonlinear and inconsistent relationship between rainfall and L-IADFs. These findings highlight the critical role of latesummer precipitation variability in shaping tree growth patterns and wood density in southeastern China. Given the expected increase in precipitation variability under climate change, regional forest ecosystems may become more vulnerable. This study provides valuable insights for forest management strategies to enhance resilience and mitigate climate-related risks.

KEYWORDS

xylogenesis, wood anatomy, autumn growth peak, L-IADFs, extreme climate events, Southeastern China

1 Introduction

In the recent decades, global warming has led to significant disruptions in seasonal precipitation patterns, intensifying extreme rainfall events and prolonging dry periods (Iles et al., 2024; IPCC, 2021; Feng et al., 2013). These climatic shifts have profound implications for forest ecosystems, as they affect the dynamics of wood formation processes, contributing to an increased tree mortality and forest degradation (Feldman et al., 2024; Zhou et al., 2024; Allen et al., 2010). Wood formation is sensitive to fluctuations in temperature and precipitation, which directly influence xylem development and lead to significant variations in growth patterns (Rathgeber et al., 2016; Rossi et al., 2016). Understanding how these climate-induced changes affect wood formation is critical for predicting the resilience and dynamics of forest ecosystems under global warming.

In temperate and alpine regions, xylem growth typically follows a unimodal pattern, with earlywood formation in spring and latewood development in late summer, followed by growth cessation in autumn (Huang et al., 2023; Rossi et al., 2006b). However, in regions with more complex climates, such as the Mediterranean and humid subtropical areas, trees often experience summer droughts, leading to distinct growth dynamics and anatomical patterns in xylem formation (Zlobin, 2022; Vieira et al., 2019). Summer droughts in these regions can induce bimodal growth patterns, with growth peaks in both spring and autumn (Häusser et al., 2023; Wang C. S. et al., 2023; Jiang et al., 2021). Autumn growth peaks are generally associated with the resumption of precipitation, which stimulates cambial activity and promotes latewood cell enlargement (Niccoli et al., 2024; Zalloni et al., 2016).

Autumn growth peaks often correspond to the formation of latewood intra-annual density fluctuations (L-IADFs) in the Mediterranean, characterized by earlywood-like cells with large lumens and thin walls within latewood. These fluctuations are typically triggered by abrupt changes in moisture availability (Balzano et al., 2019). In humid subtropical regions, where there is a pronounced shift from summer drought to autumn rainfall (Ma et al., 2021; Xu et al., 2019), climatic fluctuations drive adaptive changes in xylem width and structure, significantly influencing wood formation dynamics (Häusser et al., 2023; Zhang et al., 2020). As indicators of environmental variability, L-IADFs provide insights into how trees respond to climatic stressors such as droughts and sudden changes in moisture, reflecting their growth strategies and adaptive capacity (Niccoli et al., 2024; Gao et al., 2021a). These wood structure traits highlight the ability of trees to adjust their architecture for efficient water transport and mechanical support-key factors for forest resilience in a changing climate (Zalloni et al., 2016). Understanding these traits is crucial for predicting the long-term evolution of forests under future climate change.

Despite extensive research on the frequency of IADFs in tree rings, the underlying physiological mechanisms driving their formation remain poorly understood. Most studies have focused on documenting the occurrence and frequency of IADFs, with limited attention given to the physiological processes and environmental drivers behind their formation (Wang S.J. et al., 2023; De Micco et al., 2016). This knowledge gap hinders our understanding of the factors influencing L-IADFs formation, especially in regions with significant precipitation variability. Recent advances in xylem kinetics offer promising approaches to bridge this gap by analyzing the timing and rates of xylem cell formation alongside anatomical changes. This integrated approach can provide deeper insights into the processes of cell division, enlargement, and wall thickening (Cuny et al., 2014), helping to identify critical periods when climatic factors, such as precipitation, promote or inhibit xylem formation. For example, Gao et al. (2021b) demonstrated that water scarcity alters both the timing of cell division and the rate of cell enlargement, which ultimately influences xylem morphology. These findings emphasize the importance of linking wood anatomy and xylem dynamics to better understand L-IADF formation and its response to climatic drivers.

This study focuses on two conifer species in humid subtropical China: *Cunninghamia lanceolata* and *Cryptomeria fortunei*. These species were chosen due to their widespread distribution and ecological importance in the region. We aim to investigate the anatomical characteristics of xylem cells and analyze the dynamics of xylem formation in response to variability in late-summer precipitation. Using microcores collected over three consecutive years, we explore how late-summer precipitation influences xylem cell development, with particular attention to autumn growth peaks and L-IADF formation. We hypothesize that: (1) the amount and timing of late-summer precipitation control the intensity and duration of the autumn growth peak; (2) continuous heavy precipitation promotes cell enlargement and L-IADFs formation; and (3) a growth peak in autumn does not always correspond to L-IADFs formation.

2 Materials and methods

2.1 Study area

This study was conducted in the Gushan Mountains, a humid subtropical region of southeastern China (26.05°N, 119.38°E, 450 m a.s.l.), strongly influenced by the East Asian monsoon (Figure 1A). The forest canopy is predominantly composed of Cunninghamia lanceolata (C. lanceolata) and Cryptomeria fortune (C. fortunei), and contains minor species of Pinus massoniana and Schima superba (Zheng et al., 2021). The area experiences a humid subtropical climate, with a mean annual temperature of 20.1°C (1951-2020). July experiences the highest monthly temperature of 34.29°C, while January is the coldest month at 8.16°C. The area receives an average annual precipitation of 1387 mm, 72% of which occurs between March and August. A typical summer drought occurs in July (Figure 1B). Daily climate data, including precipitation, temperature, soil moisture, and relative humidity, were collected from the Fuzhou Meteorological Station (26.08°N, 119.28°E, 85 m a.s.l.), located 11.2 km away from the study site. Data were accessed from the China Meteorological Administration (https://weather.cma.cn/).



2.2 Tree selection and sample collection

Seven healthy C. lanceolata and five C. fortunei trees were selected for monitoring stem radial growth. The selected trees were of similar dominance and had well-developed crowns. The C. lanceolata trees had an average diameter at breast height (DBH) of 27.4 \pm 2.7 cm (mean \pm SD), a height of 7.9 \pm 1.0 m, and an estimated age of 35 ± 2 years (mean \pm SD). The C. fortunei trees had an average DBH of 28.4 \pm 4.8 cm (mean \pm SD), a height of 12.8 \pm 1.5 m, and an estimated age of 31.6 \pm 1.4 years (mean \pm SD). Microcores were collected at 7-10-day intervals during the growing seasons (March to December) from 2021 to 2023 at breast height (1.3 m) using a Trephor (Rossi et al., 2006a). Samples were immediately preserved in a 9:1 solution of 70% alcohol and acetic acid. Each core, 2 mm in diameter and 2-3 cm in length, contained two intact tree rings and adjacent phloem. Sampling points were systematically arranged around the trunk to minimize interference, maintaining a minimum spacing of 5 cm between successive samples. In the laboratory, samples were dehydrated using an alcohol gradient (70%, 90%, 95%, and 100%), cleared with limonene, and embedded in paraffin. Transverse sections (8 µm thick) were cut with a rotary microtome (HistoCore BIOCUT, LEICA, Wetzlar, Germany) and stained with safranin and Astra blue (Rossi et al., 2003; Rossi et al., 2006b).

2.3 Xylem formation dynamics

Transverse sections were examined under visible and polarized light using a 200× optical microscope (DM750P, LEICA, Wetzlar, Germany) to identify xylem cells at different differentiation stages. Cambium cells were recognized as flattened cells with thin walls, while enlarging cells appeared larger with irregular diameters. Wallthickening cells were identified by their birefringence under polarized light, and mature cells were characterized by fully developed structures, which stained red with safranin (Li et al., 2017). The onset of xylem differentiation was marked by the appearance of at least one row of enlarging cells in spring. Xylem cell division cessation in autumn was defined by the absence of enlarging cells, and differentiation was considered complete when no wall-thickening cells were present (Rossi et al., 2006b). Cell counts at each developmental stage were recorded during every sampling interval and correlated with the day of the year (DOY) to create a timeline of xylem development.

2.4 Quantification of xylem anatomical parameters

Digital images of xylem sections were captured at 200× magnification using a microscope with an integrated camera (MC190 HD, LEICA, Wetzlar, Germany). Anatomical parameters, such as radial lumen diameter (LD), cell wall thickness (CWT), and tracheid diameter (TD, where TD = LD + 2×CWT), were measured along five radial files using ImageJ software (National Institutes of Health, Bethesda, MD, USA). Mature cells were classified as earlywood (4×CWT/LD < 1) or latewood (4×CWT/LD ≥ 1) based on Mork's criterion (Denne, 1989). To assess intra-annual variability, each growth ring was divided into 10 equal radial sections. Anatomical parameters for all cells within each section were averaged, enabling segment-wise analysis of growth ring anatomical features. This approach facilitated the identification of intra-annual variations in tracheid characteristics (Silvestro et al., 2023).

2.5 Data analysis and statistics

To account for variations in growth rates across different trunk directions (Wang C. S. et al., 2023), xylem cell counts were standardized using the following formula:

$$nci = ncmi \times rwm/rws$$
 (1)

where nc_i is the standardized number of xylem cells, ncm_i is the measured number of cells, rw_m is the mean width of preceding tree rings across all samples, and rw_s is the width of preceding tree rings for the specific sample (Rossi et al., 2006b).

Year	Growing season		Early grow	ving season	Lare growing season	
	Temperature (°C)	Precipitation (mm)	Temperature (°C)	Precipitation (mm)	Temperature (°C)	Precipitation (mm)
2021	0.86	117.54	1.01	-62.31	0.73	179.86
2022	0.68	-173.54	-0.04	126.41	1.25	-299.95
2023	0.77	659.32	0.82	-88.22	0.73	747.55

TABLE 1 Difference from average value during 1980-2020 at Fuzhou meteorological station during 2021-2023 growing season.

Growing season (March-November), early growing season (March-June), late growing season (July-November).

The Gompertz function was applied to analyze xylem formation kinetics, estimating the duration of cell enlargement (d_E) and wall thickening (d_{WT}) phases. Rates of cell enlargement ($r_E=TD/d_E$) and wall thickening ($r_{WT}=2\times CWT/d_{WT}$) were calculated from the anatomical parameters of the percentage of the growth ring (Gao et al., 2021b; Palombo et al., 2018).

Vapor Pressure Deficit (VPD) was calculated using air temperature (T) and relative humidity (RH) as follows:

$$VPD = 0.6108 \times e^{(17.27T/(237.3+T)) \times (1-RH)}$$
(2)

Monthly climate indices and key phenological timings (2021– 2023) were compared using ANOVA following normality tests. Statistical analyses were conducted using IBM SPSS Statistics (version 25.0, SPSS Inc., Chicago, USA).

3 Results

3.1 Weather conditions during the study years

From March to November in 2021 to 2023, the climate deviated significantly from the 1980–2020 baseline. During the growing seasons, average temperatures were 0.86°C higher in 2021, 0.68°C higher in 2022, and 0.77°C higher in 2023. Precipitation exhibited substantial variability: an increase of 117.54 mm in 2021, a decrease of 173.54 mm in 2022, and a 50% increase (659.32 mm) in 2023.

Precipitation fluctuations were more pronounced from July to November. Compared to the 1980–2020 average, rainfall increased by 179.86 mm in 2021, decreased by 299.95 mm in 2022, and surged by 747.55 mm in 2023 (Table 1). A severe drought occurred between August and October 2022, leading to extremely low soil moisture and a high VPD. In contrast, 2023 experienced two consecutive typhoons, which brought significant rainfall, alleviating the effects of high temperatures and drought. A similar pattern was observed in 2021, although with less rainfall over a shorter period. Consequently, the moisture conditions during August–September differed significantly (p < 0.05) across the three study years (Table 2).

3.2 Xylem formation dynamics

Xylem formation dynamics exhibited notable variation across the years, though the overall patterns remained consistent for both *C. lanceolata* and *C. fortunei*. During the growing season, cambial cell numbers exhibited slight fluctuations, ranging from 3 to 5 in *C. lanceolata* and 3 to 6 in *C. fortunei* over the three years (2021– 2023). Notable inter-annual variability was observed in the dynamics of cell enlargement. In 2021, both species exhibited a bimodal growth pattern, with a primary peak in spring (around DOY 120) and a smaller second peak in autumn (around DOY 240), most pronounced in *C. fortunei*. In 2023, the bimodal pattern was even more pronounced, with a distinct autumn growth peak.

TABLE 2 Monthly average temperature, soil moisture and VPD of late growing season during 2021-2023.

Index	Year	July	August	September	October	November
Temperature	2021	30.03 ^a	28.57 ^a	28.89 ^a	24.63 ^a	18.26 ^a
(°C)	2022	29.81 ^a	30.76 ^b	28.33 ^a	23.81 ^a	20.19 ^b
	2023	29.89 ^a	28.78 ^a	28.16 ^a	23.85 ^a	19.71 ^b
Soil moisture	2021	35.42 ^a	38.59 ^a	34.51 ^a	35.2 ^a	35.84 ^a
(Kg/m ²)	2022	34.89 ^a	30.64 ^b	29.68 ^b	27.3 ^b	35.02 ^{ab}
	2023	35.87 ^a	38.01 ^a	36.99 ^c	34.6 ^a	34.2 ^b
VPD	2021	1.03 ^a	0.51 ^a	0.88 ^a	0.59 ^a	0.38 ^a
(kPa)	2022	1.17 ^a	1.37 ^b	1.14 ^b	0.82 ^b	0.38 ^a
	2023	1.02 ^a	0.64 ^a	0.61 ^c	0.55 ^a	0.54 ^b

Different letters indicate significant differences between years (p < 0.05).

In contrast, 2022 followed a unimodal pattern, with a single peak in spring. Cell wall thickening dynamics followed a similar trend to cell enlargement, though the peak occurred later due to the time required for wall thickening. Cell maturation followed an S-shaped curve, with the highest cell production recorded in 2023 and the lowest in 2022 (Figure 2).

3.3 Cambial phenology

Significant inter-annual variations in xylem formation, particularly during late summer and autumn, resulted in marked differences in cambial phenology. Due to road closures to Gushan Mountain from March to April 2022, data on the initiation of cambial activity were unavailable. In 2021, cambial activity began on DOY 95 \pm 7 for *C. lanceolata* and on DOY 89 \pm 7 for *C. fortunei*. In 2023, activity began on DOY 93 \pm 10 for C. lanceolata and on DOY 86 \pm 9 for *C. fortunei*, with no significant difference in onset timing between the two years. However, the cessation of xylem cell division and the end of the growing season showed considerable variation across years, except for C. fortunei in 2021 and 2023, where no significant differences were observed (Figure 3). Drought periods of varying intensity during late summer and autumn in 2022 and 2023 significantly advanced the timing of xylem cell division cessation (Figure 4). For C. lanceolata, cell division ceased on DOY 300 \pm 3 in 2021, occurring 60 days earlier in 2022 and 17 days earlier in 2023. The end of the growing season shifted from DOY 324 \pm 13 in 2021, advancing 44 days in 2022 and 13 days in 2023. For C. fortunei, the trend was consistent, with cell division ending 56 days earlier in 2022 and 17 days earlier in 2023 compared to DOY 300 \pm 3 in 2021. The end of the growing season advanced by 51 days in 2022 and 8 days in 2023, relative to DOY 335 \pm 4 in 2021 (Figure 3).

3.4 Xylem anatomical traits

Xylem cell anatomy exhibited significant inter- and intraannual variability. In C. lanceolata, the lumen diameter ranged from 2.45 µm to 39.6 µm. It consistently decreased from 10% of the growth ring, except for a modest enlargement at 80% across the tree ring in 2023 (Figure 5A). Cell wall thickness increased, starting at 2.18 µm and reaching a maximum of 5.30 µm at about 90% of the way across the tree ring, with a slight reduction at the final part. However, in 2023, a slight decrease in wall thickness was observed at 80% of the growth ring (Figure 5C). For C. fortunei, the lumen



FIGURE 2

Number of cambium (A-C), enlarging (D-F), wall thickening (G-I), and mature (J-L) cells of C. lanceolata (blue) and C. fortunei (red) during 2021-2023 at the Gushan Mountain. Error bars indicate the standard deviation between trees



Cambial phenology of *C. lanceolata* and *C. fortunei* during the 2021–2023 reported as the timing of onset and ending of xylem cell division, and the ending of the growing season. Different letters indicate significant differences between years (p<0.05). Error bars indicate the standard deviation.

diameter was largest at 20% of the growth ring across all three years (35.87, 39.57, 36.32 μ m, respectively), after which it consistently decreased (2.44, 2.75, 2.91 μ m, respectively). However, by 2023, cells had expanded at 70%–80% of the tree ring (Figure 5B). The cell wall thickness ranged from 1.73 μ m to 4.81 μ m. In 2023, thickness initially increased, followed by a temporary decrease at 80% of the growth ring before rising again (Figure 5D).

3.5 Kinetics of cell development

In 2023, heavy late-summer rainfall promoted the latewood cell enlargement rate (r_E), which doubled, thus forming L-IADFs. For *C. lanceolata*, r_E remained consistently high during earlywood development (averaging 6.42, 4.03, and 6.06 µm d⁻¹), while the cell wall thickening rate (r_{WT}) gradually increased (averaging 0.33, 0.26, and 0.48 µm d⁻¹). During latewood formation, r_E declined sharply as r_{WT} accelerated, peaking at 90% of the growth ring before tapering off. Notably, in 2023, an abrupt rise in r_E was observed at 80% of the growth ring (7.44 µm d⁻¹), resulting in a larger lumen diameter and relatively thinner cell wall compared to other latewood cells (Figures 6A, C). For *C. fortunei*, 20% of the growth

ring exhibited the highest r_E (7.18, 6.27, and 7.49 µm d⁻¹, respectively), which then declined sharply. During latewood formation, r_E typically decreased gradually. However, a significant increase in r_E was observed between 70% and 80% of the tree ring in 2023. The rate increased by 125% (compared to the average of the other two years), reaching 6.57 µm d⁻¹. This resulted in cells with enlarged lumens and thinner walls, resembling earlywood cells. The r_{WT} steadily increased from 10% to 90% of the tree ring, followed by a decrease in the final part. However, in 2023, r_{WT} decelerated at 80%–90% of the tree ring, while it continued to rise in the final part (Figures 6B, D).

4 Discussion

4.1 Growth dynamics under different precipitation regimes

The bimodal growth pattern, characterized by two distinct phases of xylem cell production, has been reported in both Mediterranean and humid subtropical regions (Zheng et al., 2021; Huang et al., 2018; Camarero et al., 2010). This pattern is primarily



shading indicates a drier period.

driven by seasonal fluctuations in moisture availability, particularly during spring and autumn (Jiang et al., 2021; Garcia-Forner et al., 2019). In our study, both C. lanceolata and C. fortunei exhibited distinct growth peaks in spring and autumn, with growth stagnating during the dry summer months (Figure 2). This bimodal growth

pattern aligns with the typical rainfall regime of humid subtropical China, suggesting that bimodal xylem production is a characteristic feature of conifer species in this region.

Inter-annual variability in precipitation from 2021 to 2023 significantly influenced xylem division during the late growing



during 2021-2023. Thin lines indicate earlywood cells, dotted lines indicate IADFs, and thick lines indicate latewood cells. Lines show the mean values and shaded areas show one standard deviation. EW, earlywood cell; LW, latewood.

season. In 2023, reduced precipitation towards the end of the growing season led to an earlier cessation of cell division (Figure 3). However, abundant late-summer rainfall in the same year triggered a more pronounced autumn growth peak. In contrast, the lack of late-summer precipitation in 2022 resulted in early cessation of xylem growth, eliminating the autumn growth peak (Figure 4B). These inter-annual variations highlight the adaptability of trees to changing environmental conditions. Specifically, under drought stress, trees tend to adopt a conservative growth strategy to maintain physiological stability, while under wetter conditions, they accelerate growth to maximize resource allocation (Tumajer et al., 2021). In three years of





The relative position across the growth ring for the rate of cell enlarging (A, B) and the rate of cell wall thickening (C, D) of C. lanceolata and C. fortunei during 2021-2023. Thin lines indicate earlywood cells, dotted lines indicate IADFs, and thick lines indicate latewood cells. EW, earlywood cell: LW. latewood

observations, we found no significant inter-annual difference in the onset of xylem cell division, suggesting that the previous year's conditions had little direct impact on the current year's xylem phenology (Figure 3). Nevertheless, even without pronounced phenological shifts, the severe drought in 2022 could have indirectly influenced growth in 2023 by depleting nutrient reserves and reducing cambial activity (Zlobin, 2024).

These growth dynamics underscore the critical role of water availability in regulating xylem production. During drought conditions, cambial activity is often downregulated, limiting cell division and elongation due to reduced turgor pressure and decreased carbohydrate allocation to the cambium (Wang et al., 2024). Conversely, under wetter conditions, increased water availability stimulates cambial activity, promoting more extensive cell division and elongation, which leads to the autumn growth peak (Campelo et al., 2021; De Micco et al., 2016). The timing and intensity of rainfall are thus pivotal in determining the amplitude of the autumn growth peak, highlighting the sensitivity of xylem formation to moisture variability.

4.2 Impacts of extreme climate on xylem structure

Extreme climatic events, such as severe droughts and heavy rainfall, can induce plastic adjustments in wood formation dynamics, leading to changes in xylem structure (Zlobin, 2022; Fonti et al., 2010). Variability in precipitation directly affects rates of cell enlargement and wall thickening, which ultimately shape xylem morphology (Silvestro et al., 2023; Cuny et al., 2014). These structural modifications reflect the trees' adaptive responses to environmental stress. However, prolonged droughts or intense rainfall can disrupt typical growth patterns and result in structural anomalies in the xylem (Niccoli et al., 2024; Zhang et al., 2020).

In 2022, a severe drought from August to October reduced soil moisture and increased vapor pressure deficit (VPD), which severely restricted cambial activity (Table 2; Du et al., 2024; Salomón et al., 2022; He et al., 2024). Consequently, both C. lanceolata and C. fortunei exhibited a unimodal growth pattern, with xylem production limited to the early growing season. This drought-induced cessation of cell division prevented the formation of an autumn growth peak and led to significantly reduced xylem production (Figure 2). In contrast, the substantial rainfall events in late July and early September 2023 reactivated cambial activity and accelerated cell division. The subsequent increase in water availability during latewood formation facilitated cell enlargement, leading to the development of L-IADFs. L-IADFs, characterized by large lumen diameters and thin cell walls, are crucial for enhancing water storage and improving xylem water transport efficiency under humid conditions (Niccoli et al., 2024; Bing et al., 2022).

L-IADFs formation is a key adaptive response that optimizes water transport under fluctuating moisture conditions. By enlarging the xylem lumina, trees increase hydraulic conductivity, thereby reducing resistance to water flow and ensuring efficient water transport. This structural adaptation is essential for sustaining metabolic functions during periods of high humidity and enables trees to cope effectively with moisture variability. However, cellular structure is influenced not only by variations in moisture but also by the integrated interplay of plant physiological mechanisms. For instance, water potential, cell turgor, and plant hormones (e.g., auxin and cytokinins) work together to regulate xylem cell expansion and wall thickening. Ren et al. (2015) demonstrated that reduced soil water potential diminishes cell turgor, thereby inhibiting radial expansion in vascular tissues. Buttò et al. (2020) showed that exogenous application of cytokinins under moderate drought conditions can partially restore cambial activity, highlighting the intricate hormonal interplay under water stress. Meanwhile, Niccoli et al. (2024) found that rapid rehydration following a drought can trigger compensatory growth spurts, culminating in the formation of anomalous density bands (L-IADFs). Consistent with these findings, the sustained rainfall during late summer and autumn of 2023 in our study not only reactivated cambial division but also provided the conditions necessary for accelerated cell enlargement, ultimately producing L-IADFs.

4.3 Bimodal growth dynamics do not guarantee the occurrence of L-IADFs

While both 2021 and 2023 exhibited bimodal growth patterns, L-IADFs were not observed in 2021 (Figure 5; Supplementary Figure S1), which contrasts with findings from previous studies (Niccoli et al., 2024; Campelo et al., 2021; Liu et al., 2018). This discrepancy suggests that bimodal growth dynamics do not automatically result in L-IADFs formation, as these two phenomena are not directly correlated. Timely water replenishment during late summer and early autumn can reactivate cambial activity, promoting xylem cell division and leading to a second growth peak (Camarero et al., 2010). However, although cambial activity resumed in 2021, the rate of cell enlargement was insufficient to promote L-IADFs formation, indicating that the presence of an autumn growth peak does not always lead to L-IADFs.

In contrast, the abundant and sustained precipitation during late summer and autumn in 2023 not only stimulated cambial cell division but also significantly accelerated xylem cell enlargement, resulting in the formation of L-IADFs (Figure 4; Figure 5). These results emphasize that a bimodal growth pattern alone is insufficient to predict the formation of L-IADFs. Rather, the combination of sufficient moisture availability and the rate of cell expansion is crucial for L-IADFs formation. Therefore, while bimodal growth dynamics indicate cambial reactivation, the structural modifications necessary for L-IADFs formation depend on both precise water replenishment and the tree's physiological capacity for rapid cell enlargement. Nevertheless, our observations are limited to just three years. Consequently, pinpointing a definitive water threshold for L-IADFs formation remains challenging. Future work involving longer-term observations and controlled experiments-coupled with real-time monitoring of physiological indicators-may help clarify whether such a threshold exists. This approach would not only deepen our understanding of L-IADFs formation but also offer empirical support for more robust modeling efforts.

Understanding these adaptive mechanisms is crucial for predicting the long-term impacts of climate change on forest ecosystems. Prolonged droughts or unpredictable rainfall patterns can disrupt traditional growth cycles, leading to structural anomalies in xylem formation. These anomalies may result in increased fluctuations in wood density, which can negatively affect xylem hydraulic efficiency and carbon allocation within trees-critical processes for forest health and productivity. Frequent occurrences of IADFs may indicate instability in wood formation driven by environmental stress, which could signal a decrease in forest ecosystem resilience. This instability reflects the diminished capacity of trees to maintain stable growth patterns and adapt to fluctuating environmental conditions (Fang and Zhang, 2024; Battipaglia et al., 2010). Future research should explore the physiological thresholds beyond which forest species are unable to recover from drought stress or tolerate extreme climatic events. Insights into these thresholds will be vital for developing forest management strategies that enhance the resilience of subtropical forests under the pressures of global climate change.

5 Conclusions

This study highlights the crucial role of late-summer precipitation in shaping the growth dynamics and xylem structure of Cunninghamia lanceolata and Cryptomeria fortunei in humid subtropical China. Our findings demonstrate that sufficient late-summer rainfall promotes a bimodal growth pattern, whereas the autumn drought of 2022 shortened the growing season by approximately two months, eliminating the secondary growth peak and resulting in an unusual unimodal growth pattern. In contrast, continuous late-summer rainfall in 2023 accelerated the enlargement rate of latewood cells, leading to the formation of L-IADFs, characterized by large lumens and thin cell walls. Importantly, our results show that the occurrence of a secondary growth peak does not necessarily lead to the formation of L-IADFs, as observed in 2021. Extreme climatic events, including prolonged droughts and heavy rainfall, can disrupt typical growth patterns and induce structural anomalies in xylem formation. These disruptions highlight the sensitivity of trees to moisture variability, which can impair hydraulic efficiency and affect overall growth stability. Understanding the complex responses of tree growth and xylem structure to fluctuating precipitation regimes is essential for predicting forest dynamics and developing strategies to protect forest health. As climate models forecast increasing precipitation variability, future research should focus on the long-term effects of extreme climate events on wood formation and forest growth. Insights from such research will be critical for developing adaptive forest management strategies aimed at enhancing the resilience of subtropical forests to the impacts of global climate change.

Data availability statement

The original contributions presented in the study are included in the article/Supplementary Material. Further inquiries can be directed to the corresponding authors.

Author contributions

CW: Conceptualization, Data curation, Formal Analysis, Investigation, Writing – original draft. ZZ: Data curation, Investigation, Supervision, Writing – review & editing. JG: Funding acquisition, Supervision, Writing – review & editing. FZ: Funding acquisition, Software, Writing – review & editing. SR: Supervision, Writing – review & editing. KF: Conceptualization, Funding acquisition, Project administration, Writing – review & editing, Supervision.

Funding

The author(s) declare that financial support was received for the research and/or publication of this article. This work was funded by the National Science Foundation of China (42301058, 42425101 and 42130507), the Basic Scientific Research Projects of Fujian Provincial Public Welfare Scientific Research Institutes (2024R1002004). CW received the State Scholarship Funding provided by the China Scholarship Council (202408350075) to conduct this research.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Generative AI statement

The author(s) declare that no Generative AI was used in the creation of this manuscript.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: https://www.frontiersin.org/articles/10.3389/fpls.2025.1568882/ full#supplementary-material

SUPPLEMENTARY FIGURE 1

Wood anatomy in tree-rings of *Cu. lanceolata* (A-C) and *Cr. fortunei* (D-F) sampled at the Gushan Mountain during 2021 (A, D), 2022 (B, E) and 2023 (C, F) in humid subtropical China. Arrows indicate the L-IADF.

References

Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., et al. (2010). A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *For. Ecol. Manage.* 259, 660–684. doi: 10.1016/j.foreco.2009.09.001

Balzano, A., Battipaglia, G., and De Micco, V. (2019). Wood-trait analysis to understand climatic factors triggering intra-annual density-fluctuations in co-occurring Mediterranean trees. *IAWA J.* 40, 241–258. doi: 10.1163/22941932-40190220

Battipaglia, G., De Micco, V., Brand, W. A., Linke, P., Aronne, G., Saurer, M., et al. (2010). Variations of vessel diameter and $\delta^{13}C$ in false rings of *Arbutus unedo* L. reflect different environmental conditions. *New Phytol.* 188, 1099–1112. doi: 10.1111/j.1469-8137.2010.03443.x

Bing, X. J., Fang, K. Y., Gong, X. Y., Wang, W. Z., Xu., C. X., Li, M. H., et al. (2022). The intra-annual intrinsic water use efficiency dynamics based on an improved model. *Clim. Change* 172, 16. doi: 10.1007/s10584-022-03368-1

Buttò, V., Deslauriers, A., Rossi, S., Rozenberg, P., Shishov, V., and Morin, H. (2020). The role of plant hormones in tree-ring formation. *Trees* 34, 315–335. doi: 10.1007/ s00468-019-01940-4

Camarero, J. J., Olano, J. M., and Parras, A. (2010). Plastic bimodal xylogenesis in conifers from continental Mediterranean climates. *New Phytol.* 185, 471–480. doi: 10.1111/j.1469-8137.2009.03073.x

Campelo, F., Ribas, M., and Gutiérrez, E. (2021). Plastic bimodal growth in a Mediterranean mixed-forest of Quercus ilex and Pinus halepensis. *Dendrochronologia* 67, 125836. doi: 10.1016/j.dendro.2021.125836

Cuny, H. E., Rathgeber, C. B. K., Frank, D., Fonti, P., and Fournier, M. (2014). Kinetics of tracheid development explain conifer tree-ring structure. *New Phytol.* 203, 1231–1241. doi: 10.1111/nph.12871

De Micco, V., Campelo, F., De Luis, M., Bräuning, A., Grabner, M., Battipaglia, G., et al. (2016). Intra-annual density fluctuations in tree rings: how, when, where, and why? *IAWA J.* 37, 232–259. doi: 10.1163/22941932-20160132

Denne, M. P. (1989). Definition of latewood according to mork, (1928). *IAWA J.* 10, 59–62. doi: 10.1163/22941932-90001112

Du, M. C., Xu, C., Wang, A., Lv, P. C., Xu, Z. Q., and Zhang, X. L. (2024). Different drought recovery strategy between Larix spp. and *Quercus mongolica* in temperate forests. *Sci. Total Environ.* 938, 173521. doi: 10.1016/j.scitotenv.2024.173521

Fang, O. Y., and Zhang, Q. B. (2024). Stress triggers tree-growth rebound in global forests. *Agric. For. Meteorol.* 359, 110285. doi: 10.1016/j.agrformet.2024.110285

Feldman, A. F., Feng, X., Felton, A. J., Konings, A. G., Knapp, A. K., Biederman, J. A., et al. (2024). Plant responses to changing rainfall frequency and intensity. *Nat. Rev. Earth Environ.* 5, 276–294. doi: 10.1038/s43017-024-00534-0

Feng, X., Porporato, A., and Rodriguez-Iturbe, I. (2013). Changes in rainfall seasonality in the tropics. *Nat. Clim. Change* 3, 811–815. doi: 10.1038/nclimate1907

Fonti, P., Von Arx, G., García-González, I., Eilmann, B., Sass-Klaassen, U., Gärtner, H., et al. (2010). Studying global change through investigation of the plastic responses of xylem anatomy in tree rings. *New Phytol.* 185, 42–53. doi: 10.1111/j.1469-8137.2009.03030.x

Gao, J. N., Rossi, S., and Yang, B. (2021a). Origin of intra-annual density fluctuations in a semi-arid area of northwestern China. *Front. Plant Sci.* 12. doi: 10.3389/ fpls.2021.777753

Gao, J. N., Yang, B., Peng, X. M., and Rossi, S. (2021b). Tracheid development under a drought event producing intra-annual density fluctuations in the semi-arid China. *Agric. For. Meteorol.* 308–309, 108572. doi: 10.1016/j.agrformet.2021.108572

Garcia-Forner, N., Vieira, J., Nabais, C., Carvalho, A., Martínez-Vilalta, J., and Campelo, F. (2019). Climatic and physiological regulation of the bimodal xylem formation pattern in Pinus pinaster saplings. *Tree Physiol.* 39, 2008–2018. doi: 10.1093/treephys/tpz099

Häusser, M., Aryal, S., Barth, J. A. C., Bendix, J., Garel, E., van Geldern, R., et al. (2023). Xylem formation patterns from Mediterranean to subalpine climate conditions reveal high growth plasticity of pine species on Corsica. *Trees* 37, 1027–1039. doi: 10.1007/s00468-023-02402-8

He, C., Collins, M., Zhou, T. J., Jiang, X. W., Wu, P. L., and Dunstone, N. (2024). Contrasting East Asian climate extremes in 2020 and 2022 tied to zonal flow. *Environ. Res. Lett.* 19, 104019. doi: 10.1088/1748-9326/ad6a72

Huang, J. G., Guo, X. L., Rossi, S., Zhai, L. H., Yu, B. Y., Zhang, S. K., et al. (2018). Intra-annual wood formation of subtropical Chinese red pine shows better growth in dry season than wet season. *Tree Physiol.* 38, 1225–1236. doi: 10.1093/treephys/tpy046

Huang, J. G., Zhang, Y. L., Wang, M. H., Yu, X. H., Deslauriers, A., Fonti, P., et al. (2023). A critical thermal transition driving spring phenology of Northern Hemisphere conifers. *Glob. Chang Biol.* 29, 1606–1617. doi: 10.1111/gcb.16543

Iles, C. E., Samset, B. H., Sandstad, M., Schuhen, N., Wilcox, L. J., and Lund, M. T. (2024). Strong regional trends in extreme weather over the next two decades under high- and low-emissions pathways. *Nat. Geosci.* 17, 845–850. doi: 10.1038/s41561-024-01511-4

IPCC (2021). Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press).

Jiang, Y. Y., Zhang, X. Q., Chhin, S., and Zhang, J. G. (2021). A bimodal pattern and age-related growth of intra-annual wood cell development of chinese fir in subtropical China. *Front. Plant Sci.* 12. doi: 10.3389/fpls.2021.757438

Li, X. X., Liang, E. Y., Gričar, J., Rossi, S., Čufar, K., and Ellison, A. M. (2017). Critical minimum temperature limits xylogenesis and maintains treelines on the southeastern Tibetan Plateau. *Sci. Bull.* 62, 804–812. doi: 10.1016/j.scib.2017.04.025

Liu, X. S., Nie, Y. Q., and Wen, F. (2018). Seasonal dynamics of stem radial increment of pinus Taiwanensis hayata and its response to environmental factors in the lushan mountains, Southeastern China. *Forests* 9, 387. doi: 10.3390/f9070387

Ma, Z., Guo, Q., Yang, F., Chen, H., Li, W., Lin, L., et al. (2021). Recent changes in temperature and precipitation of the summer and autumn seasons over Fujian province, China[J]. *Water.* 13 (14), 1900. doi: 10.3390/w13141900

Niccoli, F., Kabala, J. P., Pacheco-Solana, A., and Battipaglia, G. (2024). Impact of intraannual wood density fluctuation on tree hydraulic function: insights from a continuous monitoring approach. *Tree Physiol.* 44, tpad145. doi: 10.1093/treephys/tpad145

Palombo, C., Fonti, P., Lasserre, B., Cherubini, P., Marchetti, M., and Tognetti, R. (2018). Xylogenesis of compression and opposite wood in mountain pine at a Mediterranean treeline. *Ann. For. Sci.* 75, 93. doi: 10.1007/s13595-018-0773-z

Rathgeber, C. B. K., Cuny, H. E., and Fonti, P. (2016). Biological basis of tree-ring formation: A crash course. *Front. Plant Sci.* 7. doi: 10.3389/fpls.2016.00734

Ren, P., Rossi, S., Gricar, J., Liang, E. Y., and Cufar, K. (2015). Is precipitation a trigger for the onset of xylogenesis in Juniperus przewalskii on the north-eastern Tibetan Plateau? *Ann. Bot.* 115, 629–639. doi: 10.1093/aob/mcu259

Rossi, S., Anfodillo, T., and Menardi, R. (2006a). Trephor: A New Tool for Sampling Microcores from tree stems. *IAWA J.* 27, 89–97. doi: 10.1163/22941932-90000139

Rossi, S., Deslauriers, A., and Anfodillo, T. (2006b). Assessment of cambial activity and xylogenesis by microsampling tree species: an example at the alpine timberline. *IAWA J.* 27, 383–394. doi: 10.1163/22941932-90000161

Rossi, S., Deslauriers, A., and Morin, H. (2003). Application of the Gompertz equation for the study of xylem cell development. *Dendrochronologia* 21, 33–39. doi: 10.1078/1125-7865-00034

Rossi, S., Anfodillo, T., Čufar, K., Cuny, H. E., Deslauriers, A., Fonti, P., et al. (2016). Pattern of xylem phenology in conifers of cold ecosystems at the Northern Hemisphere [J]. *Global Change Biology*. 22 (11), 3804–3813. doi: 10.1111/gcb.13317

Salomón, R. L., Peters, R. L., Zweifel, R., Sass-Klaassen, Ute, G. W., Stegehuis, A. I., et al. (2022). The 2018 European heatwave led to stem dehydration but not to consistent growth reductions in forests. *Nat. Commun.* 13, 28. doi: 10.1038/s41467-021-27579-9

Silvestro, R., Zeng, Q., Buttò, V., Sylvain, J. D., Drolet, G., Mencuccini, M., et al. (2023). A longer wood growing season does not lead to higher carbon sequestration. *Sci. Rep.* 13, 4059. doi: 10.1038/s41598-023-31336-x

Tumajer, J., Shishov, V. V., Ilyin, V. A., and Camarero, J. J. (2021). Intra-annual growth dynamics of Mediterranean pines and junipers determines their climatic adaptability. *Agric. For. Meteorol.* 311, 108685. doi: 10.1016/j.agrformet.2021.108685

Vieira, J., Moura, M., Nabais, C., Freitas, H., and Campelo, F. (2019). Seasonal adjustment of primary and secondary growth in maritime pine under simulated climatic changes. *Ann. For. Sci.* 76, 84. doi: 10.1007/s13595-019-0865-4

Wang, S. J., Deng, Y., Gao, L. L., Zhang, Y. H., Shi, X. Y., and Gou, X. H. (2023). Influence of monsoon anomalies on intra-annual density fluctuations of Chinese pine in the Loess Plateau. *Int. J. Biometeorol* 67, 847–856. doi: 10.1007/s00484-023-02459-7

Wang, F., Zhang, J. Z., Fonti, P., Sun, Q. P., Wang, Y. T., Zhang, F., et al. (2024). Effects of site aridity and species on stand transpiration in high-elevation dryland ecosystems. *J. Hydrol.* 630, 130662. doi: 10.1016/j.jhydrol.2024.130662

Wang, C. S., Zheng, Z. P., Zhou, F. F., Liu, X. S., Fonti, P., Gao, J. N., et al. (2023). Intraannual dynamic of opposite and compression wood formation of Pinus massoniana Lamb. in humid subtropical China. *Front. For. Glob. Change* 6. doi: 10.3389/ffgc.2023.1224838

Xu, T., Sun, X. S., Hong, H., Wang, X. Y., Cui, M. Y., Lei, G. L., et al. (2019). Stable isotope ratios of typhoon rains in Fuzhou, Southeast China, during 2013–2017. J. Hydrol. 570, 445–453. doi: 10.1016/j.jhydrol.2019.01.017

Zalloni, E., De Luis, M., Campelo, F., Novak, K., De Micco, V., Di Filippo, A., et al. (2016). Climatic signals from intra-annual density fluctuation frequency in mediterranean pines at a regional scale. *Front. Plant Sci.* 7. doi: 10.3389/fpls.2016.00579

Zhang, J. Z., Alexander, M. R., Gou, X. H., Deslauriers, A., Fonti, P., Zhang, F., et al. (2020). Extended xylogenesis and stem biomass production in Juniperus przewalskii Kom. during extreme late-season climatic events. *Ann. For. Sci.* 77, 99. doi: 10.1007/s13595-020-01008-1

Zheng, Z. P., Zhou, F. F., Fonti, P., Ren, P., Li, X. X., Miao, G. F., et al. (2021). Intra-Annual Wood Formation of Cryptomeria fortunei and Cunninghamia lanceolata in Humid Subtropical China. Front. Ecol. Evol. 9. doi: 10.3389/ fevo.2021.733974

Zhou, F. F., Dong, Z. P., Fang, K. Y., Chen, D. L., Tang, H., Ou, T. H., et al. (2024). Summer heat induced the decline of Pinus Taiwanensis forests at its southern limit in humid Subtropical China. *Agric. For. Meteorol.* 350, 109974. doi: 10.1016/j.agrformet.2024.109974

Zlobin, I. E. (2022). Linking the growth patterns of coniferous species with their performance under climate aridization. *Sci. Total Environ.* 831, 154971. doi: 10.1016/j.scitotenv.2022.154971

Zlobin, I. E. (2024). Tree post-drought recovery: scenarios, regulatory mechanisms and ways to improve. *Biol. Rev.* 99, 1595–1612. doi: 10.1111/brv.13083