



#### **OPEN ACCESS**

EDITED BY Costanza Colombi, Stella Maris Foundation (IRCCS), Italy

REVIEWED BY
B. Blair Braden,
Arizona State University, United States
Samantha Harker,
Arizona State University, United States,
in collaboration with reviewer BB
Arianna Bentenuto,
University of Trento, Italy

\*CORRESPONDENCE
Aleksandra Grzeszak

☑ aleksandra.grzeszak@uw.edu.pl

RECEIVED 18 February 2025 ACCEPTED 19 June 2025 PUBLISHED 18 July 2025 CORRECTED 14 August 2025

#### CITATION

Grzeszak A and Pisula E (2025) Experiences of females on the autism spectrum through the perspective of minority stress theory: a review. Front. Psychiatry 16:1578963. doi: 10.3389/fpsyt.2025.1578963

#### COPYRIGHT

© 2025 Grzeszak and Pisula. This is an openaccess article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

# Experiences of females on the autism spectrum through the perspective of minority stress theory: a review

Aleksandra Grzeszak\* and Ewa Pisula

Department of Health and Rehabilitation Psychology, Faculty of Psychology, University of Warsaw, Warsaw, Poland

**Introduction:** People on the autism spectrum, especially females, are at high risk of co-occurring psychiatric and psychological conditions, suicidal ideation, and psychiatric hospitalization. The minority stress theory offers a framework for understanding mental health disparities by focusing on the influence of social factors. The current review aims to synthesize the literature on the experiences of autistic females in terms of the unique, chronic and socially based stressors that emerge at the intersection of sex/gender and autism.

**Methods:** We conducted a literature search in a systematic way on the Web of Science and Scopus databases, applying the specified inclusion and exclusion criteria. We chose a narrative manner of data presentation.

**Results:** Based on the evidence from 104 studies of various methodologies, we show that the burdens encountered by autistic females are more than the sum of strains common for the females and for autistic individuals. Females on the autism spectrum face also specific challenges related to their not meeting social gendered expectations and the stereotypical image of autism.

**Discussion:** This intersectional disadvantage can enhance our understanding of the adverse mental health outcomes observed in this population. We emphasize key directions and provide methodological recommendations for future research in this area. Additionally, we underscore the importance of enhancing practitioners' understanding of the unique characteristics of female autism and the specific vulnerabilities faced by this group to improve support and outcomes.

#### KEYWORDS

autism, females, life experiences, mental health, minority stress, social experiences, stigma

#### 1 Introduction

Autism spectrum disorder (ASD) is defined as a complex neurodevelopmental condition, characterized by lifelong challenges in social communication and interpersonal interactions, as well as repetitive, rigid patterns of behavior and interests (1). Since the identification of autism is based only on behavioral indicators, its conceptualization has evolved with the accumulation of data. Diagnostic criteria have been broadened, including different patterns of traits grouped under the term autism spectrum (2, 3). This has contributed to an increase in the frequency of diagnoses of ASD, amongst others in female individuals (3).

The male-to-female ratio in the autistic population is commonly reported as 3-4:1 (4). Increasingly, it is recognized that this asymmetry may be due to the underdiagnosis of girls and women, rather than an accurate reflection of the number of females on the autism spectrum (3–5). Nonetheless, females remain a minority within the diagnosed population. Consequently, the experiences of autistic females have been the subject of minimal research until recently, when the literature on this topic has begun to expand considerably. At this stage, there is a need to review prior research and chart directions for future scientific inquiry.

#### 1.1 The minority stress and autism

The minority stress model (6) is an elaboration of the social stress theory, which links mental health disparities with disadvantaged social status. Social stressors are understood as stigma-related experiences encountered by socially marginalized groups, compounded by a deficiency of coping resources (7). In following sections, we provide the rationale behind the application of this theory in analyzing the data on autistic females, and the overview of the model.

#### 1.1.1 The applicability of the social stress theory

The premise of the social stress theory is the existence of health disparities between disadvantaged and advantaged groups (7). Being on the autism spectrum is linked with poorer mental (8) as well as physical health outcomes compared to the general population (9). These health inequities contribute to the observed premature mortality in the autistic population due to chronic illnesses, medication side effects, or suicide (10, 11). Numerous studies have demonstrated a heightened risk of co-occurring anxiety disorders, depressive disorders, suicidal tendencies, and psychiatric hospitalization among females on the autism spectrum (12–17). Additionally, evidence suggests that autistic females experience poorer physical health (18).

The neurodiversity movement originated in autistic self-advocacy groups and is based on the assumption of equality and appreciation of differences in the human brain functioning (19). It promotes the notion that autistic individuals are an integral part of society and that autism should be understood not solely as a medical condition but also as an essential component of personal identity (19, 20). The neurodiversity movement draws attention to the social

situation of people on the spectrum – employing the term "neurominority". This term refers to a part of the population that shares not only challenges related to the way their nervous system works, but also experiences of social marginalization (21). Consequently, employing theories that analyze stigma-related processes may provide a deeper understanding of autistic individuals' life experiences and their effects on mental health (19, 21–23).

The stigmatization process begins with the labeling of a person and attribution of negative stereotypes to the label (24). Examples of unfavorable stereotypes applied to autistic individuals include "weird", "loner", "violent", or "retarded" (25–28). Positive associations, such as high intelligence or exceptional skills in specific domains, also exist (27, 29). In addition to the label itself, some behaviors that are common for autism but contrary to neurotypical social norms (e.g., inadequate facial expressions or tone of speaking, stimming, poor eye contact, not following the rules of conversation, etc.) are stigmatized. Some research indicates that stigma towards these behaviors is even greater than stigma towards the label of autism itself (30, 31).

Gender-related factors have a major influence on the formation of social structures and institutions, and prejudices and stereotypes play important roles in this process (32). The theory of intersectionality posits that disadvantaged identities do not simply combine additively, but they interact dynamically and reciprocally, producing a specific constellation of experiences (7, 33). While primarily used to analyze the experiences of Black females (33), the intersectionality framework enhances our understanding of challenges faced by females on the autism spectrum (34), including Black autistic females (35).

#### 1.1.2 The minority stress model - an overview

The minority stress model was primarily used to study sexual and gender minorities (6), but has been shown applicable in disability studies (36) and autism research (21). This theoretical framework states that minorities face unique and persistent challenges that are rooted in social hierarchies and go beyond general life stressors (6).

Minority stressors range from external (distal) to internal (proximal) in relation to the individual. The most distal are the actual events of discrimination and/or victimization. These events can manifest as individual acts of discrimination or violence in interpersonal relations, such as social exclusion or bullying, which are commonly experienced by autistic persons (37, 38). Additionally, there are more permanent situations in relation to institutions that are reported by individuals on the autism spectrum. These include instances of lack of necessary accommodations in the workplace or educational setting (39–41), as well as unmet healthcare needs (42, 43).

Living in a stigmatizing environment can affect individuals even in the absence of a concrete event of discrimination or victimization (6, 44). Expectation of rejection, a more proximal stressor, involves fear, insecurity, and vigilance that negatively impact self-concept and mental health (6, 45). It has been shown that autistic people anticipate negative reactions of others (21, 25, 46).

Concealing one's minority identity is yet another, even more proximal stressor (6). Given that certain minority identities, such as autism, are not immediately discernible, individuals may employ concealment as a means of safeguarding themselves from violence and discrimination. Concealment of autism can be understood as hiding the fact of diagnosis as well as behaving in such a way as to pass as non-autistic (22, 47-51). The latter phenomenon is known as masking or camouflaging, and is widely recognized as more prominent in females than in males on the autism spectrum, with the influence of gender stereotypes and higher social motivation considered among the possible explanations for the difference (5, 48-50, 52) Masking of autistic traits is one of the factors that may account for under-diagnosis of females (4, 5). The broad topic of camouflaging and its consequences has been elaborated in many works and reviews (5, 22, 47, 49-51) and is beyond the scope of this paper. These two strategies of stigma management are often utilized jointly; however, disclosing one's diagnostic status does not necessarily lead to abandoning all masking behaviors (22, 53). Besides, most of those who disclose do so selectively and the question of if and to whom to disclose the diagnosis is a psychologically tasking dilemma (25, 44).

Internalized stigma is conceptualized as the most proximal stressor (6). Due to awareness of stigma and exposure to prejudice-related events, the individual incorporates into their self some negative judgments that are harmful to their self-concept and self-esteem (6, 54). While individuals on the autism spectrum exhibited lower levels of internalized stigma compared to patients with serious mental illness (55, 56), studies specifically designed to investigate this phenomenon among autistic individuals have demonstrated that negative perceptions about autism influence the way they think about themselves (25, 26, 44).

The minority stress model assumes that the influence of stressors on mental health outcomes is modified by minority identity – its prominence, valence, and integration with other identities, as well as coping and social support resources arising from the identity (6). As indicated by Botha (57) in her qualitative study aiming to create the first measure of autistic community connectedness, autistic identity enables one to be part of a community, and this community reciprocally helps to develop a positive identity. Cooper, Cooper, Russell, and Smith (58) found that positive valuation of autism is positively linked to higher collective self-esteem in individuals who identify with the autistic community.

Botha and Frost (21) were the first to use the social stress framework to study potential causes of poor mental health and wellbeing outcomes in the autistic population. Utilizing adapted scales of minority stress, they demonstrated that despite controlling for general stress exposure, minority stress significantly predicted poorer well-being and higher psychological distress in individuals on the autism spectrum. Their study provides preliminary results that may indicate a relationship between minority stress and gender, but the data did not allow for further exploration (21).

#### 1.2 Aim of the review

In the late 2000s, research into the topic of sex/gender differences in autism presentation saw a notable increase (59, 60) and, starting with the important work of Cridland et al. in 2013 (61), the area of research on the life experiences of autistic females has grown substantially. The purpose of this review is to elucidate findings on the experiences of females on the autism spectrum, focusing on unique, chronic, and socially determined stressors resulting from the intersection of sex/gender and autism. We chose the narrative review approach, as it enables the extraction of information from a variety of studies not necessarily connected directly to minority stress theory and the synthesis thereof in the context of the theory (62).

#### 2 Method

Despite choosing the narrative manner of data presentation, we approached the literature search in a systematic way. The first author conducted the search on the Web of Science and Scopus databases using the queries containing combinations of alternative terms referring to: (1) autism, (2) sex/gender differences, (3) experiences, stigma, and minority stressors, linked with Boolean operators. A detailed presentation of the keywords used is shown in Table 1.


The literature search was conducted on 12-13 February 2024 and was narrowed to original research articles published in English in peer-reviewed journals since 2013. Additional search was undertaken on 19 November 2024, aimed at supplementing the query with works published in the meantime. We included studies involving females with formal autism diagnoses as well as self-diagnosed females. We decided to incorporate studies researching children, adolescent, and adult populations because the minority status of autistic females appears early in their lives and affects identity, mental health and well-being since then. We included articles describing quantitative, qualitative, and mixed methods studies. The records were pre-screened for duplicates, then the first author screened titles and abstracts. The exclusion criteria at this stage were: (1) research without female autistic participants or

TABLE 1 Search terms used in the review.

Sex/gender	Autism	Experiences, stigma, minority stressors	
women OR females OR girls OR "sex difference*" OR "gender difference*" OR "with ASD" OR "with ASD" OR "with Asperger's" OR neurodivergent OR autis* OR Asperger*		experience* OR stigma* OR label* OR stereotype* OR prejudice* OR rejection* OR discrimination OR inequit* OR inequalit* OR sexis* OR victimization OR violence OR bullying OR concealment OR disclosure OR identity	

with the number or female participants not allowing to check for sex differences; (2) studies in which autistic women were an indistinguishable part of a wider sample; (3) case studies; (4) research exclusively on the symptoms or autistic traits presentation; (5) research focusing on parents' or caregivers' experiences; (6) research testing psychometric tools; (7) research solely on the prevalence of mental and physical health problems. The last criterion stems from the assumptions of minority stress theory. This theory applies to explain disparities in mental health

related to social factors (that is, social stressors) rather than all differences. Since the purpose of our study was to review research on those experiences of women on the autism spectrum that can be considered minority stressors, we only included studies that examined the relationship between mental health and experiences. The prevalence of mental health problems is the subject of other reviews (8, 16). We decided to exclude also the studies focused on camouflaging of autistic traits. The PRISMA flow chart illustrates the reasons for exclusion of retrieved records for



which the full text has been assessed (Figure 1). Any doubts about the inclusion of articles were resolved collaboratively by us. In the end, a total of 104 articles were included in the review.

## 3 Autistic females' social experiences and mental health

Two methodological approaches are commonly used to examine minority stress theory. Between-group analyses focus on examining the effect of social status on mental health by comparing the prevalence of psychiatric symptoms or disorders in minority and non-minority groups. Within-group analyses investigate the

relations between stress processes (e.g., levels of exposure to stigmarelated experiences or coping resources) and mental health outcomes (7). Only six quantitative studies incorporated in our review included mental health measures as one of the variables. A summary of the results of these studies is presented in Table 2.

# 3.1 Between-group comparisons: autistic vs non-autistic females, autistic females vs autistic males

Three studies compared the occurrence of psychiatric diagnoses or internalizing symptoms between autistic and non-autistic

TABLE 2 Quantitative studies that included mental health measures as one of the variables (n = 6).

Authors and year	Country	Sample	Topic – type of experience	Mental health measure	Relevant findings about mental health
Cazalis, Reyes, Leduc, & Gourion (2022) (68)	France	women of the French autistic community ( $n = 225$ ), age > 18	Sexual victimization	Question about co- occurrence of psychiatric disorders Question about reported consequences of the assault in the next 6 months	Participants who have encountered sexual violence demonstrated higher prevalence of mental health problems, particularly PTSD
DaWalt et al. (2020) (67)	United States	high school students with educational classification of autism, ( <i>n</i> = 547; 76 females, 471 males), age 14-21; parents' reports	Social participation	Question about current mental health diagnoses	Females demonstrated higher prevalence of internalizing disorders (the variable of presence of mental health diagnoses was not included in further analyses on social participation rates)
Greenlee, Winter, & Marcovici (2020) (66)	United States	adolescents with autism diagnosis ( <i>n</i> = 105; 50 females, 55 males), age 13-17, and their primary caregivers	Peer victimization	Revised Children's Anxiety and Depression Scale (RCADS) – Short Version, subscales Anxiety and Depression	Girls reported more symptoms of anxiety and depression than boys Relational violence predicted anxiety symptoms in girls, but not in boys
Hampton, Allison, Baron-Cohen & Holt (2023) (63)	United Kingdom	384 autistic and 490 non-autistic mothers, age > 18	Childbirth and postnatal experiences	Question about being diagnosed with postnatal depression Question about being diagnosed with postnatal anxiety	Postnatal depression and anxiety were more prevalent in autistic than in non-autistic mothers (items on postnatal depression and disorders were not included in further multivariate analyses)
O'Connor, van den Bedem, Blijd-Hoogewys, Stockmann, & Rieffe, (2022) (65)	Netherlands	pre-adolescents and adolescents: autistic (n = 104; 18 girls, 84 boys) and non-autistic (n = 202; 111 girls, 91 boys), age 9-16	Friendship	Child Depression Inventory Child Symptom Inventory (CSI) – generalized anxiety disorder subscale	Autistic participants reported more depression and anxiety symptoms, no effect of gender In all groups depression negatively correlated with positive friendship quality (PFQ) and positively with negative friendship quality For the whole sample, anxiety correlated negatively with PFQ, but in sub-sample of autistic females anxiety and PFQ correlated positively
Pohl et al. (2020) (64)	Participants from different countries (mostly Western)	355 autistic and 132 non-autistic mothers	Pregnancy, childbirth, and motherhood	Question about having additional psychiatric or psychological diagnosis Question about experiencing prenatal depression Question about experiencing postnatal depression	Additional psychiatric or psychological diagnosis were more prevalent in autistic mothers More autistic than non-autistic mothers reported experiencing prenatal and postnatal depression

groups. In two studies on perinatal and motherhood experiences, mothers on the autism spectrum demonstrated higher prevalence of perinatal depression (63, 64) and postnatal anxiety (63) than non-autistic mothers. Moreover, other psychiatric problems were found to be more prevalent in autistic than non-autistic mothers (64). In these two studies, mothers on the autism spectrum also reported more negative experiences with professional care. There were no within-groups analyses testing whether more negative experiences with perinatal care providers were related to more adverse mental health outcomes (63, 64). In research on friendship quality, autistic pre-adolescents and adolescents exhibited more symptoms of depression and anxiety than their non-autistic peers (65).

Three studies explored mental health differences between males and females on the autism spectrum. While O'Connor et al. (65) found no gender differences in rates of internalizing symptoms, Greenlee, Winter, and Marcovici (66) identified higher levels of depression and anxiety among autistic girls experiencing peer victimization compared to autistic boys. Another study reported that female adolescents were more likely than males to have diagnoses of internalizing disorders according to parental reports (67).

The studies mentioned above add to the body of research that indicates that females on the autism spectrum may be at greater risk of psychiatric disorders, compared to non-autistic females (63–65) and autistic males (66, 67). However, the only study incorporating both gender and neurotype variables found significant differences between neurotypes but no gender differences in rates of depression and anxiety symptoms (65). Of note, this study involved adolescents diagnosed by clinicians in accordance with formal criteria and only 18 girls.

### 3.2 Within-group analyses: social experiences and mental health

Three studies examined the relationships between social experiences and mental health. Sexual victimization heightened the number of co-occurring mental health problems - precisely posttraumatic stress disorder (PTSD) was more prevalent in autistic females who had survived sexual assault in comparison to those who had not have such experience (68). Bullying, specifically relational violence, predicted symptoms of anxiety in girls, but not in boys on the autism spectrum (66). Depression and anxiety correlated negatively with positive friendship experiences and positively with negative friendship experiences in autistic and non-autistic (pre-)adolescents. Only autistic girls had a positive correlation between anxiety and positive friendship experiences (65). The results of these studies suggest that experiencing victimization is associated with poorer mental health (66, 68). Therefore, the relationship between socially-based experiences and mental health outcomes is marked. Interestingly, even if there are no differences in rates of these experiences between females and males on the autism spectrum, there are differences in relations between those experiences and mental health symptoms. Autistic females may be more prone than autistic males to detrimental effects of relational violence (66). Additionally, in this group, even engagement in positive social interactions may be paid for with more immense anxiety (65). Although only the study of O'Connor et al. (65) included both between-group and within-group analyses, all findings presented above support the potential value of analyzing the nature and the extent of social stressors faced by females on the autism spectrum. In the next part of this work, we will outline a landscape of possible relations between autism, sex/gender, social environment, and wellbeing and mental health, based on the currently available data. The structure of the following text mirrors Meyer's minority stress processes, starting from events external to the individual, gradually approaching identity-related experiences and ending with experiences that may modify the impact of minority stressors on mental health.

#### 4 Discrimination

#### 4.1 Diagnosis and support services

Females obtain autism diagnosis later than males (52, 69-71), even if they do not differ from males in levels of autistic characteristics (71). There are many females, who were not diagnosed until adulthood and they believe that this delay had prevented them from accessing adequate support and harmed their well-being (29, 72-82). Late-diagnosed autistic females describe the road to diagnosis as long and winding due to clinicians' stereotypical, male-centric view of autism and their dismissive attitude towards women seeking diagnosis (29, 73, 75, 76, 78, 80, 81, 83-87). Moreover, being primarily diagnosed with other condition is another factor that hinders autism diagnosis in females (29, 70, 75, 77, 78, 82, 84, 85). Also, more intensive camouflaging, observed to a greater extent in females than in males, was associated with a later time to diagnosis (52, 71), and this relationship may be stronger in women than in men, although further research on larger samples is needed (71). In qualitative studies, autistic females indicate that this was one of the barriers to both referral for diagnosis (29, 85), and during the diagnostic process if providers did not create conditions for feeling safe enough to "unmask" (80, 87). Females also report facing greater barriers to accessing post-diagnosis support services than males (88) and frequently evaluate these services as inadequate or insufficient (79, 80, 89, 90).

#### 4.2 Education

Lack of adequate support in educational settings is reported retrospectively by females diagnosed later in life (76, 80, 86, 91) but also by students with formal diagnosis (92, 93). They cite teachers' lack of understanding, pressure on academic achievements and attendance, and insufficient accommodations as significant obstacles to reaching their full academic potential (80, 86, 91–93). Autistic females reported not receiving adequate support not only in learning, but – even more frequently – in emotional and social functioning (72, 86) and the percentage of females dissatisfied with the level of support in their educational experiences was significantly higher than in the male sample (72). Sometimes, an unaccommodating mainstream school environment leads to exclusion from school to alternative settings

which provide a more individualized learning environment. However, such settings often host many students with emotional, behavioral, and social difficulties, making them potentially challenging for vulnerable autistic girls (93, 94).

#### 4.3 Employment

Research suggests that females and males on the autism spectrum do not differ significantly in rates of employment (95-97) or in average hourly wages (95). Although the evidence of sex/gender differences in average weekly work hours in autistic adults population are ambiguous (95, 96), females tend to prefer part-time work (72, 96) or even withdrawal from the job market, a choice not necessarily linked with parenting responsibilities (96). More pronounced differences are noted between autistic and nonautistic females, with autistic females less likely than the general female population to have paid employment (72). Autistic females are also more likely to experience job instability, underemployment, difficulties in gaining and maintaining jobs, and negative work experiences compared to non-autistic females, but there were no significant differences between autistic females and males (97, 98). These negative employment experiences can therefore be considered common to all people on the spectrum (97).

Differences between males and females on the autism spectrum in vocational experiences may be subtle. For instance, co-occurring anxiety and depression, although less common in males, were found to be a greater barrier to employment in this group (95). In turn, autistic females report a stressful mismatch of their social skills and preferences with stereotypical gendered expectations formulated by coworkers and superiors (72, 99–101).

#### 4.4 Healthcare

To date, no significant gender differences have been found in the rates of unmet medical services among autistic adults (88). However, healthcare needs among females on the autism spectrum are more likely to go unmet compared to non-autistic females. The findings concerning perinatal, and perimenopausal care indicate that professionals' lack of knowledge about autism and dismissive or even stigmatizing attitudes towards autistic individuals translate into lower quality of care, poorer treatment outcomes, and higher stress and anxiety (64, 75, 102–108). These barriers are also faced by females on the autism spectrum who experience eating disorders, hindering their recovery (109, 110).

#### 5 Victimization

Research on victimization examines various forms of violence and uses different methodologies. The data from register-based studies indicate that females with autism diagnosis are at increased risk of child maltreatment (111), and serious physical violence in adolescence and adulthood (112) compared to non-autistic

individuals and males with autism diagnosis. Self-report studies show that being a victim of physical violence is most prevalent among autistic females compared to both non-autistic females and autistic males (113, 114). In contrast, other self-report study suggests the importance of diagnosis rather than sex/gender in the experience of various forms of victimization (115). The two most researched forms of victimization encountered by people on the spectrum are bullying and sexual violence.

#### 5.1 Bullying

Autistic status, but not sex/gender, was found to be associated with higher rates of experiencing recurrent and intentional social aggression in educational and work settings (72, 97, 100, 115-119). While the overall rates of peer victimization do not appear to differ between females and males on the autistic spectrum, there are some nuances. Autistic males may be rejected and victimized more blatantly, while peer aggression between females may consist of less pronounced behaviors such as ignoring or exclusion (66, 120, 121). In their narratives, autistic females describe various form of peer violence: physical (86, 93, 94, 122), verbal (61, 77, 84, 86, 90, 93, 94, 100, 123), and relational (29, 61, 74, 81, 85, 91, 122, 124–126). They point out their divergence from social norms, especially gender norms, as causes of experienced bullying (29, 73, 81, 86, 101, 122, 124, 126, 127). Some indicate other girls and women as main enforcers of these norms and the most frequent perpetrators of peer victimization (29, 61, 84, 100, 124, 128). Seeing one's own transgressions of social expectations as the cause of violence may result in greater helplessness and self-blame in relationships and therefore affect well-being and future social interactions (29, 66, 73, 86, 121).

#### 5.2 Sexual violence

Quantitative research on sexual victimization face substantive and methodological challenges, such as definitions of sexual violence, measurement tools, and reaching participants. Majority of studies demonstrate that female sex is linked with increased vulnerability to sexual victimization and being autistic female further enhances the risk (113, 114, 129, 130). When no significant differences between females and males on the autism spectrum are observed, it is because the rate of sexual violence reported by autistic males is strikingly high (115, 131). Females on the autism spectrum encounter sexual abuse in adulthood but also below the age of consent, and the frequency of experiencing sexual violence reaches 90% (68). The perpetrators are reported to be both strange men or/and those they were familiar with (29, 68, 80, 122, 132). Autistic females face sexual harassment also in the online world (133). Homosexual orientation may be an additional risk factor for autistic females' adverse sexual experiences (134). In their narratives, females on the autism spectrum identify their low social skills, gullibility, lack of assertiveness, inadequate body language, and a strong desire to be accepted as factors leading them to be vulnerable to sexual victimization (29, 73, 80, 106, 122, 132). The

role of sexual knowledge is ambiguous — low levels of sexual knowledge may increase the risk of engaging in unwanted sexual interactions (131), while having sexual knowledge and the desire to engage in sexual relationships may expose autistic females to abuse, such as exploitation by a sexual partner (129). Being a victim of a sexual assault is linked with PTSD and with re-victimization (68).

#### 6 Expectation of rejection

Qualitative analyses of social experiences reveal traces of autistic females' beliefs that others may reject them. The rejection feared by females on the autism spectrum is twofold. First, they report the anxiety of being discriminated or victimized due to autism stigma. Some feel that negative stereotypes towards autism would affect the way other people interact with them (64, 74, 76, 84, 90, 91, 100, 107, 135-137). Others describe the vigilance related to anticipated negative reactions to their behaviors that differ from social norms (83, 121, 123, 125, 128, 136, 138, 139). Second, autistic females fear the invalidation of their autistic identity and experiences because autism is stereotypically viewed as a male condition (77, 78, 87, 102, 106, 137, 140). The fear of being repelled, mistreated or disbelieved is present in relations with friends (84, 91, 121, 125, 128), partners (132), employers and coworkers (74, 76, 100, 138), healthcare professionals (64, 77, 103, 106, 107, 137), and diagnostic and support services providers (77, 78, 87, 102, 140). In a quantitative study on barriers to formal autism diagnosis, concerns about not being believed, listened to, or being accused of "making up" symptoms were the most prominent barriers, regardless of gender. However, females (and gender-diverse individuals) rated these concerns as more severe than males did (141).

#### 7 Concealment and disclosure

For autistic females, concealing their autism is a strategy for "social survival". They do it in order to avoid anticipated stigmatization (29, 74, 83, 90, 91, 94, 142) and to make the establishment of positive social relations more likely (29, 73, 78, 81, 94, 100, 128). The decision to disclose one's autistic identity may be comparable to coming out (91). Autistic females disclose their diagnosis to close connections in hopes of building more authentic relationships (74, 76, 78, 83, 84, 91, 132). In formal settings, disclosure is motivated by the need for understanding and necessary accommodations, such as those in schools, workplaces, or healthcare environments (74, 83, 90, 100, 102, 103, 107, 137, 142).

When speaking about the actual experiences of disclosure, autistic females point out both positive and negative consequences. Some of them describe growth in personal relations (78, 83, 91, 132). Other report getting adjustments at school, work or in healthcare facilities (83, 90, 100, 103, 137). Disclosing may give freedom of self-expression and help reduce masking (78, 83, 91). Conversely, sharing the truth is often met with disbelief or dismissal (63, 64, 77, 81, 84, 100, 107, 137, 143), and

sometimes with rejection and discrimination because of negative stereotypes concerning autism (90, 100, 102, 107, 132). Negative outness experiences may influence future decisions concerning disclosure (78, 90).

#### 8 Internalized stigma

Autistic females report internalized negative judgments related to three main areas where they feel they fail to meet societal expectations. First, they describe self-blame and feelings of being "wrong," "broken," "weird," or generally inferior due to behaviors that deviate from neurotypical norms (73–75, 78, 80, 81, 86, 91, 127, 132, 138). Some report that they had believed their autistic traits rendered them incapable of fulfilling social roles, such as those of a partner (132), parent (102), or employee (100, 138). Some present self-blame about their negative social experiences, including victimization (73, 78, 80, 86, 122). Second, they express feelings of failing to meet expectations of stereotypical femininity (76, 122). Third, they describe how their divergence from male-centered stereotypes of autism contributes to feelings of being "fussy," "impostors," or undeserving of support and accommodation services (76, 77, 87, 89).

#### 9 Identity and connectedness

Autistic identity can be viewed from a personal as well as a social perspective (144–146). Following Meyer's conceptualization, we consider autistic identity as a personal identification with the diagnostic label of the autism spectrum, which can be more or less personally important, positive, and integrated with other aspects of one's self-concept (6). The degree to which an individual identifies with a group of people on the autism spectrum we refer to as community connectedness. It includes a sense of belonging and shared experiences, as well as behavioral involvement in activities to benefit the community (53, 57, 147). As shown below, these phenomena are intertwined.

#### 9.1 Autistic identity

Receiving an autism diagnosis enables autistic females to renounce guilt and shame and develop more positive identity (73, 75–77, 79–84, 90, 91, 102, 106, 127, 132, 135, 148). This process often involves rejecting a medical, deficit-centered view of autism and embracing its positive aspects, such as uniqueness, good memory, creativity, empathy, and passion for certain topics (80, 81, 83, 91, 135). An additional aspect may be the opportunity to push back against the adverse impacts of failed gender expectations on personal identity (29, 75, 83, 149). Furthermore, feeling validated in their way of functioning can reduce autistic camouflaging, enabling them to replace a sense of false identity with a more authentic self (73, 128). While females on the autism spectrum underline the importance of autism in their self-image (81, 83), they

also point out the way in which it is integrated with other aspects of their personalities, creating complex identities (80, 124, 142).

#### 9.2 Community connectedness

Positive peer relationships are vital but challenging for autistic individuals, and loneliness is a common experience (65, 67, 86, 121, 125-128, 139, 150-153). Female peer connections often involve complex social rules, which may create a particularly harmful gap between the social skills of autistic females and the demands of social interactions within their gender group (65, 67, 86, 120, 121, 152, 153). Links between autistic identity and social connectedness are two-way. Positive autistic identity enables one to connect with autistic community (73, 75-77, 102, 106). In turn, connectedness with other autistic individuals enhances appraisal of one's autistic traits (29, 73-78, 102, 106, 138, 151). It also offers a sense of belonging (29, 73, 75, 77, 78, 80, 81, 84, 102, 135, 136, 142, 143), less effortful social interactions (75, 81, 135, 143), and informational and emotional support (81, 84, 102, 136, 137, 140, 143). Having an authentic and relevant self-image also facilitates connections with neurotypical individuals, improving social relationships and increasing access to social support (77, 86, 106, 135, 138).

The works included in the review from which it was possible to draw information on identity and community connectedness are qualitative studies and do not include cross-gender comparisons. A review of quantitative studies on autistic identity by Davies et al. (145) indicated sparse and inconclusive results on the relationship between autistic identity and sex/gender. Thus, based on the data collected in our review, we can only emphasize that the diagnostic delay in females on the spectrum hinders the development of a positive identity and community connectedness. Some papers included in our review involved samples composed of both females with a formal autism diagnosis and self-diagnosed. Among the studies discussed in this section, only three included female participants without a formal diagnosis. Two of these studies analyzed material posted on blogs and forums, in which verification of diagnosis status would have been impossible anyway (77, 84). The study of Harmens et al. (76) combined self-identified and pending diagnosis participants into one group. The remaining studies, conducted using interviews, required a formal diagnosis. As a result, it is impossible to know whether there are any differences in the experience of autistic identity and community connectivity depending on diagnostic status. At the same time, this indicates the importance of formal diagnosis as validation of females' autistic selves.

#### 10 Discussion

This review aimed to synthesize the social experiences of females on the autism spectrum using the minority stress model as a conceptual framework. Basing on relatable and methodologically sound studies including samples of autistic

females, we can conclude that autistic females are disadvantaged compared to two populations within which they are minorities: neurotypical females and autistic males. This disadvantage is of intersectional nature. The social stress encountered by females on the autism spectrum is not a simple sum of burdens common for autistic people and females in general. Females on the autism spectrum also face unique strains arising from the particular content of the stigma towards autistic females: not feminine enough and not properly autistic. This stigma, combined with sexism and broader autism-related stigma, translates into lived experiences of discrimination and victimization, leading to expectations of rejection, concealment and selective disclosure, and internalized stigma. Moreover, these chronic social stressors hinder access to coping resources, such as necessary adjustments and social support.

Minority stress of females on the autism spectrum overlaps with other strains common for females. Pregnancy, childbirth, and menopause are physically and mentally difficult transitional periods requiring medical assistance. These are often the situations that expose autistic females to minority stressors such as discrimination or expectation of rejection, and dilemmas about disclosure.

In summary, the experiences of autistic females go beyond their individual characteristics and life events, and are rooted in societal notions of normality, autism, and femininity. Therefore, the minority stress theory provides a useful lens through which to examine the impact of these experiences on mental health.

Most of the studies included in this review were based on relatively small samples of women on the autism spectrum, which limits the extent to which their findings can be generalized to the broader autistic female population. An additional challenge lies in the complexities of diagnosing autism in women. Access to formal diagnosis is often restricted, leading many individuals to self-identify as being on the spectrum. This self-identified group—those without a formal diagnosis—is frequently underrepresented in research. Furthermore, girls and women who do receive a diagnosis may exhibit a more "male-typical" pattern of autistic traits, as existing diagnostic tools are largely designed around such presentations. Consequently, the picture that emerges from the reviewed literature is inevitably incomplete, particularly with regard to the diversity of the autistic population and variations in diagnostic status.

The shared experiences of minority stress among women on the autism spectrum, along with the associated mental health challenges, may significantly shape the clinical presentation of autism in this population. As highlighted in the review, autistic women are frequently exposed to similar social and cultural pressures, which influence their functioning and may contribute to the development of a distinct female autism phenotype. Research on minority stress in this group offers valuable insights into the mechanisms underlying the masking of autistic traits. The social pressure to conform to normative expectations often compels these individuals to conceal their neurodivergent characteristics—efforts that can carry a substantial cost to their psychological well-being. This dynamic creates a complex interplay in which attempts to

avoid stigma and rejection may, paradoxically, lead to negative mental health outcomes.

#### 10.1 Directions for future research

Although we retrieved 104 relevant articles concerning social experiences of autistic females, only six explored the relationships between those experiences and mental health. Therefore, we pinpoint this gap as an important direction of future research. Moreover, cross-sectional designs of these studies do not allow for conclusions on causality. The relations can be bidirectional, for instance, higher levels of anxiety may increase one's vulnerability to violence (66). Longitudinal studies are needed to capture the directions of the impact. Identifying factors that mitigate negative mental health outcomes would be of particular practical and clinical value. These factors could be, for instance, positive autistic identity and social connectedness (53, 57, 154).

Almost all studies included in this review recruited participants without intellectual disability. Thus, the conclusions drawn in this work can only be generalized to this part of the autistic female population. It is important to design studies enabling to explore social experiences of autistic females with intellectual disabilities and the impact of these experiences on mental health and wellbeing in this group. Using adequate prompts and accessible research tools is a good practice allowing participants with intellectual disability to share their experiences (122).

Sampling presents another challenge in research on females on the autism spectrum. Some studies recruit only formally diagnosed participants, potentially overlooking the experiences of those undiagnosed, a significant portion of the autistic female population (1). To overcome this limitation, other studies recruit participants who identify as autistic, regardless of formal diagnosis. This approach, however, carries the risk of inaccuracies in self-diagnosis. As in both cases researchers rely mostly on self-declaration of one's diagnostic or identity status, it is important to use psychometric measures to confirm the higher rate of autistic traits. Including the variable of diagnostic status in analyses allows for the identification of potential differences in the experiences of diagnosed and undiagnosed participants.

Most studies included in this review were conducted in Western countries, mainly the USA, UK, and Western Europe. As societal perceptions of the norm (also gender norms) and stigmatizing attitudes are deeply rooted in culture (24, 32), studies on social stress of autistic females conducted in other cultures would allow to identify differences and similarities in this regard (138).

Minority stress affects the autistic population as a whole, regardless of sex or gender (21). Although findings on sex/gender differences in mental health indicate higher prevalence of internalizing symptoms in autistic females than males, they are not unanimous. In addition, the issue of underdiagnosis of depression in males due to its atypical presentation and predominance of externalizing symptoms is becoming more

widely recognized (155, 156). As such, the hypothesis that autistic females are more disadvantaged than autistic males warrants testing using tools that examine minority stress variables and account for both internalizing and externalizing disorders.

Lastly, the above-average prevalence of transgender identity among autistic individuals (157–159) highlights the need to differentiate experiences. Reviewed studies adopted different sex/gender terminology and defined the samples studied in varying ways. For this reason, we have standardized the terminology used, without referring to the aspect of gender identity. It should be noted that gender diversity is more often reported by females on the autism spectrum (157–159). Moreover, some of the experiences of autistic females and autistic gender-diverse people may overlap (122, 141). However, the latter group faces unique minority stress (160), of which the manifestations and effects should be explored separately.

#### 10.2 Practical implications

Knowledge of risk and protective factors for autistic females' mental health and well-being is of great importance in clinical practice. It can be used to fine-tune diagnostic and support services, as well as educational and workplace environments in order to counterbalance the challenges met by girls and women on the autism spectrum.

We propose the following measures to improve the situation of females on the autism spectrum:

- 1. Including information on female presentation of autistic characteristics and camouflaging autism in the curricula of psychological, medical, and pedagogical studies in order to counteract the bias toward autism as a male condition (161–163).
- Creating accessible and reliable information resources for parents on the characteristics of autism in girls, also taking into account the issues of difficulties in adjusting to gender expectations (164, 165).
- 3. Using adequate diagnostic tools and in-depth information from diverse sources in the assessment of female individuals, including methods that evaluate camouflaging (166, 167).
- 4. Establishing post-diagnostic strength-based support programs aimed at increasing knowledge, self-awareness, and positive identity, as well as assisting in the decisions about disclosure. Such interventions can take the form of peer support groups, which can increase social connectedness (168, 169).

Many of autistic females' experiences of discrimination were encountered in contact with professionals in clinical and educational settings. As the diagnosis of autism spectrum disorder is a key code for autistic females to access resources to cope with detrimental effects of social stressors, the elimination of barriers to

receiving it early is a critical issue. Changing the way society perceives females on the autism spectrum is a long-term task; however, raising awareness among researchers and practitioners about the specificity of the female autism experience is within reach.

#### Data availability statement

The original contributions presented in the study are included in the article/Supplementary Material. Further inquiries can be directed to the corresponding author.

#### **Author contributions**

AG: Conceptualization, Funding acquisition, Writing – original draft, Writing – review & editing. EP: Conceptualization, Methodology, Supervision, Writing – review & editing.

#### **Funding**

The author(s) declare that financial support was received for the research and/or publication of this article. This work was supported by the Faculty of Psychology, University of Warsaw, from the funds awarded by the Ministry of Science and Higher Education in the form of a subsidy for the maintenance and development of research potential in 2024 (501-D125-01-1250000, zlec. 5011000193). The funds covered the cost of language proofreading of the article. The Article Processing Fee was subsidized in the internal grant system within the programme of the Ministry of Science and Higher Education "Initiative of Excellence - Research University" at the University of Warsaw, Actions I.2.4 (PSP 501-D125-20-0001241) and IV.4.1 (PSP 501-D125-20-0004410).

#### Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

#### Correction note

A correction has been made to this article. Details can be found at: 10.3389/fpsyt.2025.1672758.

#### Generative Al statement

The author(s) declare that no Generative AI was used in the creation of this manuscript.

#### Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

#### Supplementary material

The Supplementary Material for this article can be found online at: https://www.frontiersin.org/articles/10.3389/fpsyt.2025.1578963/full#supplementary-material

#### References

- 1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders (5th ed., Text Revision). Washington, DC: American Psychiatric Publishing (2022).
- 2. Lombardo MV, Lai MC, Baron-Cohen S. Big data approaches to decomposing heterogeneity across the autism spectrum. *Mol Psychiatry*. (2019) 24:1435–50. doi: 10.1038/s41380-018-0321-0
- 3. Russell G, Stapley S, Newlove-Delgado T, Salmon A, White R, Warren F, et al. Time trends in autism diagnosis over 20 years: a UK population-based cohort study. *J Child Psychol Psychiatry Allied Discip.* (2022) 63:674–82. doi: 10.1111/jcpp.13505
- 4. Loomes R, Hull L, Mandy WPL. What is the male-to-female ratio in autism spectrum disorder? A systematic review and meta-analysis. *J Am Acad Child Adolesc Psychiatry*. (2017) 56:466–74. doi: 10.1016/j.jaac.2017.03.013
- 5. Tubío-Fungueiriño M, Cruz S, Sampaio A, Carracedo A, Fernández-Prieto M. Social camouflaging in females with autism spectrum disorder: A systematic review. *J Autism Dev Disord*. (2021) 51:2190–9. doi: 10.1007/s10803-020-04695-x
- Meyer IH. Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: Conceptual issues and research evidence. *Psychol Bull.* (2003) 129:674–97. doi: 10.1037/0033-2909.129.5.674
- 7. Schwartz S, Meyer IH. Mental health disparities research: The impact of within and between group analyses on tests of social stress hypotheses. *Soc Sci Med.* (2010) 70:1111–8. doi: 10.1016/j.socscimed.2009.11.032
- 8. Lai MC, Kassee C, Besney R, Bonato S, Hull L, Mandy W, et al. Prevalence of cooccurring mental health diagnoses in the autism population: a systematic review and metaanalysis. *Lancet Psychiatry*. (2019) 6:819–29. doi: 10.1016/S2215-0366(19)30289-5

- 9. Cashin A, Buckley T, Trollor JN, Lennox N. A scoping review of what is known of the physical health of adults with autism spectrum disorder. *J Intellect Disabil.* (2018) 22:96–108. doi: 10.1177/1744629516665242
- 10. Jokiranta-Olkoniemi E, Gyllenberg D, Sucksdorff D, Suominen A, Kronström K, Chudal R, et al. Risk for premature mortality and intentional self-harm in autism spectrum disorders. *J Autism Dev Disord.* (2021) 51:3098–108. doi: 10.1007/s10803-020-04768-x
- 11. Smith DaWalt L, Hong J, Greenberg JS, Mailick MR. Mortality in individuals with autism spectrum disorder: Predictors over a 20-year period. *Autism.* (2019) 23:1732–9. doi: 10.1177/1362361319827412
- 12. Angell AM, Deavenport-Saman A, Yin L, Zou B, Bai C, Varma D, et al. Sex differences in co-occurring conditions among autistic children and youth in Florida: A retrospective cohort study (2012–2019). *J Autism Dev Disord*. (2021) 51:3759–65. doi: 10.1007/s10803-020-04841-5
- 13. Gu Z, Dawson G, Engelhard M. Sex differences in the age of childhood autism diagnosis and the impact of co-occurring conditions. *Autism Res.* (2023) 16:2391–402.
- 14. McGillivray JA, Evert HT. Exploring the effect of gender and age on stress and emotional distress in adults with autism spectrum disorder. *Focus Autism Other Dev Disabl.* (2018) 33:55–64. doi: 10.1177/1088357614549317
- 15. Oswald TM, Winter-Messiers MA, Gibson B, Schmidt AM, Herr CM, Solomon M. Sex differences in internalizing problems during adolescence in autism spectrum disorder. *J Autism Dev Disord.* (2016) 46:624–36. doi: 10.1007/s10803-015-2608-1

- 16. Sedgewick F, Leppanen J, Tchanturia K. Gender differences in mental health prevalence in autism. *Adv Autism.* (2020) 7:208–24. doi: 10.1108/AIA-01-2020-0007
- 17. So P, Wierdsma AI, Van Boeijen C, Vermeiren RRJM, Mulder NCL. Gender differences between adolescents with autism in emergency psychiatry. *Autism.* (2021) 25:2331–40. doi: 10.1177/13623613211019855
- 18. Kassee C, Babinski S, Tint A, Lunsky Y, Brown HK, Ameis SH, et al. Physical health of autistic girls and women: a scoping review. *Mol Autism*. (2020) 11. doi: 10.1186/s13229-020-00380-z
- 19. Kapp SK, Gillespie-Lynch K, Sherman LE, Hutman T. Deficit, difference, or both? Autism and neurodiversity. *Dev Psychol.* (2013) 49:59–71. doi: 10.1037/a0028353
- 20. Bagatell N. From cure to community: Transforming notions of autism. *Ethos.* (2010) 38:33–55. doi: 10.1111/j.1548-1352.2009.01080.x
- 21. Botha M, Frost DM. Extending the minority stress model to understand mental health problems experienced by the autistic population. *Soc Ment Health.* (2020) 10:20–34. doi: 10.1177/2156869318804297
- 22. Perry E, Mandy W, Hull L, Cage E. Understanding camouflaging as a response to autism-related stigma: A social identity theory approach. *J Autism Dev Disord*. (2021) 52:800–10. doi: 10.31234/osf.io/7w2pe
- 23. Rivera RA, Bennetto L. Applications of identity-based theories to understand the impact of stigma and camouflaging on mental health outcomes for autistic people. *Front Psychiatry.* (2023) 14:1–12. doi: 10.3389/fpsyt.2023.1243657
- 24. Link BG, Phelan JC. Conceptualizing stigma. Annu Rev Sociol. (2001) 27:363–85. doi:  $10.1146/\mathrm{annurev.soc.}$ 27.1.363
- 25. Botha M, Dibb B, Frost DM. Autism is me": an investigation of how autistic individuals make sense of autism and stigma. *Disabil Soc.* (2022) 37:427–53. doi: 10.1080/09687599.2020.1822782
- 26. Humphrey N, Lewis S. Make me normal": The views and experiences of pupils on the autistic spectrum in mainstream secondary schools. Autism. (2008) 12:23–46. doi: 10.1177/1362361307085267
- 27. Wood C, Freeth M. Students' Stereotypes of autism. *J Educ Issues*. (2016) 2:131–40. doi: 10.5296/jei.v2i2.9975
- 28. Treweek C, Wood C, Martin J, Freeth M. Autistic people's perspectives on stereotypes: An interpretative phenomenological analysis. *Autism.* (2019) 23:759–69. doi: 10.1177/1362361318778286
- 29. Bargiela S, Steward R, Mandy W. The experiences of late-diagnosed women with autism spectrum conditions: an investigation of the female autism phenotype. *J Autism Dev Disord*. (2016) 46:3281–94. doi: 10.1007/s10803-016-2872-8
- 30. Gillespie-Lynch K, Daou N, Obeid R, Reardon S, Khan S, Goldknopf EJ. What contributes to stigma towards autistic university students and students with other diagnoses? *J Autism Dev Disord*. (2021) 51:459–75. doi: 10.1007/s10803-020-04556-7
- 31. Butler RC, Gillis JM. The impact of labels and behaviors on the stigmatization of adults with asperger's disorder. *J Autism Dev Disord*. (2011) 41:741-9. doi: 10.1007/s10803-010-1093-9
- 32. Glick P, Fiske ST. An ambivalent alliance: hostile and benevolent sexism as complementary justifications for gender inequality. *Am Psychol.* (2001) 56:109–18.
- 33. Crenshaw K. Demarginalizing the intersection of race and sex: A black feminist critique of antidiscrimination doctrine, feminist theory, and antiracist politics. *Univ Chic Leg Forum.* (1989) 139:39–52.
- 34. Fox J. The impact of intersectional disadvantage on autistic women's experiences of interpersonal violence: A narrative review. *Autism Adulthood*. (2024) 00:1–12. doi: 10.1089/aut.2023.0100
- 35. Diemer MC, Gerstein ED, Regester A. Autism presentation in female and Black populations: Examining the roles of identity, theory, and systemic inequalities. *Autism.* (2022) 26:1931–46. doi: 10.1177/13623613221113501
- 36. Lund EM. Examining the potential applicability of the minority stress model for explaining suicidality in individuals with disabilities. *Rehabil Psychol.* (2021) 66:183–91. doi: 10.1037/rep0000378
- 37. Humphrey N, Hebron J, Humphrey N, Hebron J. Bullying of children and adolescents with autism spectrum conditions: a 'state of the field 'review conditions: a 'state of the field 'review. *Int J Incl Educ.* (2015) 0:1–18. doi: 10.1080/13603116.2014.981602
- 38. Schroeder JH, Cappadocia MC, Bebko JM, Pepler DJ, Weiss JA. Shedding light on a pervasive problem: A review of research on bullying experiences among children with autism spectrum disorders. *J Autism Dev Disord*. (2014) 44:1520–34. doi: 10.1007/s10803-013-2011-8
- 39. Davies J, Heasman B, Livesey A, Walker A, Pellicano E, Remington A. Autistic adults' views and experiences of requesting and receiving workplace adjustments in the UK. *PloS One.* (2022) 17:5–15. doi: 10.1371/journal.pone.0272420
- 40. Goddard H, Cook A. I spent most of freshers in my room"—A qualitative study of the social experiences of university students on the autistic spectrum. *J Autism Dev Disord.* (2022) 52:2701–16. doi: 10.1007/s10803-021-05125-2
- 41. Hasson L, Keville S, Gallagher J, Onagbesan D, Ludlow K, Hasson L, et al. Inclusivity in education for autism spectrum disorders: Experiences of support from the and young people on the autism spectrum Inclusivity in education for autism spectrum disorders: Experiences of support from the perspective of parent/carers, scho. *Int J Dev Disabil.* (2024) 70:201–12. doi: 10.1080/20473869.2022.2070418

- 42. Nicolaidis C, Raymaker D, Mcdonald K, Dern S, Boisclair WC, Ashkenazy E, et al. Comparison of healthcare experiences in autistic and non-autistic adults: A cross-sectional online survey facilitated by an academic-community partnership. *J Gen Intern Med.* (2013) 28:761–9. doi: 10.1007/s11606-012-2262-7
- 43. Walsh C, Connor PO, Walsh E, Lydon S. A systematic review of interventions to improve healthcare experiences and access in autism. *Rev J Autism Dev Disord*. (2021) 10:185–202. doi: 10.1007/s40489-021-00279-2
- 44. Han E, Scior K, Avramides K, Crane L. A systematic review on autistic people's experiences of stigma and coping strategies. *Autism Res.* (2022) 15:12–26. doi: 10.1002/aur.2652
- 45. Link BG. Understanding labeling effects int the area of mental disorders: an assessment of the effects of expectations of rejection. *Am Sociol Rev.* (1987) 52:96–112. doi: 10.2307/2095395
- 46. Johnson TD, Joshi A. Dark clouds or silver linings? A stigma threat perspective on the implications of an autism diagnosis for workplace well-being. *J Appl Psychol.* (2016) 101:430–49. doi: 10.1037/apl0000058
- 47. Beck JS, Lundwall RA, Gabrielsen T, Cox JC, South M. Looking good but feeling bad: "Camouflaging" behaviors and mental health in women with autistic traits. *Autism.* (2020) 24:809–21. doi: 10.1177/1362361320912147
- 48. Hull L, Petrides KV, Mandy W. The female autism phenotype and camouflaging: a narrative review. *Rev J Autism Dev Disord*. (2020) 7:306–17. doi: 10.1007/s40489-020-00197-9
- 49. Lai MC, Lombardo MV, Ruigrok ANV, Chakrabarti B, Auyeung B, Szatmari P, et al. Quantifying and exploring camouflaging in men and women with autism. *Autism.* (2017) 21:690–702. doi: 10.1177/1362361316671012
- 50. Libsack EJ, Keenan EG, Freden CE, Mirmina J, Iskhakov N, Krishnathasan D, et al. A systematic review of passing as non-autistic in autism spectrum disorder. *Clin Child Family Psychol Rev.* (2021) 24:783–812. doi: 10.1007/s10567-021-00365-1
- 51. Pearson A, Rose K. A conceptual analysis of autistic masking: understanding the narrative of stigma and the illusion of choice.  $Autism\ Adulthood$ . (2021) 3:52–60. doi: 10.1089/aut.2020.0043
- 52. McQuaid GA, Lee NR, Wallace GL. Camouflaging in autism spectrum disorder: Examining the roles of sex, gender identity, and diagnostic timing. *Autism.* (2022) 26:552–9. doi: 10.1177/13623613211042131
- 53. Cage E, Cranney R, Botha M. Brief report: does autistic community connectedness moderate the relationship between masking and wellbeing? *Autism Adulthood.* (2022) 4:247–53. doi: 10.1089/aut.2021.0096
- 54. Corrigan PW, Rafacz J, Rüsch N. Examining a progressive model of self-stigma and its impact on people with serious mental illness. *Psychiatry Res.* (2011) 189:339–43. doi: 10.1016/j.psychres.2011.05.024
- 55. Dubreucq J, Plasse J, Gabayet F, Faraldo M, Blanc O, Chereau I, et al. Self-stigma in serious mental illness and autism spectrum disorder: Results from the REHABase national psychiatric rehabilitation cohort. *Eur Psychiatry*. (2020) 63. doi: 10.1192/j.eurpsy.2019.12
- 56. Bachmann CJ, Höfer J, Kamp-Becker I, Küpper C, Poustka L, Roepke S, et al. Internalised stigma in adults with autism: A German multi-center survey. *Psychiatry Res.* (2019) 276:94–9. doi: 10.1016/j.psychres.2019.04.023
- $\,$  57. Botha M. Autistic community connectedness as a buffer against the effects of minority stress.  $\it Univ Surrey.$  (2020).
- 58. Cooper R, Cooper K, Russell AJ, Smith LGE, Cooper K. I 'm Proud to be a Little Bit Different ": The Effects of Autistic Individuals 'Perceptions of Autism and Autism Social Identity on Their Collective Self esteem. *J Autism Dev Disord*. (2021) 51:704–14. doi: 10.1007/s10803-020-04575-4
- 59. Kirkovski M, Enticott PG, Fitzgerald PB. A review of the role of female gender in autism spectrum disorders. *J Autism Dev Disord*. (2013) 43:2584–603. doi: 10.1007/s10803-013-1811-1
- 60. Attwood T. *The complete guide to Asperger's syndrome*. London: Jessica Kingsley Publishers (2007).
- 61. Cridland EK, Jones SC, Caputi P, Magee CA. Being a girl in a boys' world: Investigating the experiences of girls with autism spectrum disorders during adolescence. *J Autism Dev Disord*. (2014) 44:1261–74. doi: 10.1007/s10803-013-1985.6
- 62. Baumeister RF, Leary MR. Writing narrative literature reviews. *Rev Gen Psychol.* (1997) 1:311–20. doi: 10.1037/1089-2680.1.3.311
- 63. Hampton S, Allison C, Baron-Cohen S, Holt R. Autistic people's perinatal experiences II: A survey of childbirth and postnatal experiences. *J Autism Dev Disord.* (2023) 53:2749–63. doi: 10.1007/s10803-022-05484-4
- 64. Pohl AL, Crockford SK, Blakemore M, Allison C, Baron-Cohen S. A comparative study of autistic and non-autistic women's experience of motherhood. *Mol Autism.* (2020) 11:3. doi: 10.1186/s13229-019-0304-2
- 65. O'Connor RAG, van den Bedem N, Blijd-Hoogewys EMA, Stockmann L, Rieffe C. Friendship quality among autistic and non-autistic (pre-) adolescents: Protective or risk factor for mental health? *Autism*. (2022) 26:2041–51. doi: 10.1177/13623613211073448
- 66. Greenlee JL, Winter MA, Marcovici IA. Brief report: gender differences in experiences of peer victimization among adolescents with autism spectrum disorder. *J Autism Dev Disord.* (2020) 50:3790–9. doi: 10.1007/s10803-020-04437-z

- 67. DaWalt LS, Taylor JL, Bishop S, Hall LJ, Steinbrenner JD, Kraemer B, et al. Sex differences in social participation of high school students with autism spectrum disorder. *Autism Res.* (2020) 13:2155–63. doi: 10.1002/aur.2348
- 68. Cazalis F, Reyes E, Leduc S, Gourion D. Evidence that nine autistic women out of ten have been victims of sexual violence. *Front Behav Neurosci.* (2022) 16:852203. doi: 10.3389/fnbeh.2022.852203
- 69. Goldblum JE, McFayden TC, Bristol S, Putnam OC, Wylie A, Harrop C. Autism prevalence and the intersectionality of assigned sex at birth, race, and ethnicity on age of diagnosis. *J Autism Dev Disord*. (2023) 54:3777–91. doi: 10.1007/s10803-023-06104-5
- 70. Kentrou V, Livingston LA, Grove R, Hoekstra RA, Begeer S. Perceived misdiagnosis of psychiatric conditions in autistic adults. *eClinicalMedicine*. (2024) 71:102586. doi: 10.1016/j.eclinm.2024.102586
- 71. Milner V, Colvert E, Hull L, Cook J, Ali D, Mandy W, et al. Does camouflaging predict age at autism diagnosis? A comparison of autistic men and women. *Autism Res.* (2023) 17(3):626–36. doi: 10.1002/aur.3059
- 72. Baldwin S, Costley D. The experiences and needs of female adults with high-functioning autism spectrum disorder. *Autism.* (2016) 20:483–95. doi: 10.1177/1362361315590805
- 73. Seers K, Hogg R. Fake it 'till you make it": Authenticity and wellbeing in late diagnosed autistic women. Fem Psychol. (2022) 33:23-41. doi: 10.1177/09593535221101455
- 74. Wilson RB, Thompson AR, Rowse G, Smith R, Dugdale AS, Freeth M. Autistic women's experiences of self-compassion after receiving their diagnosis in adulthood. *Autism.* (2023) 27:1336–47. doi: 10.1177/13623613221136752
- 75. Grove R, Clapham H, Moodie T, Gurrin S, Hall G. 'Living in a world that's not about us': The impact of everyday life on the health and wellbeing of autistic women and gender diverse people. *Women's Heal*. (2023) 19:1–16. doi: 10.1177/17455057231189542
- 76. Harmens M, Sedgewick F, Hobson H. Autistic women's diagnostic experiences: Interactions with identity and impacts on well-being. *Women's Heal.* (2022) 18. doi: 10.1177/17455057221137477
- 77. Harmens M, Sedgewick F, Hobson H. The quest for acceptance: A blog-based study of autistic women's experiences and well-being during autism identification and diagnosis. *Autism Adulthood.* (2022) 4:42–51. doi: 10.1089/aut.2021.0016
- 78. Leedham A, Thompson AR, Smith R, Freeth M. 'I was exhausted trying to figure it out': The experiences of females receiving an autism diagnosis in middle to late adulthood. *Autism.* (2020) 24:135–46. doi: 10.1177/1362361319853442
- 79. Lynch J, Murphy A, Cezar Da Cruz D. A phenomenological exploration of the occupational identity of late-diagnosed autistic women in the United Kingdom. *Discov Psychol.* (2024) 4:123. doi: 10.1007/s44202-024-00234-2
- 80. Milner V, McIntosh H, Colvert E, Happé F. A qualitative exploration of the female experience of autism spectrum disorder (ASD). *J Autism Dev Disord*. (2019) 49:2389–402. doi: 10.1007/s10803-019-03906-4
- 81. Nikolova-Fontaine K, Snæfríður Þóra Egilson. 'Even the Way I Make My Coffee is Autistic ': Meaning of Autism in the Lives of Middle-Aged Women prior to and After their Diagnoses. *Scand J Disabil Res.* (2023) 25:269–81. doi: 10.16993/sidr.1003
- 82. Powell T, Parker J, Kitson H, Rogalewski M. "It was like the final piece in the puzzle for me": A qualitative study on the experiences of autistic women initially diagnosed with borderline personality disorder. *Autism Adulthood.* (2024) 6:428–37. doi: 10.1089/aut.2023.0031
- 83. Seers K, Hogg RC. 'You don't look autistic': A qualitative exploration of women's experiences of being the 'autistic other.' *Autism.* (2021) 25:1553–64. doi: 10.1177/1362361321993722
- 84. Haney JL, Cullen JA. Learning about the lived experiences of women with autism from an online community. *J Soc Work Disabil Rehabil*. (2017) 16:54–73. doi: 10.1080/1536710X.2017.1260518
- 85. Hamdani Y, Kassee C, Walker M, Lunsky Y, Gladstone B, Sawyer A, et al. Roadblocks and detours on pathways to a clinical diagnosis of autism for girls and women: A qualitative secondary analysis. *Women's Heal.* (2023) 19. doi: 10.1177/17455057231163761
- 86. Curd J, Nguyen TST. I was always on the outside, watching quietly ": Autistic women re fl ect on school experiences. *Humanit Soc Sci Commun.* (2024) 11:1148. doi: 10.1057/s41599-024-03632-y
- 87. Murphy S, Flower RL, Jellett R. Women seeking an autism diagnosis in Australia: A qualitative exploration of factors that help and hinder. Autism. (2023) 27:808–21. doi: 10.1177/13623613221117911
- 88. Koffer Miller KH, Cooper DS, Song W, Shea LL. Self-reported service needs and barriers reported by autistic adults: Differences by gender identity. *Res Autism Spectr Disord.* (2022) 92:101916. doi: 10.1016/j.rasd.2022.101916
- 89. Tint A, Weiss JA, Lunsky Y. Identifying the clinical needs and patterns of health service use of adolescent girls and women with autism spectrum disorder. *Autism Res.*  $(2017)\ 10:1558-66$ . doi: 10.1002/aur.1806
- 90. Josefsson H, Taneja Johansson S. Adolescent autistic girls navigating their diagnosis in Swedish secondary school. *Front Educ.* (2024) 9:1461054. doi: 10.3389/feduc.2024.1461054

- 91. Zakai-Mashiach M. I was like a bird without wings": autistic women's retrospective experiences in general schools. *J Autism Dev Disord*. (2023) 53:4258–70. doi: 10.1007/s10803-022-05717-6
- 92. Jacobs P, Beamish W, McKay L. Please listen to us: Adolescent autistic girls speak about learning and academic success. *Improv Sch.* (2021) 24:196–209. doi: 10.1177/1365480220973112
- 93. Sproston K, Sedgewick F, Crane L. Autistic girls and school exclusion: Perspectives of students and their parents. *Autism Dev Lang Impair*. (2017) 2:1–14. doi: 10.1177/2396941517706172
- 94. Cook A, Ogden J, Winstone N. Friendship motivations, challenges and the role of masking for girls with autism in contrasting school settings. *Eur J Spec Needs Educ.* (2018) 33:302–15. doi: 10.1080/08856257.2017.1312797
- 95. Sung C, Sánchez J, Kuo HJ, Wang CC, Leahy MJ. Gender differences in vocational rehabilitation service predictors of successful competitive employment for transition-aged individuals with autism. *J Autism Dev Disord*. (2015) 45:3204–18. doi: 10.1007/s10803-015-2480-z
- 96. Taylor JL, Dawalt LS, Marvin AR, Law JK, Lipkin P. Sex differences in employment and supports for adults with autism spectrum disorder. *Autism.* (2019) 23:1711–9. doi: 10.1177/1362361319827417
- 97. Hayward SM, Mcvilly KR, Stokes MA. Always a glass ceiling. "Gender or autism; the barrier to occupational inclusion. *Res Autism Spectr Disord.* (2018) 56:50–60. doi: 10.1016/j.rasd.2018.09.001
- 98. Hayward SM, McVilly KR, Stokes MA. I would love to just be myself': what autistic women want at work. *Autism Adulthood*. (2019) 1:297–305. doi: 10.1089/aut.2019.0020
- 99. Gemma N. Reconceptualising 'reasonable adjustments' for the successful employment of autistic women. *Disabil Soc.* (2023) 38:944–62. doi: 10.1080/09687599.2021.1971065
- 100. Nagib W, Wilton R. Examining the gender role in workplace experiences among employed adults with autism: Evidence from an online community. *J Vocat Rehabil.* (2021) 55:27–42. doi: 10.3233/JVR-211144
- 101. Hayward SM, Stokes MA, McVilly KR. Short report: Gendered workplace social interaction processes in autism. *Res Dev Disabil.* (2022) 129:104310. doi: 10.1016/j.ridd.2022.104310
- $102.\,$  Dugdale AS, Thompson AR, Leedham A, Beail N, Freeth M. Intense connection and love: The experiences of autistic mothers. Autism.~(2021)~25:1973-84. doi: 10.1177/13623613211005987
- 103. Hampton S, Man J, Allison C, Aydin E, Baron-Cohen S, Holt R. A qualitative exploration of autistic mothers' experiences I: Pregnancy experiences. *Autism.* (2023) 27:1271–82. doi: 10.1177/13623613221132435
- 104. Karavidas M, de Visser RO. It's not just in my head, and it's not just irrelevant": autistic negotiations of menopausal transitions. *J Autism Dev Disord*. (2022) 52:1143–55. doi: 10.1007/s10803-021-05010-y
- 105. Moseley RL, Druce T, Turner-Cobb JM. 'When my autism broke': A qualitative study spotlighting autistic voices on menopause. Autism. (2020) 24:1423–37. doi: 10.1177/1362361319901184
- 106. Moseley RL, Druce T, Turner-Cobb JM. Autism research is 'all about the blokes and the kids': Autistic women breaking the silence on menopause. *Br J Health Psychol.* (2021) 26:709–26. doi: 10.1111/bjhp.12477
- 107. Lum M, Garnett M, O'Connor E. Health communication: A pilot study comparing perceptions of women with and without high functioning autism spectrum disorder. *Res Autism Spectr Disord*. (2014) 8:1713–21. doi: 10.1016/j.rasd.2014.09.009
- 108. Moore L, Foley S, Larkin F. Understanding the experiences of receiving and providing maternity care for autistic adults: A Multi-perspectival Interpretative Phenomenological Analysis study. *Autism.* (2024) 29(2):435–46. doi: 10.1177/13623613241274518
- 109. Babb C, Brede J, Jones CRG, Elliott M, Zanker C, Tchanturia K, et al. 'It's not that they don't want to access the support ... it's the impact of the autism': The experience of eating disorder services from the perspective of autistic women, parents and healthcare professionals. *Autism.* (2021) 25:1409–21. doi: 10.1177/1362361321991257
- 110. Babb C, Brede J, Jones CRG, Serpell L, Mandy W, Fox J. A comparison of the eating disorder service experiences of autistic and non-autistic women in the UK. *Eur Eat Disord Rev.* (2022) 30:616–27. doi: 10.1002/erv.2930
- 111. Fisher MH, Epstein RA, Urbano RC, Vehorn A, Cull MJ, Warren Z. A population-based examination of maltreatment referrals and substantiation for children with autism spectrum disorder. *Autism.* (2019) 23:1335–40. doi: 10.1177/1362361318813998
- 112. Ghirardi L, Kuja-Halkola R, Pettersson E, Sariaslan A, Arseneault L, Fazel S, et al. Neurodevelopmental disorders and subsequent risk of violent victimization: Exploring sex differences and mechanisms. *Psychol Med.* (2023) 53:1510–7. doi: 10.1017/S0033291721003093
- 113. Gibbs V, Hudson J, Hwang YI, Arnold S, Trollor J, Pellicano E. Experiences of physical and sexual violence as reported by autistic adults without intellectual disability: Rate, gender patterns and clinical correlates. *Res Autism Spectr Disord.* (2021) 89:101866. doi: 10.1016/j.rasd.2021.101866

- 114. Gibbs V, Hudson J, Pellicano E. The extent and nature of autistic people's violence experiences during adulthood: A cross-sectional study of victimisation. *J Autism Dev Disord*. (2023) 53:3509–24. doi: 10.1007/s10803-022-05647-3
- 115. Weiss JA, Fardella MA. Victimization and perpetration experiences of adults with autism. Front Psychiatry. (2018) 9:203. doi: 10.3389/fpsyt.2018.00203
- 116. Libster N, Knox A, Engin S, Geschwind D, Parish-Morris J, Kasari C. Personal victimization experiences of autistic and non-autistic children. *Mol Autism.* (2022) 13. doi: 10.1186/s13229-022-00531-4
- 117. Matthias C, LaVelle JM, Johnson DR, Wu YC, Thurlow ML. Exploring predictors of bullying and victimization of students with autism spectrum disorder (ASD): findings from NLTS 2012. *J Autism Dev Disord*. (2021) 51:4632–43. doi: 10.1007/s10803-021-04907-y
- 118. Ochi M, Kawabe K, Ochi S, Miyama T, Horiuchi F, Ueno S. School refusal and bullying in children with autism spectrum disorder. *Child Adolesc Psychiatry Ment Health.* (2020) 14. doi: 10.1186/s13034-020-00325-7
- 119. Schrooten I, Scholte RHJ, Cillessen AHN, Hymel S. Participant roles in bullying among dutch adolescents with autism spectrum disorders. *J Clin Child Adolesc Psychol.* (2018) 47:874–87. doi: 10.1080/15374416.2016.1138411
- 120. Dean M, Kasari C, Shih W, Frankel F, Whitney R, Landa R, et al. The peer relationships of girls with ASD at school: Comparison to boys and girls with and without ASD. *J Child Psychol Psychiatry Allied Discip.* (2014) 55:1218–25. doi: 10.1111/jcpp.12242
- 121. Sedgewick F, Hill V, Pellicano E. 'It's different for girls': Gender differences in the friendships and conflict of autistic and neurotypical adolescents. *Autism.* (2019) 23:1119–32. doi: 10.1177/1362361318794930
- 122. Kanfiszer L, Davies F, Collins S. 'I was just so different': The experiences of women diagnosed with an autism spectrum disorder in adulthood in relation to gender and social relationships. *Autism.* (2017) 21:661–9. doi: 10.1177/1362361316687987
- 123. Shalit L, Cochavit E, Roginsky E. Exploring music in the everyday lives of autistic women: An Interpretative Phenomenological Analysis. *Nord J Music Ther.* (2024) 34(3):225–44. doi: 10.1080/08098131.2024.2396105
- 124. Mo S, Viljoen N, Sharma S. The impact of socio-cultural values on autistic women: An interpretative phenomenological analysis. *Autism.* (2022) 26:951–62. doi: 10.1177/13623613211037896
- 125. Myles O, Boyle C, Richards A. The social experiences and sense of belonging in adolescent females with autism in mainstream school. *Educ Child Psychol.* (2019) 36:8–21. doi: 10.53841/bpsecp.2019.36.4.8
- 126. Ryan C, Coughlan M, Maher J, Vicario P, Ryan C, Coughlan M, et al. Perceptions of friendship among girls with Autism Spectrum Disorders. *Eur J Spec Needs Educ.* (2021) 36:393–407. doi: 10.1080/08856257.2020.1755930
- 127. Vine Foggo RS, Webster AA. Understanding the social experiences of adolescent females on the autism spectrum. *Res Autism Spectr Disord.* (2017) 35:74–85. doi: 10.1016/j.rasd.2016.11.006
- 128. Tierney S, Burns J, Kilbey E. Looking behind the mask: Social coping strategies of girls on the autistic spectrum. *Res Autism Spectr Disord.* (2016) 23:73–83. doi: 10.1016/j.rasd.2015.11.013
- 129. Joyal CC, Carpentier J, McKinnon S, Normand CL, Poulin M-H. Sexual knowledge, desires, and experience of adolescents and young adults with an autism spectrum disorder: an exploratory study. *Front Psychiatry*. (2021) 12:1–19. doi: 10.3389/fpsyt.2021.685256
- 130. Pecora LA, Hancock GI, Mesibov GB, Stokes MA. Characterising the sexuality and sexual experiences of autistic females. *J Autism Dev Disord*. (2019) 49:4834–46. doi: 10.1007/s10803-019-04204-9
- 131. Brown-Lavoie SM, Viecili MA, Weiss JA. Sexual knowledge and victimization in adults with autism spectrum disorders. *J Autism Dev Disord*. (2014) 44:2185–96. doi: 10.1007/s10803-014-2093-y
- 132. Kock E, Strydom A, O'Brady D, Tantam D. Autistic women's experience of intimate relationships: the impact of an adult diagnosis. *Adv Autism.* (2019) 5:38–49. doi: 10.1108/AIA-09-2018-0035
- 133. Macmillan K, Berg T, Just M, Stewart ME. Online safety experiences of autistic young people: An Interpretative Phenomenological Analysis. *Res Autism Spectr Disord*. (2022) 96:101995. doi: 10.1016/j.rasd.2022.101995
- 134. Pecora LA, Hancock GI, Hooley M, Demmer DH, Attwood T, Mesibov GB, et al. Gender identity, sexual orientation and adverse sexual experiences in autistic females. *Mol Autism.* (2020) 11:1–17. doi: 10.1186/s13229-020-00363-0
- 135. Paricos A, Sturrock A, Twomey K, Leadbitter K. I 'm not mad, bad, and dangerous ... simply wired differently": Exploring factors contributing to good quality of life with autistic women. *Res Autism Spectr Disord*. (2024) 112:102338. doi: 10.1016/j.rasd.2024.102338
- 136. Thom-Jones S, Melgaard I, Gordon CS. Autistic women's experience of motherhood: A qualitative analysis of reddit. *J Autism Dev Disord*. (2024) 55:2539–49. doi: 10.1007/s10803-024-06312-7
- 137. Hampton S, Man J, Allison C, Aydin E, Baron-Cohen S, Holt R. A qualitative exploration of autistic mothers' experiences II: Childbirth and postnatal experiences. *Autism.* (2022) 26:1165–75. doi: 10.1177/13623613211043701

- 138. Sunagawa M. How much of my true self can I show? social adaptation in autistic women: a qualitative study. *BMC Psychol.* (2023) 11. doi: 10.1186/s40359-023-01192-5
- 139. Sedgewick F, Crane L, Hill V, Pellicano E. Friends and lovers: the relationships of autistic and neurotypical women. *Autism Adulthood.* (2019) 1:112–23. doi: 10.1089/aut.2018.0028
- 140. Tint A, Weiss JA. A qualitative study of the service experiences of women with autism spectrum disorder. *Autism.* (2018) 22:928–37. doi: 10.1177/1362361317702561
- 141. Lewis LF. A mixed methods study of barriers to formal diagnosis of autism spectrum disorder in adults. *J Autism Dev Disord*. (2017) 47:2410–24. doi: 10.1007/s10803-017-3168-3
- 142. Morgan R. How do adolescent autistic girls construct self-concept and social identity? A discourse analysis. *Educ Psychol Pract.* (2023) 39:178–200. doi: 10.1080/02667363.2023.2181316
- 143. Garcia-Molina I. When I need help, I ask my friends": experiences of Spanish autistic women when disclosing their late diagnosis to family and friends. *Front Psychol.* (2024) 15:1342352. doi: 10.3389/fpsyg.2024.1342352
- 144. Cooper K, Smith LGE, Russell A. Social identity, self-esteem, and mental health in autism. *Eur J Soc Psychol.* (2017) 47:844–54. doi: 10.1002/ejsp.2297
- 145. Davies J, Cooper K, Killick E, Sam E, Healy M, Thompson G, et al. Autistic identity: A systematic review of quantitative research. *Autism Res.* (2024) 17:874–97. doi: 10.1002/aur.3105
- 146. McDonald TAM. Discriminative and criterion validity of the autism spectrum identity scale (ASIS). *J Autism Dev Disord*. (2017) 47:3018–28. doi: 10.1007/s10803-017-3221-2
- 147. Frost DM, Meyer IH. Measuring community connectedness among diverse sexual minority populations. J Sex Res. (2012) 49:36–49. doi: 10.1080/00224499.2011.565427
- 148. Donovan J, Chiatti BD, McKeever A, Bloch JR, Gonzales MS, Birati Y. Yes, I can bond." Reflections of autistic women's mothering experiences in the early postpartum period. *Women's Heal.* (2023) 19:1–13. doi: 10.1177/17455057231175312
- 149. Kourti M, MacLeod A. I don't feel like a gender, I feel like myself": autistic individuals raised as girls exploring gender identity. *Autism Adulthood.* (2019) 1:52–9. doi: 10.1089/aut.2018.0001
- 150. Libster N, Knox A, Engin S, Geschwind D, Parish-Morris J, Kasari C. Sex differences in friendships and loneliness in autistic and non-autistic children across development. *Mol Autism.* (2023) 14. doi: 10.1186/s13229-023-00542-9
- 151. Webster AA, Garvis S. The importance of critical life moments: An explorative study of successful women with autism spectrum disorder. *Autism.* (2017) 21:670–7. doi: 10.1177/1362361316677719
- 152. Sedgewick F, Hill V, Yates R, Pickering L. Gender differences in the social motivation and friendship experiences of autistic and non-autistic adolescents. *J Autism Dev Disord.* (2016) 46:1297–306. doi: 10.1007/s10803-015-2669-1
- 153. Dean M, Chang YC, Shih W, Orlich F, Kasari C. Social engagement and loneliness in school-age autistic girls and boys. *Women's Heal.* (2023) 19:1–12. doi: 10.1177/17455057231170973
- 154. Corden K, Brewer R, Cage E. Personal identity after an autism diagnosis: relationships with self-esteem, mental wellbeing, and diagnostic timing. *Front Psychol.* (2021) 12. doi: 10.3389/fpsyg.2021.699335
- 155. Shi P, Yang A, Zhao Q, Chen Z, Ren X, Dai Q. A hypothesis of gender differences in self-reporting symptom of depression: implications to solve under-diagnosis and under-treatment of depression in males. *Front Psychiatry*. (2021) 12:589687. doi: 10.3389/fpsyt.2021.589687
- 156. Oliffe JL, Rossnagel E, Seidler ZE, Kealy D, Ogrodniczuk JS, Rice SM. Men's depression and suicide. *Curr Psychiatry Rep.* (2019) 21:103. doi: 10.1007/s11920-019-1088-v
- 157. Cooper K, Smith LGE, Russell AJ. Gender identity in autism: sex differences in social affiliation with gender groups. *J Autism Dev Disord.* (2018) 48:3995–4006. doi: 10.1007/s10803-018-3590-1
- 158. Dewinter J, De Graaf H, Begeer S. Sexual orientation, gender identity, and romantic relationships in adolescents and adults with autism spectrum disorder. *J Autism Dev Disord.* (2017) 47:2927–34. doi: 10.1007/s10803-017-3199-9
- 159. George R, Stokes MA. Gender identity and sexual orientation in autism spectrum disorder. *Autism.* (2018) 22:970–82. doi: 10.1177/1362361317714587
- 160. George R, Stokes MA. A quantitative analysis of mental health among sexual and gender minority groups in ASD. J Autism Dev Disord. (2018) 48:2052–63. doi: 10.1007/s10803-018-3469-1
- 161. Brickhill R, Atherton G, Piovesan A, Cross L. Autism, thy name is man: Exploring implicit and explicit gender bias in autism perceptions. *PloS One.* (2023) 18: e0284013. doi: 10.1371/journal.pone.0284013
- 162. Ward TB, Curtis C, Seehagen S. Investigating the effects of perceived student gender on primary school teachers' recognition of autism. *Psychol Sch.* (2022) 59:1363–76. doi: 10.1002/pits.22667
- 163. Whitlock A, Fulton K, Lai MC, Pellicano E, Mandy W. Recognition of girls on the autism spectrum by primary school educators: an experimental study. *Autism Res.* (2020) 13:1358–72. doi: 10.1002/aur.2316

164. Anderson J, Marley C, Gillespie-Smith K, Carter L, MacMahon K. When the mask comes off: Mothers' experiences of parenting a daughter with autism spectrum condition. *Autism.* (2020) 24:1546–56. doi: 10.1177/1362361320913668

- 165. McCormick S, Thomas G. Exploring parents' experiences of working with educational psychologists to support their autistic daughters. *Educ Psychol Pract.* (2024) 41:1–23. doi: 10.1080/02667363.2024.2414456
- 166. Muggleton JTB, MacMahon K, Johnston K. Exactly the same but completely different: A thematic analysis of Clinical Psychologists' conceptions of Autism across genders. *Res Autism Spectr Disord.* (2019) 62:75–84. doi: 10.1016/j.rasd.2019.03.004
- 167. Tsirgiotis JM, Young RL, Weber N. A mixed-methods investigation of diagnostician sex/gender-bias and challenges in assessing females for autism spectrum disorder. *J Autism Dev Disord*. (2022) 52:4474–89. doi: 10.1007/s10803-021-05300-5
- 168. Crane L, Hearst C, Ashworth M, Davies J, Hill EL. Supporting newly identified or diagnosed autistic adults: an initial evaluation of an autistic-led programme. J Autism Dev Disord. (2021) 51:892–905. doi: 10.1007/s10803-020-04486-4
- 169. Crompton CJ, Hallett S, McAuliffe C, Stanfield AC, Fletcher-Watson S. A group of fellow travellers who understand\*: interviews with autistic people about post-diagnostic peer support in adulthood. *Front Psychol.* (2022) 13. doi: 10.3389/fpsyg.2022.831628