



OPEN ACCESS

EDITED AND REVIEWED BY

Ahmed El Nemr,
National Institute of Oceanography and
Fisheries (NIOF), Egypt

*CORRESPONDENCE

Cheng Yan
✉ cheng_yan@cug.edu.cn
Carla Viegas
✉ carla.viegas@estesl.ipl.pt
Yunping Han
✉ yphan@rcees.ac.cn
Annalaura Carducci
✉ annalaura.carducci@unipi.it

SPECIALTY SECTION

This article was submitted to
Environmental Health and Exposome,
a section of the journal
Frontiers in Public Health

RECEIVED 29 November 2022

ACCEPTED 13 January 2023

PUBLISHED 17 February 2023

CITATION

Yan C, Viegas C, Han Y and Carducci A (2023)
Editorial: Bioaerosol emission characteristics
and the epidemiological, occupational, and
public health risk assessment of waste and
wastewater management.
Front. Public Health 11:1111457.
doi: 10.3389/fpubh.2023.1111457

COPYRIGHT

© 2023 Yan, Viegas, Han and Carducci. This is
an open-access article distributed under the
terms of the [Creative Commons Attribution
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that the
original publication in this journal is cited, in
accordance with accepted academic practice.
No use, distribution or reproduction is
permitted which does not comply with these
terms.

Editorial: Bioaerosol emission characteristics and the epidemiological, occupational, and public health risk assessment of waste and wastewater management

Cheng Yan^{1*}, Carla Viegas^{2,3,4*}, Yunping Han^{5,6*} and
Annalaura Carducci^{7*}

¹School of Environmental Studies, China University of Geosciences, Wuhan, China, ²H&TRC - Health & Technology Research Center, ESTeSL - Escola Superior de Tecnologia da Saúde, Instituto Politécnico de Lisboa, Lisbon, Portugal, ³NOVA National School of Public Health, Public Health Research Centre, Comprehensive Health Research Center, CHRC, NOVA University Lisbon, Lisbon, Portugal, ⁴Comprehensive Health Research Center (CHRC), Lisbon, Portugal, ⁵State Key Laboratory of Environmental Aquatic Chemistry, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing, China, ⁶University of Chinese Academy of Sciences, Beijing, China, ⁷Laboratory of Hygiene and Environmental Virology, Department of Biology, University of Pisa, Pisa, Italy

KEYWORDS

wastewater treatment, bioaerosol, source apportionment, emission release principle, occupational and public health risk assessment

Editorial on the Research Topic

[Bioaerosol emission characteristics and the epidemiological, occupational, and public health risk assessment of waste and wastewater management](#)

Why is important to assess occupational exposure to bioaerosols

Waste management industries are vital to achieve the Sustainable Development Goals (SDGs) suggested by World Health Organization (1). Besides SDGs achievement, also the circular economy is dependent of the waste sector. The circular economy intends to re-use the existing resources instead of disposing materials that are no longer useful (2). Thus, the circular economy is very dependent of an increased workforce dedicated to waste management. Nonetheless, while better waste management is being predicted as a critical contribute to reduce health outcomes and environmental negative impacts, the bioaerosols' occupational exposure in waste sector is being neglected leading to negative outcomes on workers' health (1).

The scientific community and stakeholders should be also aware of the foreseen increased exposure to microbes and antimicrobial resistance in different environments that will lead to a boost of exposure to bioaerosols (for instance mycotoxins and climate-sensitive infectious diseases) in the waste sector, mainly due to climate change (3, 4).

Overview of the published papers

Aiming to assess health risks due to bioaerosol emission and air pollution, four articles contributed with their results. [Kontró et al.](#) found that bioaerosol levels were high especially in the composting compared to bioenergy producing facilities. Endotoxin, *A. fumigatus* and *Streptomyces* spp. detected in bioaerosols were also observed in the nasal passages of the workers, indicating that bioaerosols in composting plant has great potential to harm to workers' health.

[Pascale et al.](#) revealed high contamination levels and large microbial heterogeneity both for PM and bioaerosol samples. They also found *Bacillus* spp., *Saccharopolyspora* spp., and *Thermomyces* spp. may be suggested as indicators of biological contamination in composting plants. In addition, this work showed that using multiple assays is helpful to ensure the needed sensitivity and accuracy of bioaerosol detection.

[Pan et al.](#) assessed airborne bacterial community in electronic waste dismantling site and a waste transfer station based on culture-dependent and culture-independent methods. Bacterial communities in waste-associated bioaerosols were predominated by *Proteobacteria* spp. and *Bacteroidetes* spp.. One-third of the species in these genera were uncultured. Differences community structure existed in airborne bacterial diversity among different sampling sites, showing that waste-associated environments have unique bacterial diversity.

[Bai et al.](#) used the Taiwan Longitudinal Health Insurance Database and the Taiwan Air Quality Monitoring Database to conduct a retrospective cohort study to investigate whether air pollution increases the risk of uveitis. Overall, 175,489 subjects were linked to their nearby air quality monitoring stations; air pollution was significantly associated with incidental uveitis, especially at high total hydrocarbon (THC) and CH₄ levels; and uveitis risk increased with increasing NO_x and THC levels.

For public health risk assessment of wastewater, [Lin et al.](#) reviewed the effects of water pollution on human health and disease heterogeneity. Eighty-five relevant papers were selected. The results shown that the impact of water pollution on human health is significant, although there may be regional, age, gender, and other differences in degree. Taken together, diarrhea caused by enteroviruses in aquatic environments is the most common disease caused by water pollution. Therefore, some suggestions about strengthening water intervention management and carrying out intervention water quality measures have been put forward.

In addition, for health risk of waste management, [Ruppen et al.](#) reported that Hwange, Western Zimbabwe used a community-based monitoring to identify sources of pollution and related these to past and present mining activities in a river downstream of a coal mining area. The primary source of acid mine drainage came from abandoned underground mine sites. Concentrations of Mn, Ni, and As were exceeding national fresh water guidelines and international drinking water standards. Results showed that this community-based monitoring offers a promising approach to establish a high-quality dataset for assessing risks.

Main findings and research needed

The exposure to bioaerosols in solid or liquid waste processing facilities and their surrounding can lead to

different types of adverse effects. Although they are well-documented by epidemiological surveys and numerous studies have monitored the air microbial contamination in these settings, a clear link between the environmental microbial pollution and diseases of workers or population is still lacking.

To better understand the relations between bioaerosol concentration and composition and risks for human health future research is needed, addressing several points:

- 1) Epidemiological studies and dose-response estimation. The evaluation of relationship between microbial contamination and health problems is very complex, because of the diversity of microorganisms in the bioaerosol and the variability of their distribution according to the area and time, as well as the difference in human susceptibility and reactions. Large and long lasting cohort studies, monitoring the air microbial contamination during the follow up could provide answers, but they are hampered by the complexity and costs. Moreover, the current use of personal protective equipment would hide the real risk.
- 2) Exposure assessment. The choice of appropriate indicators or index pathogens should take in account the aim of the study and their measure should be easy, timely, and representative of risk. For safety purposes total bacterial and fungal counts and fecal indicators are commonly used. Nevertheless, the lack of epidemiological studies hampers the definition of limits for an acceptable risk. On the other hand, the study of microbial composition of bioaerosol with cultural or non-cultural methods can show the presence and concentration of pathogens, allowing a more specific risk assessment (5, 6). Nevertheless, the proposed methods are highly variable and often lacking validation and standardization for risk assessment purposes. Thus, further studies are needed taking in account sensitivity and specificity, besides the real meaning of analytical results, e.g., of non-cultural methods toward the pathogens infectivity.
- 3) Risk assessment. The application of the qualitative or semi-quantitative methods can be useful for risk management, to apply preventive and protective measures, but only the Quantitative Microbial Risk Assessment (QMRA) can allow defining acceptable limits and to simulate and evaluate different scenarios (7). Moreover, interactions between bioaerosol infective components and air particulate and gaseous pollutants should be better studied to clarify their role in enhancing or reducing the microbial risk.

Author contributions

CY, CV, YH, and AC contributed to conception and design of the study, organized the database, performed the statistical analysis, wrote the first draft of the manuscript, wrote sections of the manuscript. All authors contributed to manuscript revision, read, and approved the submitted version.

Acknowledgments

H&TRC author gratefully acknowledge the FCT/MCTES national support through the UIDB/05608/2020 and the UIDP/05608/2020.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships

that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

1. Viegas C, Pena P, Dias M, Gomes B, Cervantes R, Carolino E, et al. Microbial contamination in waste collection: unveiling this Portuguese occupational exposure scenario. *J Environ Manage.* (2022) 314:115086. doi: 10.1016/j.jenvman.2022.115086
2. European Public Service Association. *Waste Management in Europe. Good Jobs in the Circular Economy?* (2017). Available online at: <http://gala.gre.ac.uk/id/eprint/19920> (accessed December 7, 2017)
3. Viegas C. Climate change influence in fungi. *Eur J Public Health.* (2021) 31:164–270. doi: 10.1093/eurpub/ckab164.270
4. European Centre for Disease Prevention, and Control (2021). *Extreme Rainfall and Catastrophic Floods in Western Europe.* Available online at: <https://www.ecdc.europa.eu/sites/default/files/documents/RRA-20210720-1799.pdf> (accessed July 29, 2021)
5. Lanzarini NM, Federigi I, Mata RM, Borges MDN, Saggiaro EM, Cioni L, et al. Human adenovirus in municipal solid waste leachate and quantitative risk assessment of gastrointestinal illness to waste collectors. *Waste Manage.* (2022) 138:308–17. doi: 10.1016/j.wasman.2021.12.005
6. Tian J, Yan C, Alcega SG, Hassard F, Tyrrel S, Coulon F, et al. Detection and characterization of bioaerosol emissions from wastewater treatment plants: challenges and opportunities. *Front Microbiol.* (2022) 13:958514. doi: 10.3389/fmicb.2022.958514
7. Carducci A, Donzelli G, Cioni L, Federigi I, Lombardi R, Verani M. Quantitative microbial risk assessment for workers exposed to bioaerosol in wastewater treatment plants aimed at the choice and setup of safety measures. *Int J Environ Res Public Health.* (2018) 15:1490. doi: 10.3390/ijerph15071490