



OPEN ACCESS

EDITED BY

Francisco Pitanga,
Federal University of Bahia (UFBA), Brazil

REVIEWED BY

Liliana C. Baptista,
University of Alabama at Birmingham,
United States

*CORRESPONDENCE

Alberto Jiménez-Maldonado
✉ jimenez.alberto86@uabc.edu.mx

RECEIVED 10 January 2024

ACCEPTED 11 March 2024

PUBLISHED 28 March 2024

CITATION

Jiménez-Maldonado A, Rentería I,
Johnson DK, Moncada-Jiménez J and
García-Suárez PC (2024) Physical exercise and
cognition in older adults, a scientific approach
scanty reported in Latin America and
Caribbean populations.
Front. Sports Act. Living 6:1368593.
doi: 10.3389/fspor.2024.1368593

COPYRIGHT

© 2024 Jiménez-Maldonado, Rentería,
Johnson, Moncada-Jiménez and
García-Suárez. This is an open-access article
distributed under the terms of the [Creative
Commons Attribution License \(CC BY\)](#). The
use, distribution or reproduction in other
forums is permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication in
this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Physical exercise and cognition in older adults, a scientific approach scanty reported in Latin America and Caribbean populations

Alberto Jiménez-Maldonado^{1*}, Iván Rentería¹, David K. Johnson²,
José Moncada-Jiménez³ and Patricia C. García-Suárez^{1,4}

¹Facultad de Deportes, Universidad Autónoma de Baja California, Ensenada, México, ²Department of Neurology, University of California, Davis, CA, United States, ³Human Movement Sciences Research Center (CIMOHU), University of Costa Rica, San Jose, Costa Rica, ⁴Department of Health, Sports and Exercise Sciences, University of Kansas, Lawrence, KS, United States

The advancement of public services, including the increased accessibility of health services, has led to a rise in life expectancy globally. As a result, aging populations are becoming more prevalent, raising concerns about cognitive decline. Fortunately, non-pharmacological methods, such as physical exercise, have been shown to mitigate the effects of aging on the brain. In this perspective article, we examined meta-analyses on the impact of physical exercise on cognition in older adults. The results indicate that combined exercise (i.e., aerobic plus strength training), has a significant positive effect on overall cognition and executive function. However, we found a lack of scientific studies on this topic in Latin American and Caribbean countries. Therefore, there is a pressing need for research to identify the feasibility of physical exercise interventions to improve cognitive skills in older adults from these regions.

KEYWORDS

physical exercise, general cognition, executive functions, older adults, Latin America

1 Introduction

According to the World Health Organization (WHO) (1), life expectancy has increased since the past century, and by 2030, 1 in 6 individuals will be aged 60 or more, and epidemiological projections indicate that it is expected that the world will be inhabited by 2.1 billion older adults in 2050 (2). Nevertheless, the downside of this demographic transition in Latin America and Caribbean (LAC) countries is the heterogeneous and distributed social and health disparity-related risk factors for cognitive aging and dementia, a situation that is emphasized by the inadequate infrastructure and scarce scientific approaches to support the mental health in the individuals at the Caribbean Community (3). The growing elderly's demography emphasizes that health professionals should deeply understand older adults' biology and behavior pattern changes. In this sense, it is well known that morpho-functional changes in the brain accompany human aging (4). Some of these include a decline in the hippocampus volume (4), morphological changes in the cortex regions (neocortex areas) (5, 6), and changes in the brain metabolism (7) [see Matsson et al. (7) for an extensive review].

The age-related changes mentioned above enhance older adults' susceptibility to mental health conditions such as depression, anxiety, sleep disorders (6), and cognitive decline (4), being the last condition a precursor of a more severe disorder such as the Alzheimer's Disease (AD). The prevalence of AD in the general population of the United States in 2020 was 11.3% (i.e., approximately 6 million inhabitants) and 14.0% among Hispanics (i.e., approximately 0.71 million inhabitants) (8). The number of individuals in the general population with AD is expected to increase more than twice (i.e., approximately 13.85 million individuals) by 2060. It will be 423% higher among Hispanics (i.e., approximately 3.72 million inhabitants), yet the impact will be higher in adults older than 85, especially women (8).

Besides the natural aging process's impact on the human brain, voluntary movement has also become critical in protecting and maintaining cognitive function. It is known that older adults are among the most sedentary individuals, and higher sedentary time has been directly related to an increased risk of all-cause mortality in that population (9). Neuroscientists, exercise physiologists, and diverse health professionals have made significant efforts to identify pharmacological and non-pharmacological interventions focused on strengthening the brain's health in aging individuals. Owing to relevant scientific evidence, there is a consensus that physical exercise (PE) is a feasible method to attenuate aging-induced brain damage (10, 11). Some reports have identified that PE enhances the neuronal circuit in healthy individuals; therefore, the exercise benefits are not limited to a population with mental health disorders (12). Several randomized controlled trials have been the foundation for published meta-analyses (12–16), which analyzed in detail the impact of different exercise protocols on cognitive outcomes improvements in older adults, including variables such as general cognition, executive functions, and memory. These and other conditions are responsible for supporting real-life activities that allow older adults to orchestrate the complexities of daily tasks to plan actions, organize information, think abstractly, allocate mental resources, reason, solve novel problems, adapt to new situations, and act appropriately during social interactions (12–16). The current viewpoint article discusses the evidence reported in the meta-analyses about the impact of PE on cognitive control in older adults. At the end of the article, we present a brief reflection on how some variables hinder the rapid application of the evidence published in the countries of LAC.

2 Physical exercise on general cognition

Depending on the source, general cognition, also known as General Cognitive Ability (GCA) and defined as general mental ability, has been evaluated with valid and reliable instruments such as the Mini-Mental State Examination (MMSE) and the Montreal Cognitive Assessment (MoCA) (17). Meta-analytical evidence summarized in healthy older adults showed that studies implementing multicomponent interventions (i.e., protocols that included aerobic and strength exercises) were considered the

most robust interventions for improving GCA (Figure 1A) (12–16). In the proposed molecular mechanism elicited by the multicomponent exercise interventions, the researchers highlighted neural growth factors (e.g., BDNF, IGF-1) as the main molecules synthesized during the resistance exercises and the regulators to induce more significant benefits in the brain compared with single aerobic interventions (12, 14). In addition, it was also suggested that the anaerobic resistance exercises (i.e., strength) increase the lactate concentration in the bloodstream; once in circulating blood, the lactate reaches the brain and enhances the expression of genes linked with cognition (e.g., BDNF) (18, 19). Those potential adaptations synergize with the well-identified effects induced by aerobic exercise, such as increased blood flow to the brain and upregulation of the brain's metabolism (20–22) (Figure 2A).

Together with the benefits of PE on brain functions in healthy older adults, the impact of exercise on general cognition in Mild Cognitive Impairment (MCI) patients has been another subject analyzed in published systematic reviews and meta-analyses (12, 15, 23–28). Figure 1B describes the origin of the individuals included in those studies. MCI is a condition that implicates severe cognitive decline compared with the aging effects *per se*; in this sense, it is known that MCI may continue until developing dementia and AD (29–32). The MCI (all-types) prevalence in LAC is 14.95%, values higher than those observed in high-income countries (e.g., United States and England) (33). Similar to other illnesses, several social and health factors increase the risk of developing MCI; the more commonly reported are education level, cardiometabolic diseases (i.e., hypertension, diabetes), vitamin D deficiency, and apolipoprotein E (APOE) e4 genotype (29, 32). In regard with this, was identified that LAC shows a socioeconomic environment, that increase substantially the risk to suffer MCI (33, 34). Considering MCI patients, and similar to healthy older adults, the protocols that include resistance exercises seem to be the most potent intervention to attenuate the disruption of the neuronal network linked with cognition (24, 27). Several potential mechanisms have been suggested to explain the positive effect of exercise that allows the recovery of the microenvironment in the brain, leading to attenuating the MCI progression. Some of these include the turn-up of trophic factors (e.g., BDNF, IGF-1), regulation of the redox system, increase in mitochondrial mass, angiogenesis, and neurogenesis; moreover, improved glucose metabolism was also reported (24, 35) (Figure 2B).

3 Physical exercise on executive functions

From another perspective, besides GCA, the effects of PE on executive functions (EFs) in older adults free from MCI have also been meta-analyzed (Figure 1C) (12–14, 36–38). The EFs are considered high-order cognitive processes unregulated by emotions and include selective attention, resistance to interference, working memory, mental flexibility, planning, verbal reasoning, and feedback utilization (39–42). It is known that the





FIGURE 1 Frequency of studies included in meta-analyses on the effects of physical exercise on (A) general cognition in older adults without diseases, (B) general cognition in patients with mild cognitive impairment, (C) executive functions in older adults without diseases, and (D) executive functions in patients with mild cognitive impairment.

frontal lobe and parietal subcortical regions are activated during those tasks (43). The conclusions drawn from the current evidence are equivocal. For instance, a classical meta-analysis by Colcombe et al. (16) indicated that combined exercise interventions (i.e., aerobic exercises plus strength training) have a greater effect size on cognitive tasks, including EFs. The authors emphasized the dedifferentiation hypothesis to partially explain their findings (16). The hypothesis suggests that older adults recruit additional cortical areas to compensate for losses in neural efficiency.

On the contrary, other evidence that analyzed an aged population without diagnosed cognitive impairment found a non-significant effect of aerobic and resistance training on EFs (13). Indeed, the same study reported a significant effect of mind-body exercise (Tai Chi) on some EFs (i.e., working memory and processing speed). The authors emphasized mind-body exercise features to support their results; concretely, the researchers mentioned that the aerobic, strength, and flexibility exercises included in the Tai Chi movement sessions synergize to

generate neuronal changes leading to improvements in the performance of cognitive activities, which included EFs (13). In agreement with the previous reports, others have also identified a significant effect of resistance, aerobic, and multicomponent training on EFs in healthy older adults ≥ 50 years (37). At this point, it is worth mentioning that throughout their analysis, the researchers found a pronounced benefit of resistance training on EFs (14, 37). In concordance with the above mentioned, a more recent meta-analysis concluded that PE is an effective intervention aimed at improving memory and EFs; in their study (12), the authors indicated that the BDNF synthesis, neocortical modifications, functional connectivity, and changes in the hippocampus volume as the principal adaptations to attain improvements in the cognitive performance.

On the other hand, the effects of PE on EFs in populations with MCI have also been recently meta-analyzed (Figure 1D) (12, 24, 27, 42). However, in this population, the findings are controversial; some authors reported null effects of PE (i.e., aerobic, strength, or multicomponent training programs) on EFs (12), on the



contrary, recent evidence identified that resistance exercise induces the highest impact on the EFs compared with aerobic exercise (24, 44). The MCI has several molecular sources, which involve the accumulation of beta-amyloid plaques and neurofibrillary tangles (both molecular biomarkers of AD), glucose hypometabolism, cellular senescence, and neuronal atrophy (45–47). Subsequently, the PE interventions included in the meta-analytical evidence were likely insufficient (e.g., session duration, frequency of training, and exercise intensity) to induce molecular adaptations to improve the performance of EFs in MCI patients (12). Furthermore, it is worth mentioning that the MCI has subtypes; in general, those included amnesic MCI patients, who showed poor performance in episodic memory assessments, and non-amnesic MCI patients, who did not show fail in memory but displayed poor performance in other cognitive domains such as EFs, language, and visuospatial abilities (46). In the inclusion criteria of the articles, those characteristics were not considered (24, 44), which increased the risk of bias, leading to a high heterogeneity in the findings.

4 Physical exercise in the late adulthood in Latin America and Caribbean

Despite the promising results now communicated, unfortunately, the data reported in Figure 1 indicate that the LAC populations have been poorly studied; this situation hinders

the direct applications of exercise recommendations as a non-pharmacological treatment to improve general cognition in older adults (10). In the following lines, we will emphasize some variables that could attenuate the benefits of PE on the brain functions (i.e., general cognition and EFs) in older adults from LAC countries. One reason is the education level; cross-sectional studies have constantly found a low education level in the older adult population from Mexico, Colombia, Costa Rica, and Brazil (48–51). On the contrary, in the United States, older adults show a higher education level than in LAC countries (52). In this sense, and knowing that it is a complex analysis, authors have mentioned that the education level is a moderator variable in the cognitive performance of older adults (53–56). Moreover, some reports indicated that education is more relevant for cognitive domains like memory (54). The prior information allows us to infer a lesser benefit of PE on the GCA in the older adults from LAC.

Moreover, the income differences among LAC countries and the world regions reported in Figure 1 are another factor that deserves attention. As it is well-known and in agreement with a recent World Bank (WB) report in 2023 (57), Mexico, Brazil, and Argentina show emerging markets and developing economies, while the United States, Canada, Australia, and Japan are targeted as advanced economies. These economic differences considerably impact the quality of life for the older adult population in each country. One of the potential disadvantages for the Latin American and Caribbean populations is their nutritional status. A recent report from the Food and Agriculture Organization (58) indicated that food insecurity is higher in LAC

than in Asia, North America, and Europe (58). In this sense, current evidence indicates that the nutrition level is a determinant factor for brain functionality (59, 60). In light of this, it was suggested that a protein intake deficiency disturbs the antioxidant capacity, increasing the reactive oxygen species (ROS) levels and biomolecules that might potentially harm the brain (61, 62). Indeed, basic studies reported that a low-protein diet is linked with a decline in the neurotransmitter concentration associated with cognitive functions (63).

5 Conclusions

In regard with the meta-analyses articles analyzed, few studies from LAC have been carried out, however, we do not discard that the scarce evidences until now reported in the common database can be consequence by the language, all the articles analyzed were written in English language, we consider that systematic review and quantitative approach limited to Spanish, and Portuguese language are justified to identify the pool effect of PE on the cognition skills in LAC, with this activity, the barrier language could be cope. On the other hand, studies are also needed to clarify the participation of the variables above discussed in the responses to PE interventions and, consequently, to identify specific recommendations about the practice of PE to improve the brain function in older adults from LAC Countries. Finally, we emphasize that performing little PE is always better than none, and therefore, physical activity must be included in the lifestyle of older adults. This healthy habit will induce emotional and physiological benefits and potentially attenuate the Governments' health economic burden of countries classified as emerging and low-world economies.

Data availability statement

The original contributions presented in the study are included in the article/supplementary material, further inquiries can be directed to the corresponding author/s.

Author contributions

AJ-M: Conceptualization, Funding acquisition, Writing – original draft. IR: Formal Analysis, Writing – review & editing,

Conceptualization. DJ: Formal Analysis, Writing – review & editing, Visualization. JM-J: Data curation, Formal Analysis, Writing – review & editing, Visualization. PG-S: Formal Analysis, Writing – review & editing, Visualization.

Funding

The author(s) declare financial support was received for the research, authorship, and/or publication of this article.

AJ-M received a funding. “23 Convocatoria Interna de Apoyo a Proyectos de Investigación de la Universidad Autónoma de Baja California, México” (Register Number 431/2/C/38/23). The funders have no role in decision-making or writing of the manuscript.

Acknowledgments

We greatly appreciate the effort made for the undergraduate student Manuel Arturo Celaya Romero who designed and elaborated the [Figure 2](#).

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The author(s) declared that they were an editorial board member of *Frontiers*, at the time of submission. This had no impact on the peer review process and the final decision.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

1. World Health Organization: WHO. Ageing and health. (2022). Available online at: <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health> (Accessed 10 October 2023).
2. Nations U. World Population Ageing 2019 Highlights. United Nations. (2019). Available online at: http://books.google.ie/books?id=-mz8DwAAQBAJ&pg=PP4&dq=978-92-1-148325-3&hl=&cd=1&source=gbs_api
3. Seon Q, Maharaj S, Dookeeram D, Ali K, Seemungal T. Leveraging research, community and collaboration towards robust COVID-19 mental health response in the Caribbean. *Lancet Reg Health Am.* (2023) 19:100440. doi: 10.1016/j.lana.2023.100440
4. Erickson KI, Prakash RS, Voss MW, Chaddock L, Heo S, McLaren M, et al. Brain-derived neurotrophic factor is associated with age-related decline in hippocampal volume. *J Neurosci.* (2010) 30(15):5368–75. doi: 10.1523/JNEUROSCI.6251-09.2010
5. Zhao L, Matloff W, Ning K, Kim H, Dinov ID, Toga AW. Age-related differences in brain morphology and the modifiers in middle-aged and older adults. *Cerebral Cortex.* (2019) 29(10):4169–93. doi: 10.1093/cercor/bhy300
6. Jockwitz C, Mérillat S, Liem F, Oswald J, Amunts K, Jäncke L, et al. Generalizing longitudinal age effects on brain structure—a two-study comparison approach. *Front Hum Neurosci.* (2021) 15:635687. doi: 10.3389/fnhum.2021.635687

7. Mattson MP, Arumugam TV. Hallmarks of brain aging: adaptive and pathological modification by metabolic states. *Cell Metab.* (2018) 27:1176–99. doi: 10.1016/j.cmet.2018.05.011
8. Rajan KB, Weuve J, Barnes LL, McAninch EA, Wilson RS, Evans DA. Population estimate of people with clinical Alzheimer's disease and mild cognitive impairment in the United States (2020–2060). *Alzheimer's Dementia.* (2021) 17(12):1966–75. doi: 10.1002/alz.12362
9. Rezende LFM, Rey-López JP, Matsudo VKR, Luiz ODC. Sedentary behavior and health outcomes among older adults: a systematic review. *BMC Public Health.* (2014) 14(1):333. doi: 10.1186/1471-2458-14-333
10. Erickson KI, Hillman C, Stillman CM, Ballard RM, Bloodgood B, Conroy DE, et al. Physical activity, cognition, and brain outcomes: a review of the 2018 physical activity guidelines. *Med Sci Sports Exerc.* (2019) 51:1242–51. doi: 10.1249/MSS.0000000000001936
11. Gallardo-Gómez D, del Pozo-Cruz J, Noetel M, Álvarez-Barbosa F, Alfonso-Rosa RM, del Pozo Cruz B. Optimal dose and type of exercise to improve cognitive function in older adults: a systematic review and Bayesian model-based network meta-analysis of RCTs. *Ageing Res Rev.* (2022) 76:101591. doi: 10.1016/j.arr.2022.101591
12. Sanders L, Hortobágyi T, La Bastide-van Gemert S, Van Der Zee EA, Van Heuvelen MJG. Dose-response relationship between exercise and cognitive function in older adults with and without cognitive impairment: a systematic review and meta-analysis. *PLoS One.* (2019) 14(1):e0210036. doi: 10.1371/journal.pone.0210036
13. Kelly ME, Loughrey D, Lawlor BA, Robertson IH, Walsh C, Brennan S. The impact of exercise on the cognitive functioning of healthy older adults: a systematic review and meta-analysis. *Ageing Res Rev.* (2014) 16:12–31. doi: 10.1016/j.arr.2014.05.002
14. Smith PJ, Blumenthal JA, Hoffman BM, Cooper H, Strauman TA, Welsh-Bohmer K, et al. Aerobic exercise and neurocognitive performance: a meta-analytic review of randomized controlled trials. *Psychosom Med.* (2010) 72(3):239–52. doi: 10.1097/PSY.0b013e3181d14633
15. Van Uffelen J, Paw MJMCA, Hopman-Rock M, Van Mechelen W. The effects of exercise on cognition in older adults with and without cognitive decline: a systematic review. *Clin J Sport Med.* (2008) 18(6):486–500. doi: 10.1097/jsm.0b013e3181845f0b
16. Colcombe SJ, Kramer AF. Fitness effects on the cognitive function of older adults. *Psychol Sci.* (2003) 14(2):125–30. doi: 10.1111/1467-9280.t01-1-01430
17. Sala G, Gobet F. Cognitive training does not enhance general cognition. *Trends Cogn Sci (Regul Ed).* (2019) 23:9–20. doi: 10.1016/j.tics.2018.10.004
18. Huang Z, Zhang Y, Zhou R, Yang L, Hong-Ying P. Lactate as potential mediators for exercise-induced positive effects on neuroplasticity and cerebrovascular plasticity. *Front Physiol.* (2021) 12:656455. doi: 10.3389/fphys.2021.656455
19. Yang J, Ruchti E, Petit JM, Jourdain P, Grenningloh G, Allaman I, et al. Lactate promotes plasticity gene expression by potentiating NMDA signaling in neurons. *Proc Natl Acad Sci U S A.* (2014) 111(33):12228–33. doi: 10.1073/pnas.1322912111
20. Tomoto T, Tarumi T, Chen JN, Hynan LS, Munro Cullum C, Zhang R. One-year aerobic exercise altered cerebral vasomotor reactivity in mild cognitive impairment. *J Appl Physiol.* (2021) 131(1):119–30. doi: 10.1152/jappphysiol.00158.2021
21. Thomas BP, Tarumi T, Sheng M, Tseng B, Womack KB, Munro Cullum C, et al. Brain perfusion change in patients with mild cognitive impairment after 12 months of aerobic exercise training. *J Alzheimer's Dis.* (2020) 75(2):617–31. doi: 10.3233/JAD-190977
22. Gaitán JM, Boots EA, Dougherty RJ, Oh JM, Ma Y, Edwards DF, et al. Brain glucose metabolism, cognition, and cardiorespiratory fitness following exercise training in adults at risk for Alzheimer's disease. *Brain Plast.* (2019) 5(1):83–95. doi: 10.3233/BPL-190093
23. Farhani F, Shahrbanian S, Auais M, Hekmatikar AHA, Suzuki K. Effects of aerobic training on brain plasticity in patients with mild cognitive impairment: a systematic review of randomized controlled trials. *Brain Sci.* (2022) 12(6):732. doi: 10.3390/brainsci12060732
24. Huang X, Zhao X, Li B, Cai Y, Zhang S, Wan Q, et al. Comparative efficacy of various exercise interventions on cognitive function in patients with mild cognitive impairment or dementia: a systematic review and network meta-analysis. *J Sport Health Sci.* (2022) 11:212–23. doi: 10.1016/j.jshs.2021.05.003
25. Venegas-Sanabria LC, Caverro-Redondo I, Martínez-Vizcaíno V, Cano-Gutiérrez CA, Álvarez-Bueno C. Effect of multicomponent exercise in cognitive impairment: a systematic review and meta-analysis. *BMC Geriatr.* (2022) 22(1):617. doi: 10.1186/s12877-022-03302-1
26. Yong L, Liu L, Ding T, Gao Y, Su H, Wang J, et al. Evidence of effect of aerobic exercise on cognitive intervention in older adults with mild cognitive impairment. *Front Psychiatry.* (2021) 12:713671. doi: 10.3389/fpsy.2021.713671
27. Gates N, Singh MAF, Sachdev PS, Valenzuela M. The effect of exercise training on cognitive function in older adults with mild cognitive impairment: a meta-analysis of randomized controlled trials. *American Journal of Geriatric Psychiatry.* (2013) 21(11):1086–97. doi: 10.1016/j.jagp.2013.02.018
28. Zheng G, Xia R, Zhou W, Tao J, Chen L. Aerobic exercise ameliorates cognitive function in older adults with mild cognitive impairment: a systematic review and meta-analysis of randomized controlled trials. *Br J Sports Med.* (2016) 50:1443–50. doi: 10.1136/bjsports-2015-095699
29. Petersen RC, López OL, Armstrong MJ, Getchius TSD, Ganguli M, Gloss D, et al. Practice guideline update summary: mild cognitive impairment. *Neurology.* (2018) 90(3):126–35. doi: 10.1212/wnl.0000000000004826
30. Petersen RC, Lopez O, Armstrong MJ, Getchius TSD, Ganguli M, Gloss D, et al. Practice guideline update summary: mild cognitive impairment report of the guideline development, dissemination, and implementation. *Neurology.* (2018) 90(3):126–35. doi: 10.1212/WNL.0000000000004826
31. Jongsiriyanong S, Limpawattana P. Mild cognitive impairment in clinical practice: a review article. *Am J Alzheimer's Dis Other Dement.* (2018) 33:500–7. doi: 10.1177/1533317518791401
32. Langa KM, Levine DA. The diagnosis and management of mild cognitive impairment: a clinical review. *J Am Med Assoc.* (2014) 312:2551–61. doi: 10.1001/jama.2014.13806
33. Ribeiro FS, Teixeira-Santos AC, Leist AK. The prevalence of mild cognitive impairment in Latin America and the Caribbean: a systematic review and meta-analysis. *Ageing Ment Health.* (2022) 26(9):1710–20. doi: 10.1080/13607863.2021.2003297
34. Mukadam N, Sommerlad A, Huntley J, Livingston G. Population attributable fractions for risk factors for dementia in low-income and middle-income countries: an analysis using cross-sectional survey data. *Lancet Glob Health.* (2019) 7(5):e596–603. doi: 10.1016/S2214-109X(19)30074-9
35. Barber SE, Clegg AP, Young JB. Is there a role for physical activity in preventing cognitive decline in people with mild cognitive impairment? *Age Ageing.* (2012) 41(1):5–8. doi: 10.1093/ageing/afr138
36. Turner DT, Hu MX, Generaal E, Bos D, Ikram MK, Heshmatollah A, et al. Physical exercise interventions targeting cognitive functioning and the cognitive domains in nondementia samples: a systematic review of meta-analyses. *J Geriatr Psychiatry Neurol.* (2021) 34:91–101. doi: 10.1177/0891988720915523
37. Northey JM, Cherbuin N, Pumpa KL, Smeed DJ, Rattray B. Exercise interventions for cognitive function in adults older than 50: a systematic review with meta-analysis. *Br J Sports Med.* (2018) 52:154–60. doi: 10.1136/bjsports-2016-096587
38. Young J, Angevaren M, Rusted J, Tabet N. Aerobic exercise to improve cognitive function in older people without known cognitive impairment. *Cochrane Database Syst Rev.* (2015) 2015(4):CD005381. doi: 10.1002/14651858.cd005381.pub4
39. Friedman NP, Miyake A. Unity and diversity of executive functions: individual differences as a window on cognitive structure. *Cortex.* (2017) 86:186–204. doi: 10.1016/j.cortex.2016.04.023
40. Chan RCK, Shum D, Touloupoulou T, Chen EYH. Assessment of executive functions: review of instruments and identification of critical issues. *Arch Clin Neuropsychol.* (2008) 23(2):201–16. doi: 10.1016/j.acn.2007.08.010
41. Guiney H, Machado L. Benefits of regular aerobic exercise for executive functioning in healthy populations. *Psychon Bull Rev.* (2013) 20:73–86. doi: 10.3758/s13423-012-0345-4
42. Etnier JL, Chang YK. The effect of physical activity on executive function: a brief commentary on definitions, measurement issues, and the current state of the literature. *J Sport Exerc Psychol.* (2009) 31(4):469–83. doi: 10.1123/jsep.31.4.469
43. Niendam TA, Laird AR, Ray KL, Dean YM, Glahn DC, Carter CS. Meta-analytic evidence for a superordinate cognitive control network subserving diverse executive functions. *Cogn Affect Behav Neurosci.* (2012) 12(2):241–68. doi: 10.3758/s13415-011-0083-5
44. Li H, Su W, Dang H, Han K, Lu H, Yue S, et al. Exercise training for mild cognitive impairment adults older than 60: a systematic review and meta-analysis. *J Alzheimer's Dis.* (2022) 88:1263–78. doi: 10.3233/JAD-220243
45. Anderson ND. State of the science on mild cognitive impairment (MCI). *CNS Spectr.* (2019) 24:78–87. doi: 10.1017/S1092852918001347
46. Petersen RC, Caracciolo B, Brayne C, Gauthier S, Jelic V, Fratiglioni L. Mild cognitive impairment: a concept in evolution. *J Intern Med.* (2014) 275(3):214–28. doi: 10.1111/joim.12190
47. Haroutunian V, Hoffman LB, Been MS. Is there a neuropathology difference between mild cognitive impairment and dementia? *Dialogues Clin Neurosci.* (2009) 11(2):171–9. doi: 10.31887/dcn.2009.11.2/haroutunian
48. Diaz-Venegas C, Samper-Ternent R, Michaels-Obregón A, Wong R. The effect of educational attainment on cognition of older adults: results from the Mexican health and aging study 2001 and 2012. *Ageing Ment Health.* (2019) 23(11):1586–94. doi: 10.1080/13607863.2018.1501663
49. Travassos GF, Coelho AB, Arends-Kuenning MP. The elderly in Brazil: demographic transition, profile, and socioeconomic condition. *Rev Bras Estud Popul.* (2020) 37:1–27. doi: 10.20947/S0102-3098a0129
50. Monteverde M, Peláez E, Celton D. Población y Salud en Mesoamérica Población y Salud en Mesoamérica Level of Education and Disability among the Elderly People from Buenos Aires Level of Education and Disability among the Elderly People from Buenos Aires Nivel de educación y discapacidad entre los ancianos de Buenos Aires. Available online at: <http://ccp.ucr.ac.cr/revista/>

51. Rueda-Salazar S, Spijker J, Devolder D, Albala C. The contribution of social participation to differences in life expectancy and healthy years among the older population: a comparison between Chile, Costa Rica and Spain. *PLoS One*. (2021) 16(3):e0248179. doi: 10.1371/journal.pone.0248179
52. Mathers CD, Stevens GA, Boerma T, White RA, Tobias MI. Causes of international increases in older age life expectancy. *Lancet*. (2015) 385(9967):540–8. doi: 10.1016/S0140-6736(14)60569-9
53. Alley D, Suthers K, Crimmins E. Education and cognitive decline in older Americans: results from the AHEAD sample. *Res Aging*. (2007) 29(1):73–94. doi: 10.1177/0164027506294245
54. Williams BD, Pendleton N, Chandola T. Does the association between cognition and education differ between older adults with gradual or rapid trajectories of cognitive decline? *Aging Neuropsychol Cogn*. (2022) 29(4):666–86. doi: 10.1080/13825585.2021.1889958
55. Lövdén M, Fratiglioni L, Glymour MM, Lindenberg U, Tucker-Drob EM. Education and cognitive functioning across the life span. *Psychol Sci Public Interest*. (2020) 21(1):6–41. doi: 10.1177/1529100620920576
56. Jansen MG, Geerligs L, Claassen JAHR, Overdorp EJ, Brazil IA, Kessels RPC, et al. Positive effects of education on cognitive functioning depend on clinical status and neuropathological severity. *Front Hum Neurosci*. (2021) 15:723728. doi: 10.3389/fnhum.2021.723728
57. World Bank. *Global Economic Prospects, June 2023*. Washington, DC: World Bank (2023). doi: 10.1596/978-1-4648-1951-3.
58. FAO, IFAD, UNICEF, WFP, WHO. The state of food security and nutrition in the world 2023. In: eBooks. FAO, IFAD, UNICEF, WFP, WHO (2023). <https://doi.org/10.4060/cc3017en> (Accessed 8 October 2023).
59. Rentería I, García-Suárez PC, Moncada-Jiménez J, Machado-Parra JP, De Moura Mello Antunes B, Lira FS, et al. Unhealthy dieting during the COVID-19 pandemic: an opinion regarding the harmful effects on brain health. *Front Nutr*. (2022) 9:876112. doi: 10.3389/fnut.2022.876112
60. Ximenes-Da-Silva A, Guedes RCA. Editorial: nutrients, neurotransmitters and brain energetics. *Front Neurosci*. (2020) 14. doi: 10.3389/fnins.2020.568937
61. Song K, Li Y, Zhang H, An N, Wei Y, Wang L, et al. Oxidative stress-mediated blood-brain barrier (BBB) disruption in neurological diseases. *Oxid Med Cell Longevity*. (2020) 2020:568937. doi: 10.1155/2020/4356386
62. Banks WA, Rhea EM. The blood-brain barrier, oxidative stress, and insulin resistance. *Antioxidants*. (2021) 10(11):1695. doi: 10.3390/antiox10111695
63. Satō H, Tsukamoto-Yasui M, Takado Y, Kawasaki N, Matsunaga K, Ueno S, et al. Protein deficiency-induced behavioral abnormalities and neurotransmitter loss in aged mice are ameliorated by essential amino acids. *Front Nutr*. (2020) 7:23. doi: 10.3389/fnut.2020.00023