



# Exploring Potential Regulatory Anesthetic Drugs Based on RNA Binding Protein and Constructing CESC Prognosis Model: A Study Based on TCGA Database

Ying Zheng, Xiao Wen Meng and Jian Ping Yang\*

Department of Anesthesiology, First Affiliated Hospital of Soochow University, Suzhou, China

## OPEN ACCESS

### Edited by:

Oscar Díaz-Cambronero,  
La Fe Hospital, Spain

### Reviewed by:

Qian Chen,  
Guangxi Medical University Cancer  
Hospital, China  
Zi-jian Zhang,  
Central South University, China

### \*Correspondence:

Jian Ping Yang  
szyangjp@126.com

### Specialty section:

This article was submitted to  
Surgical Oncology,  
a section of the journal  
Frontiers in Surgery

Received: 27 November 2021

Accepted: 10 March 2022

Published: 05 April 2022

### Citation:

Zheng Y, Meng XW and Yang JP  
(2022) Exploring Potential Regulatory  
Anesthetic Drugs Based on RNA  
Binding Protein and Constructing  
CESC Prognosis Model: A Study  
Based on TCGA Database.  
Front. Surg. 9:823566.  
doi: 10.3389/fsurg.2022.823566

**Objective:** To investigate the differential expression of RBPs in cervical squamous cell carcinoma (CESC), analyze the regulatory effect of narcotic drugs on RBPs, and establish the prognostic risk model of CESC patients.

**Methods:** RNA-SEQ data and clinical case data of cancer and normal samples from CESC patients were obtained from the Cancer Genome Atlas (TCGA) database and Genotype-Tissue Expression (GTEx) database. Differentially expressed RBPs were screened by R language and enriched. The CMAP database is used to predict the anesthetic drugs that regulate the differential expression of RBPs. The prognostic risk score model was constructed by COX regression analysis. Risk score of each CESC patient was calculated and divided into high-risk group and low-risk group according to the median risk score. The prediction efficiency of prognostic risk model was evaluated by Kaplan-Meier (KM) analysis and receiver operating characteristic (ROC) curve, and the correlation between prognostic risk model and clinical characteristics was analyzed. Immunohistochemistry was used to detect the expression of RNASEH2A and HENMT1 in tissues.

**Results:** There were 65 differentially expressed RBPs in CESC. Five anesthetics, including benzocaine, procaine, pentoxifyverine, and tetracaine were obtained to regulate RBPs. Survival analysis showed that seven genes were related to the prognosis of patients, and the CESC risk score model was constructed by COX regression. The risk score can be used as an independent prognostic factor. RNASEH2A and HENMT1 are up-regulated in tumors, which can effectively distinguish normal tissues from tumor tissues.

**Conclusion:** It is found that different anesthetic drugs have different regulatory effects on the differential expression of RBPs. Based on the differentially expressed RBPs, the prognostic risk score model of CESC patients was constructed. To provide ideas for the formulation of individualized precise anesthesia scheme and cancer pain analgesia scheme, which is helpful to improve the perioperative survival rate of cancer patients.


**Keywords:** cervical squamous cell carcinoma (CESC), bioinformatics, narcotic drugs, predictors, risk score


## INTRODUCTION

Cervical cancer is the fourth most common cancer among women in the world, with a high mortality rate among women in developing countries (1). As the most common tissue type of cervical cancer, cervical squamous cell carcinoma (CESC) is a serious threat to women's health, causing about 273,200 death every year (2). In recent years, with the development of cancer screening and various treatment methods such as surgery, radiotherapy and chemotherapy, the clinical prognosis of CESC has been improved to some extent. However, due to the lack of effective diagnostic methods in the early stage of the disease, the risk of metastasis and recurrence of CESC is still high and the prognosis is poor. More and more evidence shows that the abnormal expression of a variety of genes is involved in the occurrence and development of CESC (3–5). In view of the high incidence rate and high mortality rate of CESC, early detection and risk assessment are particularly important for improving the prognosis of CESC patients. Therefore, it is necessary and urgent to find new biomarkers for diagnosis, prognosis and treatment to improve the survival rate of cervical cancer patients. RNA binding proteins (RBPs) are proteins that interact with many types of RNA, including rRNAs, ncRNAs, snRNAs, miRNAs, mRNAs, tRNAs and snoRNAs. So far, more than 1,500 RBPs genes have been found in the human genome (6). These RBPs play an important role in maintaining the physiological balance of cells, especially in the process of development and stress

response. RBPs can bind to target RNA in a structure or sequence dependent manner to form RNA protein complexes, and regulate mRNA stability, RNA processing, splicing, localization, output and translation at the post transcriptional level (7). In the past decades, many studies have revealed that RBPs are abnormally expressed in tumors, affect the transformation of mRNA to protein, and participate in tumorigenesis (8–10). Among them, only a few RBPs have been deeply studied and found to play a key role in human cancer (11–13). The systematic functional study of RBPs will help us to fully understand its role in tumors.

Narcotic drugs are prescription drugs for the treatment of cancer pain. If they are used continuously, they will cause extreme physical and mental dependence, and can only be used in medical treatment and scientific research. Local anesthetics are commonly used for postoperative analgesia and local anesthesia at the surgical site of cancer patients. Local anesthesia includes intestinal nerve block, local infiltration anesthesia, surface anesthesia and so on. Studies have found that local anesthesia can reduce the stress response after surgery and reduce the inhibitory effect of stress response on the immune system (14). Local anesthesia can reduce the dosage of opiates, reduce the inhibition of opiate analgesics on the immune system, and play a certain role in tumor recurrence and metastasis. At the same time, intestinal nerve block combined with propofol can reduce interleukin (IL) 1  $\beta$ / IL-8, increase IL-10. IL-1  $\beta$ / IL-8 is considered to be a cytokine promoting tumor formation, and IL-10 is a cytokine inhibiting tumor formation (15). Local





anesthetics are commonly used in the clinic. Local anesthesia is also a commonly used anesthesia technology in clinic. Some narcotic drugs inhibit tumor growth, invasion and metastasis. Other narcotic drugs promote tumor growth, invasion and metastasis. Their mechanism may be related to regulating the immune ability of the body to the tumor. So choosing different anesthetic drugs in different preoperative periods may have different effects on tumor recurrence and invasion, and directly affect the prognosis of surgical patients. Therefore, the effects of narcotic drugs on tumors and their related mechanisms need to be further studied and discussed.

Based on the above, the RNA sequencing and clinicopathological data of CESC were downloaded from the Cancer Genome Atlas (TCGA) database and Genotype-Tissue Expression (GTEx) database. Subsequently, abnormally expressed RBPs between CESC and normal cervical tissues were identified by high-throughput bioinformatics analysis, and their potential functions and molecular mechanisms were systematically explored. This study identified some RBPs that may affect the prognosis of CESC and promoted the understanding of the molecular mechanism of CESC progression. These RBPs may provide potential biomarkers for diagnosis and prognosis.


## MATERIALS AND METHODS

### Data Download and Processing

Three hundred and nine human cervical cancer gene expression samples were downloaded from TCGA database (<https://portal.gdc.cancer.gov/>), including 3 normal samples and 306 tumor samples, and the corresponding clinical information was provided. The data of the additional 19 normal tissue samples were from GTEx database (<https://gtexportal.org/home/datasets>). RBPs were collected by Merkle et al. (16) and a total of 1542 RBP genes were obtained (**Supplementary Table 1**). The original data were preprocessed with limma software package, and the differentially expressed RBPs were included with error detection rate  $(FDR) < 0.01$  and  $|\log_{2}FC$  (foldchange)  $> 2$ . Differentially expressed RBPs were submitted to the STRING database to identify protein-protein interaction information.

### GO Enrichment and KEGG Pathway Analysis

The biological functions of these differentially expressed RBPs were comprehensively detected by Gene Ontology (GO) enrichment and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis. GO analysis terms include cellular



**FIGURE 3 |** Functional enrichment analysis of differentially expressed RBPs. **(A)** GO analysis divided RBPs into three functional groups: biological process (BP), cell composition (CC) and molecular function (MF). **(B)** Distribution of RBPs in the first 10 GO enrichment functions. **(C)** KEGG analysis of differentially expressed RBPs.

components (CC), molecular functions (MF), and biological processes (BP). R packages such as clusterProfiler and pathview are used for GO and KEGG pathway enrichment analysis. The difference was statistically significant ( $p.adjust < 0.05$ ).

### Exploration of Anesthetic Drugs With Potential Regulatory Effect


Connectivity map (CMAP) database was used to find anesthetic drugs that regulate the differential expression of RBPs. The database can use computer simulation methods to predict potential drugs that may induce or reverse biological states encoded by gene expression characteristics. Differentially expressed RBPs were classified into up-regulated group and down-regulated group and uploaded to CMAP database. The negative correlation score indicates that the drug inhibits the

expression of up-regulated genes and promotes the expression of down-regulated genes, which may reverse the cancer process.

### Screening of Prognostic Genes of RBPs and Construction of Prognostic Model

According to the amount of single gene expression, the median gene expression was used as the grouping method. Prognostic differences of different groups were analyzed by Kaplan-Meier survival analysis. *P*-value and hazard ratio (HR) with 95% confidence interval (CI) were obtained by log-rank test and univariate COX proportional hazards regression. Then, based on the differentially expressed RBPs related to prognosis, the risk score model was constructed by multivariate COX regression analysis, and the risk score was calculated. Calculation formula of prognosis model:  $riskscore = b1 \times Exp1 + b2 \times Exp2 +$






**FIGURE 5 |** CESC patient risk model based on prognosis related RBPs (A) Survival analysis was used to screen prognosis related RBP. (B) Multivariate prognostic analysis of seven prognosis related RBPs (EXO1, HENMT1, RNASEH2A, MRPL47, ZFR2, MRPS24 and NOVA1). (C) Calculation formula of comprehensive risk score of four genes. Exp stands for gene expression.

nodes of cancer and adjacent tissues were complete. Among 150 patients with cervical cancer, the results of MKI67 immunohistochemistry were collected from 115 patients. *T*-test was used to analyze the difference of gene expression between cancer and adjacent cancer. Correlation test is used to reflect the linear correlation between the expression of two genes. ROC curve was used to analyze the diagnostic effect of gene expression on cervical cancer.

## RESULTS

### RBPs in Differentially Expressed CESC Tissues Were Screened

The flow of this study is shown in **Figure 1**. The data obtained from TCGA and GTEx databases are processed by Perl and R language, and 1984 DEGs (**Figure 2A**) are obtained by “limma” package analysis, including 65 RBPs (**Figure 2B**). RBPs



were collected by Merkley et al. (16) and a total of 1542 RBPs genes were obtained (Supplementary Table 1). STRING database analysis showed that there was a relationship of protein interaction among 49 RBPs (Figure 2C).

### Expression of Differentially Expressed RBPs and Enrichment Analysis of Go and KEGG Pathways


In order to study the function and mechanism of the identified RBPs, these differentially expressed RBPs were enriched and analyzed. The GO analysis results were included in the analysis with  $P < 0.05$  as the standard, and the results were divided into BP, CC and MF groups (Figure 3A). The first 10 significant GO analyses showed that RBPs were mainly involved in ribosomal subunit, ribosome, large ribosomal subunit, structural constituent of ribosome, RNA splicing, RNA catabolic process, cytoplasmic translation, regulation of translation, double-stranded RNA binding and catalytic activity acting on RNA (Figure 3B). KEGG signaling pathway is mainly enriched in Hepatitis C, Spliceosome and Ribosome (Figure 3C).

### Potential Small Molecule Drug Screening

Among these highly significantly related molecules, benzocaine, procaine, Pentoxifyverine and tetracaine are narcotic drugs (Figure 4A). Procaine and Pentoxifyverine are negatively correlated with RBPs gene expression and have potential therapeutic effects on CESC (Figure 4B).

### Construct a Prognostic RBP Prediction Model

For the survival analysis of 65 differentially expressed RBPs, the  $P$ -value and hazard ratio (HR) with 95% confidence interval (CI) were obtained by logrank test and univariate COX proportional hazards regression (Figure 5A). According to the results of survival analysis, seven RBPs (EXO1, HENMT1, RNASEH2A, MRPL47, ZFR2, MRPS24 and NOVA1) were related to the prognosis of patients. Subsequently, seven prognostic RBPs were analyzed by multiple COX regression, in which EXO1, HENMT1, RNASEH2A and MRPS24 can be used as independent predictors of CESC prognosis (Figure 5B). Then, the prognostic risk model of CESC was constructed with the



**FIGURE 7** | Nomogram can predict the 1,3, and 5 year overall survival of patients with CESC.

above four genes. The risk score of each sample was calculated according to the risk coefficient and the expression of 4 RBPs (Figure 5C).

### Risk Model Performance Evaluation


According to the risk score formula, the risk score of 306 CESC patients was calculated, and the median score was taken as the cut-off value. The patients were divided into high-risk group and the low-risk group, with 153 cases in each group (Figure 6A). The distribution of survival time shows that the number of deaths of CESC patients in the high-risk group is more than that in the low-risk group, and the patients with shorter overall survival (OS) in the high-risk group are more than those in the low-risk group (Figure 6C). Kaplan-Meier survival analysis was conducted for patients with high-risk and low-risk groups. The survival rate of low-risk group was significantly higher than that of high-risk group (Figure 6B). The results of ROC curve show that the AUC as the prediction efficiency of risk score for 1-year, 3-year and 5-year prognosis of patients are 0.786, 0.727, and 0.722 respectively, indicating that the model has a certain ability to predict the prognosis of CESC patients (Figure 6D). The nomogram provides a graphical representation of each factor. The prognostic risk of a single patient can be calculated from the points associated with each risk factor, which can be used to predict the 1, 3, and 5 year overall survival of CESC patients (Figure 7).

### Correlation Between Risk Score and Clinical Factors

Data of TCGA-CESC data set were used to further study the correlation between patients' risk score and clinical factors in the model. Chi square test showed that there were differences in tumor grade distribution between high-risk group and low-risk group (Figure 8A). To assess whether the risk score was independent of other clinical variables, COX univariate and multivariate analyses were performed in 182 patients with complete clinical data. Univariate COX analysis showed that the T, N stages and riskScore of CESC patients were related to the prognosis (Figure 8B). Further multivariate COX regression analysis showed that riskScore, T stage and N stage were independent factors affecting OS in patients with CESC (Figure 8C).

### Validation of Prognostic Genes


The expression data analysis of the database showed that there were significant differences in the expression of seven RBPs between CESC and normal tissues, including the up-regulated expression of five genes (HENMT1, RNASEH2A, EXO1, MRPL47 and ZFR2) and the down-regulated expression of two genes (MRPS24 and NOVA1) (Figures 9A-G). In addition, the expression of RNASEH2A was positively correlated with the expression of EXO1, ZFR2, HENMT1, MRPL47, and MRPS24. The expression of HENMT1 was positively correlated with the expression of EOX1 and ZFR2 (Figure 9H). Finally, protein expression levels of RNASEH2A and HENMT1



in cervical cancer and adjacent tissues were verified by immunohistochemistry. It can be seen that the expression of RNASEH2A and HENMT1 in tumor tissues is significantly higher than that in adjacent tissues (Figures 10A,C). In the RNASEH2A immunohistochemical expression micro array, after deleting the tissue points shed during the experiment, the expression of RNASEH2A in the remaining 120 pairs of cancer and adjacent tissues was statistically analyzed (Supplementary Figure 1). Statistical analysis showed that the expression of RNASEH2A in cervical cancer was higher than that in adjacent tissues (Figure 10B). In the HENMT1 immunohistochemical expression micro array, after deleting the tissue points shed during the experiment, the expression of HENMT1 in the remaining 93 pairs of cancer and adjacent tissues was statistically analyzed (Supplementary Figure 2). Statistical analysis showed that the expression of HENMT1 in cervical cancer was higher than that in adjacent tissues (Figure 10D).

### Correlation Between RNASEH2A and HENMT1 and Clinical Markers, and Evaluation of Diagnostic Effect

GEPIA database analysis showed that RNASEH2A and HENMT1 were positively correlated with the expression of tumor marker (MKI67) (Figures 11A,D). Clinical immunohistochemical data showed that the expression of tumor marker (MKI67) in cancer tissues was stronger than that in normal tissues (Figures 11B,E). At the same time, RNASEH2A and HENMT1 were positively correlated with the expression of clinical markers (MKI67) in cancer tissues (Figures 11C,F). ROC curve analysis shows that the areas under RNASEH2A, HENMT1 and MKI67 curves are 0.92, 0.946, and 0.925, respectively (Figures 11G-I). However, RNASEH2A, HENMT1 combined with clinical markers (MKI67) calculated the largest area under the ROC curve, which was 0.992 (Figure 11J).




## DISCUSSION

CESC is one of the most common malignant tumors in women worldwide, and has a high incidence rate (16). The risk of recurrence and metastasis of CESC is high. Early diagnosis and treatment are very important to improve the prognosis of CESC patients. Therefore, it is urgent to explore new potential biomarkers that can be used for early diagnosis, targeted therapy or prognosis evaluation to improve the prognosis of CESC patients. Micro array analysis is a high-throughput technology, which can detect the expression level of thousands of genes at the same time. Nowadays, abnormal gene expression is considered to be one of the factors in the occurrence and development of CESC, and more and more studies show that some deregulated genes in CESC may become candidate biomarkers for diagnosis and prognosis (17). RBPs are proteins that can bind to a variety of RNAs and can be stably expressed in cells. Their main role is RNA processing, such as mRNA splicing and translation regulation.

In the past decades, it has been found that RBPs are closely related to the occurrence and development of many tumors (18). At present, narcotic drugs are prescription drugs for the treatment of cancer pain. Some narcotic drugs inhibit tumor growth, invasion and metastasis. Other narcotic drugs promote tumor growth, invasion and metastasis. Their mechanism may be related to regulating the immune ability of the body to the tumor. So choosing different anesthetic drugs in different preoperative periods may have different effects on tumor recurrence and invasion, and directly affect the prognosis of surgical patients. Therefore, this study analyzed the relationship between RBPs and narcotic drugs in CESC.

In this study, the gene expression profile of CESC was analyzed by bioinformatics to explore its molecular mechanism and identify important molecules that may be used as CESC biomarkers and therapeutic targets. In this study, we downloaded the gene expression profile data set of CESC from TCGA database, deeply analyzed it by bioinformatics method, and




obtained the RBPs between CESC tissue and normal tissue. The results showed that 65 RBPs were identified between CESC and normal tissues. GO functional enrichment analysis showed that DEGs were mainly involved in ribosomal subunit, ribosome, large ribosomal subunit, structural constituent of ribosome, RNA splicing, RNA catabolic process, cytoplasmic translation, regulation of translation, double-stranded RNA binding and catalytic activity acting on RNA. In addition, enrichment results of KEGG pathway showed that the enriched pathways mainly involved Hepatitis C, Spliceosome and Ribosome. Post transcriptional regulation of RNA stability is an important step in the process of gene expression. RBPs can interact with RNA to form RNA protein complexes, so as to increase the stability of target mRNA, promote gene expression and play a key role in the progress of various diseases. The above information shows that our data mining results are consistent with the existing research results. In this study, local anesthetics (benzocaine, procaine, pentoxifyverine and tetracaine) with regulatory function to 65 differentially expressed RBPs were analyzed. Among them, the regulation trend of benzocaine and tetracaine on genes is the same as that of RBPs. Although drugs have analgesic

effect, they may lead to disease deterioration by promoting cell biological function. Procaine and Pentoxifyverine may inhibit tumor progression on the basis of anesthesia and analgesia. Its effect on the tumor and its related mechanism needs to be further studied and discussed.

Survival analysis was used to screen the genes related to the prognosis of CESC patients in 65 differentially expressed RBPs. The results showed that seven RBPs (HENMT1, RNASEH2A, EXO1, MRPL47, ZFR2, NOVA1, and MRPS24) were related to the prognosis of patients. Among them, four genes (HENMT1, RNASEH2A, EXO1, and MRPS24) are independent predictors of the prognosis of CESC patients. Based on the above four genes, a CESC prognostic risk score model was constructed to further improve the reliability of the prediction results.

Hen methyltransferase 1 (HENMT1) is a methyltransferase. It is a kind of RNA composed of 24–30 nucleotides. It is produced by Dicer independent mechanism and mainly comes from transposons and other repeat elements (19, 20). The expression of HENMT1 in ovarian cancer is increased with the increase of tumor grade, which was related to the degree of malignancy (21). In this study, the expression of HENMT1 in cervical cancer



patients is up-regulated, and its high expression indicates a better prognosis. As a low-risk gene, HENMT1 may be a marker for predicting the prognosis of cervical cancer patients.

Ribonuclease H2 subunit A (RNASEH2A) is an endonuclease that specifically degrades RNA. It can remove lag strand Okazaki fragment RNA primers by mediating the process of DNA replication. RNASEH2A participates in the occurrence of human glioma by promoting glioma cell proliferation and inhibiting apoptosis (22). Over expression of RNASEH2A is positively correlated with chemoresistance of breast cancer cells (23). RNASEH2A was highly expressed in lung cells, and its knockdown inhibited the proliferation of lung cells and induced apoptosis (24). Similarly, we found that RNASEH2A is highly expressed in cervical cancer and participates in RNA catabolic process, which is expected to become a molecular diagnostic marker and therapeutic target of cervical cancer.

Exonuclease 1 (EXO1) has both 5'-3' exonuclease and 5' structure specific endonuclease activities. It plays an important role in base mismatch repair, cross injury synthesis, nucleotide

excision repair, DNA double strand break repair, meiotic recombination repair and telomere maintenance (25). It was found that the abnormal expression of EXO1 gene may affect the prognosis, survival and progress of patients with prostate cancer (26). The high expression of EXO1 in breast cancer and lung cancer may promote tumor development (27, 28). We found that EXO1 was highly expressed in cervical cancer and was verified by immunohistochemistry. EXO1 participates in catalytic activity in RNA and is expected to become a molecular diagnostic marker of CESC.

Mitochondrial ribosomal protein L47 (MRPL47) is a member of the MRPs family. Mitochondrial ribosome is composed of a small 28s subunit and a large 39s subunit. MRPL47 encodes a large subunit protein. MRPL47 gene mutation is a new high risk factor of vincristine induced peripheral neuropathy in children with acute lymphoblastic leukemia (29). MRPL47 was reported in square cell carcinoma of head and neck, which was related to the prognosis of patients (30). In this study, compared with normal samples, the expression of MRPL47 in CESC was up-regulated

and involved in the structural components of ribosomes, and its increased expression in CESC showed a good prognosis.

Nova alternative splicing regulator 1 (NOVA1) is an RNA binding protein with a specific sequence. The protein has three KH type domains that can bind to RNA. In recent years, many studies have shown that NOVA1 is involved in regulating the occurrence and development of tumors. Zhang et al. (31) have shown that noval1 plays a carcinogenic role in primary liver cancer. Kim et al. (32) showed that the expression of NOVA1 was significantly down regulated in gastric cancer, and the low expression level of noval1 was closely related to the poor prognosis of patients with gastric cancer. Similarly, NOVA1 expression was down regulated in CESC, and patients with low expression had a poor prognosis. NOVA1 plays a role in RNA splicing and is expected to become a molecular diagnostic marker of CESC.

Mitochondrial ribosomal protein S24 (MRPS24) belongs to the universal ribosomal protein US3 family. At present, the role of MRPS24 in tumorigenesis and development has not been reported. In this study, the expression of MRPS24 was down regulated in CESC, which is the structural component of ribosome and a risk factor for poor prognosis.

Finally, in order to explore the correlation between the markers predicted in this study and clinical tumor markers, the correlation between the experimental results of HENMT1, RNASEH2A and clinical tumor markers was analyzed. It was found that HENMT1 and RNASEH2A were positively correlated with the expression of tumor marker (MKI67). At the same time, compared with the diagnostic efficiency of simple tumor markers, HENMT1 and RNASEH2A combined with clinical markers (MKI67) have higher diagnostic efficiency and reduce the misdiagnosis and missed diagnosis rate of patients to a certain extent.

In conclusion, we used bioinformatics methods to deeply mine the RBPs expression profile data set of CESC. On this basis, we mined narcotic drugs (benzocaine, procaine, pentoxiverine and tetracaine) that regulate RBPs. It was found that procaine and Pentoxiverine are expected to become potential drugs for the treatment of CESC. We screened seven hub genes (HENMT1, RNASEH2A, EXO1, MRPL47, ZFR2, NOVA1 and MRPS24). Among them, the prognostic risk model was constructed based on four independent predictors (HENMT1, RNASEH2A, EXO1 and MRPS24) of the prognosis of CESC patients. The model not only provides new insights into the heterogeneity of CESC, but also has independent predictive value for unconventional

clinicopathological factors. It can provide patients with more accurate prognosis evaluation and individualized diagnosis and treatment.

## DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**, further inquiries can be directed to the corresponding author/s.

## ETHICS STATEMENT

The studies involving human participants were reviewed and approved by the Affiliated Huaian No. 1 People's Hospital of Nanjing Medical University. The patients/participants provided their written informed consent to participate in this study.

## AUTHOR CONTRIBUTIONS

YZ and JY participated in the study design, drafted the manuscript, and revised the manuscript. YZ and XM statistically analyzed the data. All authors read and approved the final manuscript.

## FUNDING

Huai'an Science and Technology support project, Grant No. HAB201927.

## ACKNOWLEDGMENTS

Thanks to tutor Yang Jianping's careful guidance and the experimental support of other relevant personnel in the Department of Anesthesiology.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fsurg.2022.823566/full#supplementary-material>

**Supplementary Table 1** | Information list of 1,542 RBPs genes.

**Supplementary Figure 1** | The expression of RNASEH2A in the remaining 120 pairs of cancer and adjacent tissues.

**Supplementary Figure 2** | The expression of HENMT1 in the remaining 93 pairs of cancer and adjacent tissues.

## REFERENCES

- Small W Jr, Bacon MA, Bajaj A, Chuang LT, Fisher BJ, Harkenrider MM, et al. Cervical cancer: A global health crisis. *Cancer*. (2017) 123:2404–12. doi: 10.1002/cncr.30667
- Siegel RL, Miller KD, Jemal A. Cancer statistics, 2018. *CA Cancer J Clin*. (2018) 68:7–30. doi: 10.3322/caac.21442
- Bahrami A, Hasanzadeh M, Shahidsales S, Farazestanian M, Hassanian SM, Moetamani Ahmadi M, et al. Genetic susceptibility in cervical cancer: From bench to bedside. *J Cell Physiol*. (2018) 233:1929–39. doi: 10.1002/jcp.26019
- Gu X, Guo XK, Chen BH, Gao XJ, Chen F, Liu Q. Prognostic and clinicopathological value of ZIC1 in patients with cervical squamous cell carcinoma. *Oncol Lett*. (2019) 18:6621–7. doi: 10.3892/ol.2019.11007
- Zhao D, Qin W, Zhao C, Long J, Li M. CXCR7, a Prognostic Biomarker in Cervical Squamous Cell Carcinoma, May Be a Screening Index for Treatment Options at Stages IB1 and IIA1. *Cancer Manag Res*. (2019) 11:10287–96. doi: 10.2147/CMAR.S228684
- Gerstberger S, Hafner M, Tuschl T. A census of human RNA-binding proteins. *Nat Rev Genet*. (2014) 15:829–45. doi: 10.1038/nrg3813

7. Masuda K, Kuwano Y. Diverse roles of RNA-binding proteins in cancer traits and their implications in gastrointestinal cancers. *Wiley Interdiscip Rev RNA*. (2019) 10:e1520. doi: 10.1002/wrna.1520
8. Pereira B, Billaud M, Almeida R. RNA-binding proteins in cancer: old players and new actors. *Trends Cancer*. (2017) 3:506–28. doi: 10.1016/j.trecan.2017.05.003
9. Legrand N, Dixon DA, Sobolewski C. AU-rich element-binding proteins in colorectal cancer. *World J Gastrointest Oncol*. (2019) 11:71–90. doi: 10.4251/wjgo.v11.i2.71
10. Chatterji P, Rustgi AK. RNA. Binding proteins in intestinal epithelial biology and colorectal cancer trends. *Mol Med*. (2018) 24:490–506. doi: 10.1016/j.molmed.2018.03.008
11. Zhang H, Wang Y, Dou J, Guo Y, He J, Li L, et al. Acetylation of AGO2 promotes cancer progression by increasing oncogenic miR-19b biogenesis. *Oncogene*. (2019) 38:1410–31. doi: 10.1038/s41388-018-0530-7
12. Jeong HM, Han J, Lee SH, Park HJ, Lee HJ, Choi JS, et al. ESRP1 is overexpressed in ovarian cancer and promotes switching from mesenchymal to epithelial phenotype in ovarian cancer cells. *Oncogenesis*. (2017) 6:e389. doi: 10.1038/oncsis.2017.89
13. Chen H, Liu J, Wang H, Cheng Q, Zhou C, Chen X, et al. Inhibition of RNA-binding protein Musashi-1 suppresses malignant properties and reverses paclitaxel resistance in ovarian carcinoma. *J Cancer*. (2019) 10:1580–92. doi: 10.7150/jca.27352
14. Tonnesen E, Wahlgreen C. Influence of extradural and general anaesthesia on natural killer cell activity and lymphocyte subpopulations in patients undergoing hysterectomy. *Br J Anaesth*. (1988) 60:500–7. doi: 10.1093/bja/60.5.500
15. Deegan CA, Murray D, Doran P, Moriarty DC, Sessler DI, Mascha E, et al. Anesthetic technique and the cytokine and matrix metalloproteinase response to primary breast cancer surgery. *Reg Anesth Pain Med*. (2010) 35:490–5. doi: 10.1097/AAP.0b013e3181ef4d05
16. Merkley MA, Hildebrandt E, Podolsky RH, Arnouk H, Ferris DG, Dynan WS, et al. Large-scale analysis of protein expression changes in human keratinocytes immortalized by human papilloma virus type 16 E6 and E7 oncogenes. *Proteome Sci*. (2009) 7:29. doi: 10.1186/1477-5956-7-29
17. Ouyang F, Liu J, Xia M, Lin C, Wu X, Ye L, et al. GINS2 is a novel prognostic biomarker and promotes tumor progression in early-stage cervical cancer. *Oncol Rep*. (2017) 37:2652–62. doi: 10.3892/or.2017.5573
18. Tang R, Jiang Y, Tan A, Ye J, Xian X, Xie Y, et al. 16S rRNA gene sequencing reveals altered composition of gut microbiota in individuals with kidney stones. *Urolithiasis*. (2018) 46:503–14. doi: 10.1007/s00240-018-1037-y
19. Lim SL, Qu ZP, Kortschak RD, Lawrence DM, Geoghegan J, Hempfling AL, et al. HENMT1 and piRNA stability are required for adult male germ cell transposon repression and to define the spermatogenic program in the mouse. *PLoS Genet*. (2015) 11:e1005620. doi: 10.1371/journal.pgen.1005782
20. Hempfling AL, Lim SL, Adelson DL, Evans J, O'Connor AE, Qu ZP, et al. Expression patterns of HENMT1 and PIWIL1 in human testis: implications for transposon expression. *Reproduction*. (2017) 154:363–74. doi: 10.1530/REP-16-0586
21. Lee E, Lokman NA, Oehler MK, Ricciardelli C, Grutzner F, A. Comprehensive molecular and clinical analysis of the piRNA pathway genes in ovarian cancer. *Cancers (Basel)*. (2020) 13:4. doi: 10.3390/cancers13010004
22. Dai B, Zhang P, Zhang Y, Pan C, Meng G, Xiao X, et al. RNaseH2A is involved in human gliomagenesis through the regulation of cell proliferation and apoptosis. *Oncol Rep*. (2016) 36:173–80. doi: 10.3892/or.2016.4802
23. Shen J, Lin S, Liu L, Wang C. Ribonuclease H2 Subunit A impacts invasiveness and chemoresistance resulting in poor survivability of breast cancer in ER dependent manner. *Am J Transl Res*. (2020) 12:2281–94.
24. Zhang J, Ma D, Kang H, Zhao J, Yang M. Long noncoding RNA LINC01287 promotes proliferation and inhibits apoptosis of lung adenocarcinoma cells via the miR-3529-5p/RNASEH2A axis under the competitive endogenous RNA pattern. *Environ Toxicol*. (2021) 36:2093–104. doi: 10.1002/tox.23325
25. Sertic S, Quadri R, Lazzaro F, Muzi-Falconi M. EXO1: A tightly regulated nuclease. *DNA Repair (Amst)*. (2020) 93:102929. doi: 10.1016/j.dnarep.2020.102929
26. Hua X, Ge S, Chen J, Zhang L, Tai S, Liang C. Effects of RNA binding proteins on the prognosis and malignant progression in prostate cancer. *Front Genet*. (2020) 11:591667. doi: 10.3389/fgene.2020.591667
27. Liu J, Zhang J. Elevated EXO1 expression is associated with breast carcinogenesis and poor prognosis. *Ann Transl Med*. (2021) 9:135. doi: 10.21037/atm-20-7922
28. Zhou CS, Feng MT, Chen X, Gao Y, Chen L, Li LD, Li DH, et al. Exonuclease 1 (EXO1) is a potential prognostic biomarker and correlates with immune infiltrates in lung adenocarcinoma. *Onco Targets Ther*. (2021) 14:1033–48. doi: 10.2147/OTT.S286274
29. Abaji R, Ceppi F, Patel S, Gagné V, Xu CJ, Spinella JF, et al. Genetic risk factors for VIPN in childhood acute lymphoblastic leukemia patients identified using whole-exome sequencing. *Pharmacogenomics*. (2018) 19:1181–93. doi: 10.2217/pgs-2018-0093
30. Hu G, Jiang Q, Liu L, Peng H, Wang Y, Li S, et al. Integrated analysis of RNA-binding proteins associated with the prognosis and immunosuppression in squamous cell carcinoma of head and neck. *Front Genet*. (2021) 11:571403. doi: 10.3389/fgene.2020.571403
31. Zhang YA, Liu HN, Zhu JM, Zhang DY, Shen XZ, Liu TT, et al. binding protein Noval promotes tumor growth *in vivo* and its potential mechanism as an oncogene may due to its interaction with GABAA Receptor- $\gamma$ 2. *J Biomed Sci*. (2016) 23:71. doi: 10.1186/s12929-016-0288-6
32. Kim EK, Yoon SO, Jung WY, Lee H, Kang Y, Jang YJ, et al. Implications of NOVA1 suppression within the microenvironment of gastric cancer: association with immune cell dysregulation. *Gastric Cancer*. (2017) 20:438–47. doi: 10.1007/s10120-016-0623-3

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Zheng, Meng and Yang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.