



# Expression and Immunogenicity of Recombinant African Swine Fever Virus Proteins Using the Semliki Forest Virus

Niran Fang<sup>1,2</sup>, Bin Yang<sup>1</sup>, Ting Xu<sup>2</sup>, Yanpeng Li<sup>3</sup>, Huimin Li<sup>4</sup>, Hanghui Zheng<sup>1</sup>, Aiguo Zhang<sup>5</sup> and Ruiai Chen<sup>1,2,3\*</sup>

<sup>1</sup> Zhaoqing Institute of Biotechnology Co. Ltd., Zhaoqing, China, <sup>2</sup> College of Veterinary Medicine, South China Agricultural University, Guangzhou, China, <sup>3</sup> Zhaoqing DaHuaNong Biology Medicine Co. Ltd., Zhaoqing, China, <sup>4</sup> Jinggangshan University, Jinggangshan, China, <sup>5</sup> Henan University of Animal Husbandry and Economy, Zhengzhou, China

African swine fever virus (ASFV) is a large DNA virus belonging to the Asfarviridae family that damages the immune system of pigs, resulting in the death or slaughter of millions of animals worldwide. Recent modern techniques in ASFV vaccination have highlighted the potential of viral replicon particles (RPs), which can efficiently express foreign proteins and induce robust cellular and humoral immune responses compared with the existing vaccines. In this study, we established a Semliki Forest virus (SFV) vector by producing replication-defective viral particles. This vector was used to deliver RPs expressing ASFV antigens. SFV-RPs expressing ASFV p32 (SFV-p32) and p54 (SFV-p54) were tested in baby hamster kidney (BHK-21) cells. Proteins expression was evaluated via western blotting and indirect immunofluorescence, while immunogenicity was evaluated in BALB/c mice. The resulting RPs exhibited high levels of protein expression and elicited robust humoral and cellular immune responses.

**Keywords:** African swine fever virus, Semliki Forest virus, replication-defective viral particles, immunogenicity, cellular immune response

## OPEN ACCESS

### Edited by:

Heather Lynne Wilson,  
University of Saskatchewan, Canada

### Reviewed by:

Qin Zhao,  
Northwest A&F University, China  
Kenneth Lundstrom,  
PanTherapeutics, Switzerland

### \*Correspondence:

Ruaii Chen  
chensa727@vip.126.com

### Specialty section:

This article was submitted to  
Veterinary Infectious Diseases,  
a section of the journal  
Frontiers in Veterinary Science

**Received:** 05 February 2022

**Accepted:** 04 April 2022

**Published:** 09 May 2022

### Citation:

Fang N, Yang B, Xu T, Li Y, Li H,  
Zheng H, Zhang A and Chen R (2022)  
Expression and Immunogenicity of  
Recombinant African Swine Fever  
Virus Proteins Using the Semliki Forest  
Virus. *Front. Vet. Sci.* 9:870009.  
doi: 10.3389/fvets.2022.870009

## INTRODUCTION

African swine fever virus (ASFV) infection affects domestic pigs and causes an acute, febrile, and highly infectious disease. No effective vaccine against ASFV exists to date, and once the infection is detected, the animals must be slaughtered, imposing a large economic burden on the swine industry. ASFV was first observed in Europe in 1957 and spread in Portugal in 1960 wherefrom the virus became established in the Iberian Peninsula and caused outbreaks across Western Europe, the Caribbean, and Brazil. Although ASFV was almost eradicated outside of Africa by the mid-1990s, in 2007 it was detected in Georgia wherefrom it spread throughout Eastern Asia and Eastern Europe, causing large-scale global economic losses. Before 2018, there were no reported cases of ASFV in China, but the government continued to monitor for the disease. In 2013, China joined the ASFV global alliance to strengthen the prevention and control of the disease. This initiated fundamental research efforts with the ultimate goal of developing effective vaccines against ASFV.

In the earliest ASFV studies, attenuated viruses were found to protect from the disease, but their efficacy in the field was not adequate. In recent years, there has been a rapid drive to develop platforms for the expression of a variety of heterologous proteins in animal cells, and there are

now many methods and vector expression platforms available for the expression of different antigens. Vector expression platforms based on nucleic acids represent important new approaches for vaccine development, and DNA vaccines have the added advantage of expressing heterologous DNA sequences (1, 2). Recent improvements in vector systems have facilitated the development of virus-based mRNA vaccines (2, 3). RNA replicon vaccines are based on a vector platform derived from the Semliki Forest virus (SFV), which can replicate autonomously and induce systemic, cellular, and mucosal immunity (4). Murgia et al. (5) has described an alphavirus vector carrying replicon particle (RP)-30 and RP-54, using Venezuelan equine encephalitis virus (VEEV) rather than SFV, and with Vero cells rather than BHK-21 cells, which helped establish humoral immunity in swine.

cDNA coding for a variety of viral proteins involved in ASFV infection, such as p32 and p54, can be cloned into the SFV vector. The p32 protein, which has a molecular weight of ~30 kDa and is expressed in the early stages of viral infection, has a good level of antigenicity and can induce a robust immune response. A p32-specific blocking antibody inhibits the internalization of the virus, indicating that this viral component is also involved in the viral entry into cells (6). p54 is a type I transmembrane protein that crosses the inner envelope of the virus particle and forms a homodimer via a cysteine bridge (7). p54 helps the virus to bind to target cells (8) and stimulates the host to produce p54-specific antibodies. In ASFV-infected cells, p54 binds to the endoplasmic reticulum and uses its membrane to form the viral particle envelope. A special cross-reaction between p54 and the dynein light chain (DLC8) mediates intracellular transport of the virus, suggesting that p54 plays an important role in viral uptake and processing (9).

In a previous study, an SFV vector coding for foreign proteins, as well as a pSFV-helper1, was used to encode the SFV structural proteins and transcribe them into mRNA *in vitro*, these recombinant RNAs were electroporated into baby hamster kidney (BHK)-21 cells to express high levels of heterologous proteins (10). In the SFV replicon vector, SFV structural genes can be replaced with heterologous genes of interest, generating a self-replicating RNA that can efficiently express the heterologous genes. Replicon RNA supplies the non-structural genes, which contain the packaging signal. The viral structural genes are provided by the helper RNA, which lacks part of the non-structural genes, including the packaging signal. Therefore, only the replicon RNA retains the packaging signal, and it is packaged into RPs through the expression of the structural genes supplied by the helper RNA. The resulting RPs can infect mammalian cells but only undergo one round of replication (10, 11). The alphavirus replicon system has been used to develop effective influenza RP vaccines, which have been evaluated in chickens, pigs, and humans and elicited complete protection against lethal avian influenza (4, 12). Compared to conventional DNA vaccines, the replicon RNA system is self-amplifying, boosting cellular and humoral responses (4). The co-expression of genes encoding Ebola and Lassa virus-specific glycoproteins was used to produce an RP vaccine that protects guinea pigs against Ebola and Lassa viral challenges, indicating that RP vaccines may protect against multiple diseases (13). Another advantage of RP vaccines is that

alphaviruses may infect many different cell lines, indicating that cells, such as Chinese hamster ovary, primary chicken embryo fibroblast, duck embryo fibroblast, or human embryonic kidney (HEK) 293 (or 293T) cells, which can yield different amounts of RPs, may be used in commercial vaccine production (4).

Here we aimed to establish an SFV vector by producing replication-defective viral particles and utilized it to deliver RPs expressing ASFV antigens. SFV-RPs expressing ASFV p32 (SFV-p32) and p54 (SFV-p54) were tested in baby hamster kidney (BHK-21) cells. Proteins expression was evaluated via western blotting and indirect immunofluorescence, while immunogenicity was evaluated in BALB/c mice.

## METHODS AND MATERIALS

### Plasmids, Cells, and Animals


pSFVCs-LacZ and pSFV-helper1 (plasmid numbers #92076 and #92073, respectively) were purchased from Addgene (Watertown, Massachusetts, USA). Fifteen 5-week-old SPF BALB/c mice were purchased from the Animal Experiment Center of Southern Medical University (Guangzhou, People's Republic of China), and raised in individually ventilated cages within the SPF animal house of Zhaoqing DaHuaNong Biology Medicine Co. Ltd. (Zhaoqing, People's Republic of China). The animal study was reviewed and approved by Animal Welfare and Ethical Censor Committee.

### Construction of SFV Vector Plasmids

On an SFV vector platform, using the pSFVCs-LacZ containing the non-structural genome of SFV (Figure 1). The LacZ-encoding gene was removed using a BamHI restriction endonuclease to insert a heterologous gene into pSFVCs-LacZ. Enhanced green fluorescent protein (EGFP)- and ASFV p32- or p54-encoding sequences were inserted into the vector plasmid to establish pSFVCs-EGFP, pSFV-EGFP, pSFV-p32, and pSFV-p54 (Figure 1) based on GenBank sequences for p32 and p54 (accession no. NP\_042786 and NP\_042818). The above heterologous proteins were inserted downstream of the 26S promoter.

### SFVCs-RP Generation

To obtain SFVCs-RP, the SFV expression vector plasmids described above were linearized using SpeI, then transcribed *in vitro* using the mMACHINE™ SP6 (Thermo Fisher Scientific, Waltham, MA, USA). pSFVCs-LacZ RNA and pSFVCs-EGFP RNA were electroporated into BHK-21 cells with pSFV-helper1 RNA at 100 V for 25 ms, respectively. BHK-21 cells were cultured in 1% fetal bovine serum (FBS)-containing Dulbecco's Modified Eagle Medium (Gibco) at 37°C until the cells adhered, then the medium was changed to serum-free medium. The viral particles were collected from the supernatant by centrifugation 36 h after co-electroporation, then filtered through a 0.22 μm Millex-GP filter to generate SFVCs-LacZ and SFVCs-EGFP RPs. β-galactosidase was detected using *in situ* β-galactosidase Staining Kit (Beyotime, Shanghai, China), and EGFP was detected using a fluorescence microscope.



## SFV-RP Generation

To avoid interference from the fusion capsid protein during the specific expression of the target protein in the follow-up animal experiments, we removed the viral capsid protein and constructed the replicon plasmid pSFV-EGFP via homologous recombination using the pEASY<sup>®</sup>-Basic Seamless Cloning and Assembly Kit (TransGen Biotech, Beijing, China). SFV-EGFP RP was then obtained using the above-described method for SFVCs-RP. Fluorescence signal intensity was compared with SFVCs-RPs 24 h after infection.

## ASFV SFV-RP Generation

To develop the plasmids expressing ASFV p32 and p54, we constructed the replicon plasmids pSFV-p32 and pSFV-p54 via homologous recombination using the pEASY<sup>®</sup>-Basic Seamless Cloning and Assembly Kit and pSFV-EGFP as described above.

pSFV-p32 and pSFV-p54 were linearized using SpeI and SpaI, respectively, and transcribed *in vitro* using the mMACHINE<sup>™</sup> SP6 (Thermo Fisher Scientific). They were then electroporated into BHK-21 cells with the helper RNA at 100 V for 25 ms. SFV-p32 and SFV-p54 RPs were collected using the same method described above.

## ASFV SFV-RP Viral Titers and Detection of Infected BHK-21 Cells

The viral supernatant was collected by centrifugation, every 6 h starting from 24 h after the replicon RNA electroporation into BHK-21 cells with helper RNA. Then, 100  $\mu$ L dilutions of viral samples were added to 96-well plates containing BHK-21 cell monolayers. Following 60 h incubation at 37°C under 5% CO<sub>2</sub>, viruses were detected based on the cytopathogenic effect (CPE) and immunofluorescence (IFA). ASFV SFV-RP titration was calculated using the Spearman-Kärber method.

## IFA

SFV-p32 and SFV-p54 viral particles were collected every 6 h starting from 24 h after co-electroporation. Cells were cultured in plates with particles for 60 h and washed with phosphate-buffered saline containing Tween (PBST) three times, fixed with formaldehyde for 10 min at 4°C, then washed with PBST and permeabilized with 0.1% Triton X-100. They were then incubated for 2 h at 37°C with 5% bovine serum albumin (BSA), followed by washing and 1 h incubation at 37°C with an anti-ASFV serum antibody (Zhaoqing DaHuaNong Biology Medicine Co. Ltd., Zhaoqing, People's Republic of China) diluted at 1:2500. The samples were then washed again and incubated for 1 h at 37°C with IFTC-conjugated AffiniPure Goat Anti-Swine IgG (H + L) (Jackson ImmunoResearch) diluted at 1:600 as the secondary antibody. They were then washed three times and observed under a fluorescence microscope.


## Western Blotting

BHK-21 cells infected with SFV-p32 and SFV-p54 (RPs) for 36 h were collected and lysed on ice for 5 min. The lysate was mixed with loading buffer (5 $\times$ ) and boiled for 10 min. The mixture was loaded onto 12% polyacrylamide gels, electrophoresed for 120 min at 80 V, then transferred to a polyvinylidene difluoride membrane via electroblotting for 90 min. The membranes were blocked with 5% BSA for 1 h at room temperature, then washed three times with PBST for 10 min each. They were incubated overnight at 4°C with an anti-ASFV serum antibody diluted at 1:1,000, then washed and incubated for 1 h at 37°C with a secondary goat anti-swine IgG (H+L) conjugated with horseradish peroxidase (ABclonal) diluted at 1:5,000. Finally, the membranes were washed three times and visualized using a FUSION FX EDGE SPECTRA imaging system (VILBER, Collégien, France).

## ASFV SFV-RP Immunization

Five-week-old BALB/c mice were immunized three times via intramuscular injections of SFV-32 RPs, SFV-54 RPs, or PBS in the lateral thigh ( $n = 5/\text{group}$ ). Each mouse was immunized at 2-week intervals at the

same dose (0.2 mL) [The SFV-p32 RP titer in BHK-21 cells was  $10^{3.0}$  median tissue culture infectious dose ( $\text{TCID}_{50}$ )/0.1 mL, and the SFV-p54 RPs titer was  $10^{3.5}$   $\text{TCID}_{50}$ /0.1 mL]. Blood samples were collected from the canthus before the first immunization and 1 week after each



**FIGURE 2** | Expression of heterologous proteins in BHK-21 cells with SFVCs-RP infected. **(A)**  $\beta$ -galactosidase expression ( $\times 200$ ). **(B)** EGFP expression ( $\times 200$ ). **(C)** Cells were infected with SFVCs-EGFP RPs ( $\times 400$ ) for 24 h. CPE was evaluated.

immunization and stored at  $-80^{\circ}\text{C}$ . Mice were monitored and weighed daily.

### Enzyme-Linked Immunosorbent Assay

ELISA for detecting anti-p32 antibody IgG was performed with the serum samples described above using the ASFV p32 Antibodies Test Kit (sandwich ELISA) according to the manufacturer's instructions (Zhaoqing DaHuaNong Biology Medicine Co. Ltd., Zhaoqing, China). Microplates were coated with the ASFV p32 antigen and the serum samples were added, followed by HRP-labeled ASFV antigen. After washing the plate, the substrate was added, and the blue color was observed. The reaction was then terminated, giving a yellow color. The specific absorption peak at 450 nm was observed using a Spectrophotometer Multiskan Skyhigh (Thermo Scientific).

ELISA for detecting anti-p54 antibody IgG was performed using AsurDx™ ASFV p54 Antibody Test Kit (BioStone Animal Health, Dallas, TX, USA). The microplates were coated with the ASFV p54 antigen. The serum samples were added and the microplates were washed five times. An anti-ASF monoclonal antibody (mAb)-biotin conjugate and HRP-tagged streptavidin were added. After incubation for 30 min at  $37^{\circ}\text{C}$ , the plate was washed and the substrate was added. The color change was observed and the reaction was terminated, which caused another color change (blue to yellow). The specific absorption peak at 450 nm was measured using a Spectrophotometer Multiskan Skyhigh (Thermo Scientific). Three parallel experiments were performed for each sample.

### Enzyme-Linked Immune Absorbent Spot Assay

Mice were euthanized 2 weeks after the third immunization. Lymphocytes were aseptically isolated from the spleens for cellular immunity experiments. Interferon (IFN)- $\gamma$  and interleukin (IL)-4 expressions were measured using anti-mouse IFN- $\gamma$ - and anti-mouse IL-4-precoated ELISPOT kits

(Dakewe Biotech Co., Ltd., Shenzhen, China). The synthesized polypeptides were p32-1, DFNKV IRAHN FIQTI; p32-2, IRAHN FIQTI HGTP; p54, ENLRQ RNTYT HKDLE (Sangon Bioengineering Co., Ltd., Shanghai, China). Each group was assigned a positive control (10  $\mu\text{L}$  PHA with resuspended cells), a negative control (without stimulant but with resuspended cells), and a background negative control (only the culture medium). Mice immunized with SFV-p32 RP (with stimulant polypeptides p32-1 and p32-2, the final concentration is 60  $\mu\text{g}/\text{mL}$ ), and SFV-p54 RP (with stimulant polypeptide p54, the final concentration is 60  $\mu\text{g}/\text{mL}$ ) (five mice each) were used. An immunospot image analyzer from the Dakewe Biotech Co., Ltd., Shenzhen, China was used.

### Statistical Analysis

SPSS software (IBM SPSS Statistics for Windows, Version 20.0, IBM Corporation, Armonk, NY, USA) was used for statistical analysis. GraphPad Prism (version 9.2, GraphPad Software Inc., CA, USA) was used to analyze data. Significance levels were represented by *P*-values (ns, non-significant; \**P* < 0.05; \*\**P* < 0.01; \*\*\**P* < 0.001).


## RESULTS

### SFVCs-RP Production



The SFVCs-LacZ and SFVCs-EGFP RPs were successfully packaged. The blue color due to  $\beta$ -galactosidase was observed in 80–90% of cells, while few control cells were stained (**Figure 2A**). EGFP signal was detected using a fluorescence microscope, and more than half of the cells emitted green fluorescence (**Figure 2B**). Cells infected for 24 h with SFVCs-LacZ and SFVCs-EGFP RPs showed evident CPE (**Figure 2C**).

### SFV-RP Production

SFV-EGFP RPs were successfully packaged. BHK-21 cells were infected with SFV-EGFP RPs and SFVCs-EGFP RPs for 24 h,



**FIGURE 3** | Fluorescence signal intensity was compared after removing the viral capsid protein. BHK-21 cells were infected with SFVCs-EGFP and SFV-EGFP RPs for 24 h, and fluorescence signal intensities were compared ( $\times 200$ ).



**FIGURE 4** | IFA and viral growth of ASFV SFV-RPs. **(A)** IFA of ASFV SFV-RPs infected BHK-21 cells after 60 h ( $\times 400$ ). **(B)** Viral growth in ASFV SFV-RPs-infected BHK-21 cells.

and fluorescence signal intensities showed that the expression of EGFP in the former was not decreased compared with the latter (Figure 3), confirming that the SFV vector could efficiently express foreign genes.

### ASFV SFV-RP Production

EGFP in the pSFV-EGFP was replaced with ASFV antigens p32 and p54 and electroporated into BHK-21 cells. SFV-p32 and SFV-p54 RPs were successfully packaged.


### IFA of Infected BHK-21 Cells and Viral Growth of ASFV SFV-RPs

BHK-21 cells infected with SFV-p32 and SFV-p54 RPs for 60 h were detected using IFA. The results showed that infected cells exhibited green fluorescence and CPE at a suitable viral titer (Figure 4A).

The titers of SFV-p32 and SFV-p54 RPs in BHK-21 cells continuously increased from 24 to 36 h post-infection and peaked ( $10^{3.0}$  TCID<sub>50</sub>/0.1 mL,  $10^{3.5}$  TCID<sub>50</sub>/0.1 mL, respectively) 36 h post-infection (Figure 4B). The subsequent decrease was related to cell death.

### ASFV SFV-RP Western Blotting

To test heterologous p32 and p54 protein expression, cells were lysed and analyzed via western blotting. The bands at 30 kDa confirmed successful packaging of the SFV-p32 and SFV-p54 RPs and efficient infection of BHK-21 cells (Figure 5).



**FIGURE 5** | Western blotting of ASFV SFV-RPs. ASFV p32 expression in SFV-p32 RP-infected BHK-21 cells (Lane 1), ASFV p54 expression in SFV-p54 RP-infected BHK-21 cells (Lane 2).

### Mouse Weight Changes After Immunization


As shown in Figure 6, the weight of the immunized group decreased slightly and began to increase gradually again 1 week later, while finally there was no difference in the weight and state of the mice compared to the PBS group after completing the third immunization. The result indicated there had no significant effect on the growth of mice after immunization with ASFV SFV-RPs.

### ELISA Results

Mice were immunized three times with SFV-p32 and SFV-p54 RPs. Serum collected 1 week after the second and third immunizations were tested using ELISA. As shown in Figure 7, SFV-p32 RP-infected mice had higher antibody levels after the third immunization compared to the second immunization. Similarly, SFV-p54 RP-infected mice produced IgG antibodies after the second and third immunizations (Figure 8). The results indicated that three immunizations induced robust antibody production.

### ELISPOT Results


Two weeks after the third immunization, IFN- $\gamma$  and IL-4 expression were detected in lymphocytes using ELISPOT (Figure 9). The results showed that SFV-p32 and -p54 RP-injected mice had higher expression of both cytokines compared to the negative and background controls.



**FIGURE 6** | Comparative weight changes of immunized and non-immunized (PBS group) mice monitored for 35 days after the first immunization. ns, non-significant ( $P > 0.05$ ).

## DISCUSSION

Our results show that replication-defective viral particles can be produced by packaging SFV replicon RNA and first-generation helper RNA. The replication-defective viral particles were generated through amplification and recombination within  $\sim 24$  h, with viral production peaking after 36 h. Due to the deletion of key sections of the SFV genome, the



**FIGURE 7** | ELISA was used to detect anti-p32 IgG in each group. Cut off calculation: cut off = average value of negative control + 0.15. When the negative control value was  $\leq 0.05$ , it was calculated as 0.05. Positive judgment: samples with OD value  $\geq$  cut off were judged as positive. Negative determination: samples with OD value  $<$  cut off were judged as negative. SFV-p32 RP-immunized mice are numbered 1–5, which are consistent with the numbers assigned to mice detected by pre-immune serum, the second immunization and the third immunization in the FIGURE.


resulting replication-defective viral particles were only capable of performing one round of replication in infected cells (10). We initially produced SFVCs-EGFP RPs and found that over 50% of the cells expressed the EGFP.

Sjöberg et al. (14) indicated that the first 102 bases of the viral capsid gene play a role as a translational enhancer, and that SFV vectors that incorporate this RNA have improved heterologous protein synthesis. However, in the present study, the expression of the target protein was not reduced after the capsid-coding section was removed, which is different from previous results (14). We believe that the difference in results may be explained in several ways. Viral structural protein synthesis might be inhibited after SFV-infection so that the translation enhancer does not play a corresponding role. It was reported that the synthesis of viral structural proteins in extensive amounts upon Sindbis-infection engages  $<25\%$  of the synthesizing capacity used by the cell for steady state protein production and host protein synthesis stops almost completely 5 h after infection (14, 15). Furthermore, it was reported that a mildly acidic pH may lead to the hydrolysis of the SFV capsid (16). In addition, it is unclear which C' gene fragment enhances translation initiation and whether the effect results from an adaptation to the specific environment of SFV-infected cells (14). Further research is required to answer this question. In any case, to prevent the interference of the capsid protein on the specific expression of the target protein in subsequent animal experiments, the fused capsid protein should be removed. Therefore, we produced SFV-EGFP RPs. Finally, we produced ASFV SFV-RPs containing p32- and p54-encoding segments, which showed high levels of protein expression, implying that we were successful in establishing the SFV vector platform.

The serum contains numerous heterologous proteins, making it difficult to package RPs produced in cells for subsequent

animal experiments. There is a need for safe and reproducible methods for vaccine development. In this study, we first used 1% FBS in the medium after co-electroporation. Once the majority of the BHK-21 had adhered, they were cultured in a serum-free medium (growth kinetic parameters including cell density, viability, maximum growth density, and viral titer are not shown in this paper). We found that this approach was helpful for the RP production, and even subsequent animal experiments. Other researchers have reported the merits of using serum-free media in the production of vaccines and other biological products (2, 17). In the future, we will optimize our culture media to support viral proliferation better.

In the present study, mice were immunized with SFV RPs containing ASFV antigens, which resulted in the robust induction of humoral and cell-mediated immune responses. Almost all mice inoculated with SFV-p32 and SFV-p54 (RPs) produced IgGs against the expressed antigen after the third immunization. Moreover, the levels of ASFV p32-specific IgG and ASFV p54-specific IgG remained high 5 months post immunization. The ELISPOT results showed that all immunized mice produced IFN- $\gamma$  and IL-4, which indicates that the SFV-based vector vaccine induced a robust cell-mediated response. Although one mouse did not produce IgG antibodies after the third round of immunization with SFV-p54 RPs, it still exhibited a cellular immune response [Figure 9 (c-3, d-3)]. This might be because ASFV SFV-RPs are more effective at inducing cellular rather than humoral immune responses. ILs mediate T and B lymphocyte activation, proliferation, and differentiation during the activation and regulation of the cellular immune response. Our data demonstrate that SFV-p32 and SFV-p54 RPs also elicited high IL-4 expression in mice, an important mediator of the Th1-type response. There is evidence that cytotoxic CD8<sup>+</sup> T lymphocytes play an important role in protection against ASFV,




**FIGURE 8** | ELISA was used to detect anti-p54 IgG and calculate the inhibition percentage (IP) in each group. **(A)** ELISA results for each group. The mean OD of the negative control was  $\geq 0.5$ . **(B)** Calculation of the IP of all samples in each group. IP of all samples was calculated as follows:  $IP (\%) = (1 - OD_{450} \text{ test sample} / \text{Mean } OD_{450} \text{ negative control}) * 100$ . The IP (%) of the positive control was  $\geq 55\%$ . Serum/plasma samples were assessed as follows:  $IP \leq 40\%$  (negative), ASF antibodies were absent in the test samples;  $IP \geq 50\%$  ASF antibodies were present in the test samples;  $50\% > IP \geq 40\%$  doubtful, rechecked in 1–2 weeks. SFV-p54 RP-immunized mice are numbered 1–5, which are consistent with the numbers assigned to mice detected by pre-immune serum, the second immunization and the third immunization in the FIGURE.

and cell-mediated response is critical for clearing the infection (5, 18).

The SFV-based vector vaccine is a defective single-cycle virus generated by substituting the structural protein-encoding genes from the SFV genome with the ASFV antigens and then electroporation this vector RNA into BHK-21 cells together with helper RNA. Numerous studies have documented the development of DNA and DNA replicon vaccines (19, 20). Although it was reported that DNA vaccines elicit humoral and cellular immune responses, they failed to protect animals from a virulent ASFV challenge (21). Moreover, RNA replicon vaccines based on the alphavirus replicon system have been evaluated in

chickens, pigs, and humans, and provided complete protection against lethal challenges (4, 12). It was reported that young pigs exhibited a protective humoral immune response following influenza hemagglutinin (HA) mRNA RPs vaccination (4, 22). In addition, robust humoral and cellular responses were elicited in mice, rabbits, and rhesus macaques which were given HA and neuraminidase (NA) mRNA RPs vaccines (4, 23). Though Murgia et al. (5) already described ASFV RP-30 and RP-54 using VEEV, which were tested for expression levels in Vero cells and immunogenicity in swine, there is a lack of relevant research on the cellular immune response. The present paper is different from Murgia et al. (5) as it focused on the design and optimization



**FIGURE 9** | ELISPOT was used to detect IFN- $\gamma$  and IL-4 expression in SFV-p32 and -p54 RP-immunized mice. **(A)** IFN- $\gamma$  spots of SFV-p32 RP-immunized mice. **(B)** IL-4 spots of SFV-p32 RP-immunized mice. **(C)** IFN- $\gamma$  spots of SFV-p54 RP-immunized mice. **(D)** IL-4 spots of SFV-p54 RP-immunized mice. For all four panels, 1–5 are the experimental groups and 6, 7, and 8 are the background control, negative control, and positive control, respectively. The numbers indicate the number of spots.

of replicon particles and the study of cellular immune response, which we believe will be useful for further immunogenicity and protection experiments in pig models.

Recent strategies have involved the modification of the alphaviral vector 5' terminal region to specifically identify cancer cells and induce their death (24). Although such research is still in its early stages, it holds promise for cancer treatment. Modified viruses can also be used in the research and development of cancer vaccines given the vast number of target proteins (25). Therapeutic alphaviral vector-derived cancer vaccines activate humoral and cellular immunity in the treatment of melanoma, as well as breast, mast cell, and prostate cancers (26). Despite the promise of viral vector vaccines, certain challenges remain

(27). In this study, there was no recovery of infectious virus by blind passaging, IFA, or plaque assay; furthermore, no clinical symptoms of infection were observed in all inoculated animals, which indicate that use of this SFV vector system is valuable to express high levels of recombinant ASFV proteins and study the immunogenicity in pigs. It was reported that SFV-helper1 system is prone to generating the wild-type virus by recombination, however, using the pSFV-helper2 and the split helper can increase the safety of the system by decreasing recombination events (28–31). Therefore, in the next stage, we will use two other systems and compare them with the SFV-helper1 system to improve applications in clinical trials. In addition, there is still room for further alphaviral vector optimization, such as replacing the SP6

promoter with a cytomegalovirus (CMV) promoter, eliminating the *in vitro* transcription step, and minimizing RNA degradation, all of which could reduce costs and save effort.

## CONCLUSION

We established an SFV vector platform to produce replication-defective viral particles containing ASFV antigens, which induced effective antigen-specific humoral and cellular immune responses. The resulting RPs exhibited excellent biosafety and elicited a robust immune response due to the replication-defective nature of the viral particles and the use of a self-replicating RNA. However, in this study, we have not included any pig models. Nevertheless, we still believe that this paper will be useful for future immunogenicity research in pig models, providing a reference for the production of RNA replicon vaccines to elicit robust humoral and cellular immune responses.

## DATA AVAILABILITY STATEMENT

The datasets presented in this study can be found in online repositories. The names of the repository/repositories and accession number(s) can be found in the article/supplementary material.

## REFERENCES

- Wolff JA, Malone RW, Williams P. Direct gene transfer into mouse muscle *in vivo*. *Science*. (1990) 247:1465–8. doi: 10.1126/science.1690918
- Suárez-Patiño SF, Bernardino TC, Núñez EGF, Astray RM, Pereira CA, Soares HR, et al. Semliki Forest Virus replicon particles production in serum-free medium BHK-21 cell cultures and their use to express different proteins. *Cytotechnology*. (2019) 71:949–62. doi: 10.1007/s10616-019-00337-y
- Mogler MA, Kamrud KI. RNA-based viral vectors. *Exp Rev Vaccines*. (2015) 14:283–312. doi: 10.1586/14760584.2015.979798
- Vander Veen R, Hank Harris DL, Kamrud KI. Alphavirus replicon vaccines. *Anim Health Res Rev*. (2012) 13:1–9. doi: 10.1017/S146625231200011
- Murgia MV, Mogler M, Certoma A, Green D, Monaghan P, Williams DT, et al. Evaluation of African swine fever (ASF) vaccine strategy incorporating priming with an alphavirus-expressed antigen followed by boosting with attenuated ASF virus. *Arch Virol*. (2019) 164:359–70. doi: 10.1007/s00705-018-4071-8
- Gomez-Puertas P, Rodriguez F, Oviedo JM, Brun A, Alonso C, Escribano JM, et al. The African swine fever virus proteins p54 and p30 are involved in two distinct steps of virus attachment and both contribute to the antibody-mediated protective immune response. *Virology*. (1998) 243:461–71. doi: 10.1006/viro.1998.9068
- Rodriguez F, Ley V, Gomez-Puertas P. The structural protein p54 is essential for African swine fever virus viability. *Virus Res*. (1996) 40:161–7. doi: 10.1016/0168-1702(95)01268-0
- García-Mayoral MF, Rodríguez-Crespo I, Bruix M. Structural models of DYNLLI with interacting partners: African swine fever virus protein p54 and postsynaptic scaffolding protein gephyrin. *FEBS Lett*. (2011) 585:53–7. doi: 10.1016/j.febslet.2010.11.027
- Hernaiz B, Diaz-Gil G, Garcia-Gallo M. The African swine fever virus dynein-binding protein p54 induces infected cell apoptosis. *FEBS Lett*. (2004) 569:224–8. doi: 10.1016/j.febslet.2004.06.001
- Liljeström P, Garoff H. A new generation of animal cell expression vectors based on the Semliki Forest Virus replicon. *Biotechnology*. (1991) 9:1356–61. doi: 10.1038/nbt1291-1356

## ETHICS STATEMENT

The animal study was reviewed and approved by Animal Welfare and Ethical Censor Committee.

## AUTHOR CONTRIBUTIONS

RC conceived the idea. NF did most of the experimental work and wrote the first draft of the manuscript. BY, HZ, and TX made the statistical analyses and did parts of the experimental work. YL assembled the data. HL and AZ advised in the process of manuscript writing. All authors contributed to the article and approved the submitted version.

## FUNDING

This study was supported by the Technology Planning Project of Guangdong Province (grant number 2019B020211004).

## ACKNOWLEDGMENTS

We thank YL for providing the anti-ASFV serum. We thank Guangming Ma and Weifeng Zhong for providing the ASFV antibody test kit.

- Berglund P, Sjöberg M, Garoff H, Atkins GJ, Sheahan BJ, Liljeström P, et al. Semliki Forest expression system: production of conditionally infectious recombinant particles. *Bio Technol*. (1993) 11:916–20. doi: 10.1038/nbt0893-916
- Schultz-Cherry S, Dybing JK, Davis NL, Williamson C, Suarez DL, Johnston R. Influenza virus (A/HK/156/97) hemagglutinin expressed by an alphavirus replicon system protects chickens against lethal infection with Hong Kong-origin H5N1 viruses. *Virology*. (2000) 278:55–9. doi: 10.1006/viro.2000.0635
- Pushko P, Geisbert J, Parker M, Jahrling P, Smith J. Individual and bivalent vaccines based on alphavirus replicons protect guinea pigs against infection with Lassa and Ebola viruses. *J Virol*. (2001) 75:11677–85. doi: 10.1128/JVI.75.23.11677-11685.2001
- Sjöberg EM, Suomalainen M, Garoff H. A significantly improved Semliki Forest virus expression system based on translation enhancer segments from the viral capsid gene. *Biotechnology*. (1994) 12:1127–31. doi: 10.1038/nbt1194-1127
- Garry RF. Sindbis virus-induced inhibition of protein synthesis is partially reversed by medium containing an elevated potassium concentration. *J Gen Virol*. (1994) 75 (Pt 2):411–5. doi: 10.1099/0022-1317-75-2-411
- Schlegel A, Schaller J, Jentsch P, Kempf C. Semliki forest virus core protein fragmentation: its possible role in nucleocapsid disassembly. *Biosci Rep*. (1993) 13:333–47. doi: 10.1007/BF01150478
- Fontana D, Kratje R, Etcheverrigaray M, Prieto C. Immunogenic virus-like particles continuously expressed in mammalian cells as a veterinary rabies vaccine candidate. *Vaccine*. (2015) 33:4238–46. doi: 10.1016/j.vaccine.2015.03.088
- Takamatsu HH, Denyer MS, Lacasta A, Stirling CM, Argilaguat JM, Netherton CL, et al. Cellular immunity in ASFV responses. *Virus Res*. (2013) 173:110–21. doi: 10.1016/j.virusres.2012.11.009
- Kohno A, Emi N, Kasai M, Tanimoto M, Saito H. Semliki Forest virus-based DNA expression vector: transient protein production followed by cell death. *Gene Ther*. (1998) 5:415–8. doi: 10.1038/sj.gt.3300589
- Ljungberg K, Whitmore AC, Fluet ME, Moran TP, Shabman RS, Collier ML, et al. Increased immunogenicity of a DNA-launched Venezuelan equine encephalitis virus-based replicon DNA vaccine. *J Virol*. (2007) 81:13412–23. doi: 10.1128/JVI.01799-07

21. Argilaguet JM, Pérez-Martín E, Nofrarias M, Gallardo C, Accensi F, Lacasta A, et al. DNA vaccination partially protects against African swine fever virus lethal challenge in the absence of antibodies. *PLoS ONE*. (2012) 7:942. doi: 10.1371/journal.pone.0040942
22. Erdman MM, Kamrud KI, Harris DL, Smith J. Alphavirus replicon particle vaccines developed for use in humans induce high levels of antibodies to influenza virus hemagglutinin in swine: proof of concept. *Vaccine*. (2010) 28:594–6. doi: 10.1016/j.vaccine.2009.10.015
23. Hubby B, Talarico T, Maughan M, Reap EA, Berglund P, Kamrud KI, et al. Development and preclinical evaluation of an alphavirus replicon vaccine for influenza. *Vaccine*. (2007) 25:8180–9. doi: 10.1016/j.vaccine.2007.09.038
24. Ramachandran M, Yu D, Dyczynski M, Baskaran S, Zhang L, Lulla A, et al. Safe and effective treatment of experimental neuroblastoma and glioblastoma using systemically delivered triple micro RNA-detargeted oncolytic semliki forest virus. *Clin Cancer Res*. (2017) 23:1519–30. doi: 10.1158/1078-0432.CCR-16-0925
25. Schirmacher V, Forg P, Dalemans W, Chlichlia K, Zeng Y, Fournier P, et al. Intra-pinna anti-tumor vaccination with self-replicating infectious RNA or with DNA encoding a model tumor antigen and a cytokine. *Gene Ther*. (2000) 7:1137–47. doi: 10.1038/sj.gt.3301220
26. Ketola A, Hinkkanen A, Yongabi F, Furu P, Määttä AM, Liimatainen T, et al. Oncolytic Semliki forest virus vector as a novel candidate against unresectable osteosarcoma. *Cancer Res*. (2008) 68:8342–50. doi: 10.1158/0008-5472.CAN-08-0251
27. Smyth JW, Fleeton MN, Sheahan BJ, Atkins GJ. Treatment of rapidly growing K-BALB and CT26 mouse tumours using Semliki Forest virus and its derived vector. *Gene Ther*. (2005) 12:147–59. doi: 10.1038/sj.gt.3302390
28. Tubulekas I, Liljeström P. Suppressors of cleavage-site mutations in the p62 envelope protein of Semliki Forest virus reveal dynamics in spike structure function. *J Virol*. (1998) 72:2825–31. doi: 10.1128/JVI.72.4.2825-2831.1998
29. Smerdou C, Liljeström P. Two-helper RNA system for production of recombinant Semliki forest virus particles. *J Virol*. (1999) 73:1092–8. doi: 10.1128/JVI.73.2.1092-1098.1999
30. Rayner JO, Dryga SA, Kamrud KI. Alphavirus vectors and vaccination. *Rev Med Virol*. (2002) 12:279–96. doi: 10.1002/rmv.360
31. Ketola A, Schlesinger S, Wahlfors J. Properties of Sindbis virus vectors produced with a chimeric split helper system. *Int J Mol Med*. (2005) 15:999–1003. doi: 10.3892/ijmm.15.6.999

**Conflict of Interest:** NF, BY, HZ, and RC were employed by Zhaoqing Institute of Biotechnology Co. Ltd. YL and RC were employed by Zhaoqing DaHuaNong Biology Medicine Co. Ltd.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Fang, Yang, Xu, Li, Li, Zheng, Zhang and Chen. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.